



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2018.1D

Diagnostik und Therapie primärer und metastasierter Mammakarzinome

Komplementäre Therapie „Survivorship“

FORSCHEN
LEHREN
HEILEN



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2018.1D

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

Komplementäre Therapien Hormontherapie „Survivorship“ (Rezidiv-Prävention)

▪ Versionen 2002–2017:

Albert / Bauerfeind / Blohmer / Fersis /
Friedrich / Gerber / Göhring / Hanf /
Janni / Kümmel / Lück / von Minckwitz / Oberhoff /
Scharl / Schmidt / Schütz / Thomssen

▪ Version 2018:

Kümmel / Hanf

Screened Data Sources:


Pubmed 2012 - 01/2018

ASCO 2015 – 2016

SABCS 2015 – 2017

EBCC 2015 – 2016

Cochrane library: summary Jan. 2018:



Komplementäre Verfahren werden parallel zur konventionellen Therapie angewendet und unterscheiden sich von alternativen Verfahren dadurch, dass sie den Wert der konventionellen Verfahren nicht in Frage stellen, sondern sich als Ergänzung verstehen

Onkolleitlinienprogramm

1. Witt CM et al.. A Comprehensive Definition for Integrative Oncology. J Natl Cancer Inst Monogr (2017) 2017(52): Igx012

“Integrative oncology is a patient-centered, evidence-informed field of cancer care that utilizes mind and body practices, natural products, and/or lifestyle modifications from different traditions alongside conventional cancer treatments. Integrative oncology aims to optimize health, quality of life, and clinical outcomes across the cancer care continuum and to empower people to prevent cancer and become active participants before, during, and beyond cancer treatment.”



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2018.1D

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

Allgemein

	Oxford		
	LoE	GR	AGO
▪ CAM anstelle lokoregionärer Interventionen	2b	B	--
▪ CAM anstelle systemischer Therapie	2b	B	--
• Patienten sollten nach ihrer Nutzung von komplementären und alternativen Therapien befragt werden.			
▪ <i>Unter Systemtherapie:</i> Besondere Beachtung gilt möglichen Medikamenteninteraktionen			

1. Saquib J, Parker BA, Natarajan L, et al. Prognosis following the use of complementary and alternative medicine in women diagnosed with breast cancer. Complement Ther Med. 2012 Oct;20(5):283-90. doi: 10.1016/j.ctim.2012.04.002. Epub 2012 Apr 27.
2. Guha N, Kwan ML, Quesenberry CP, et al: Soy isoflavones and risk of cancer recurrence in a cohort of breast cancer survivors: the Life After Cancer Epidemiology study. Breast Cancer Res Treat. 2009;118(2):395–405, pmid:19221874.
3. Johnson SB et al.. Use of Alternative Medicine for Cancer and Its Impact on Survival. JNCI J Natl Cancer Inst (2018) 110(1): djx145.
4. Fremd C et al.. Use of complementary and integrative medicine among German breast cancer patients: predictors and implications for patient care within the PRAEGNANT study network. Arch Gynecol Obstet. 2017 May;295(5):1239-1245. doi: 10.1007/s00404-017-4348-2. Epub 2017 Mar 22.
5. Samuels N et al.. Unmonitored use of herbal medicine by patients with breast cancer: reframing expectations. J Cancer Res Clin Oncol (2017) 143:2267–2273
6. Smith PJ et al.. Complementary and alternative medicine use by patients receiving curative-intent chemotherapy. Asia-Pacific Journal of Clinical Oncology 2016; 12: 265–274
7. Greenlee H et al.. Association Between Complementary and Alternative Medicine Use and Breast Cancer Chemotherapy Initiation: The Breast Cancer Quality of Care (BQUAL) Study. JAMA Oncol. 2016 Sep 1;2(9):1170-6. doi: 10.1001/jamaoncol.2016.0685



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2018.1D

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

Komplementäre Therapien prä- und postoperativ

Oxford
LoE GR AGO

Präoperativ

- Hypnose (reduziert Ängste, Schmerz, Übelkeit)

1b B +

Postoperativ

Akupunktur

- bei Schmerzen, Ängstlichkeit
- bei Übelkeit, Erbrechen

1b B +/-
2b B +

Massage Therapie (bei Schmerzen)

2b C +/-

Frühzeitige postoperative Bewegungstherapie

beugt Dysfunktion der oberen Extremität vor
CAVE: vermehrt Wundsekret

1a A +

Prophylaktische Lymphdrainage

1b B -

Hypnosis

- Cramer H, Lauche R, Paul A, et al: Hypnosis in Breast Cancer Care: A Systematic Review of Randomized Controlled Trials. Integr Cancer Ther. 2015 Jan;14(1):5-15. Epub 2014 Sep 18.
- Montgomery GH, David D, Kangas M, et al. (2014) Randomized Controlled Trial of a Cognitive-Behavioral Therapy Plus Hypnosis Intervention to Control Fatigue in Patients Undergoing Radiotherapy for Breast Cancer. JCO DOI 10.1200/JCO.2013.49.3437
- Montgomery GH, Bovbjerg DH, Schnur JB et al. (2007): A randomized clinical trial of a brief hypnosis intervention to control side effects in breast surgery patients. J Nat Cancer Inst; 99:1304–1312.
- Schnur JB, Bovbjerg DH, David D et al. (2008): Hypnosis decreases presurgical distress in excisional breast biopsy patients. Anesth Analg , 106(2):440-4
- Montgomery GH, Schnur JB, Kravits K. Hypnosis for cancer care: Over 200 years young. CA Cancer J Clin. 2012 Nov 20. doi: 10.3322/caac.21165.

Acupuncture and Postoperative Nausea and Vomiting / Pain

- Mallory MJ et al.: Acupuncture in the postoperative setting for breast cancer patients: a feasibility study. Am J Chin Med. 2015;43(1):45-56.
- Chao LF et al.: The efficacy of acupoint stimulation for the management of therapy-related adverse events in patients with breast cancer: a systematic review. Breast Cancer Res Treat (2009) 118:255–267.
- Quinlan-Woodward J, Gode A, Dusek JA: Assessing the Impact of Acupuncture on Pain, Nausea, Anxiety, and Coping in Women Undergoing a Mastectomy. Oncol

- Nurs Forum. 2016 Nov 1;43(6):725-732.
4. Giron PS, Haddad CA, Lopes de Almeida: Effectiveness of acupuncture in rehabilitation of physical and functional disorders of women undergoing breast cancer surgery. Support Care Cancer. 2016 Jun;24(6):2491-6.
 5. Chiu HY, Hsieh YJ, Tsai PS. Systematic review and meta-analysis of acupuncture to reduce cancer-related pain. Eur J Cancer Care (Engl). 2017 Mar;26(2). doi: 10.1111/ecc.12457. Epub 2016 Feb 7

Massage Therapy

1. Lee SH, Kim JY, Yeo S et al: Meta-Analysis of Massage Therapy on Cancer Pain. Integr Cancer Ther. 2015 Jul;14(4):297-304.
2. Pan YQ, Yang KH, Wang YL, et al: Massage interventions and treatment-related side effects of breast cancer: a systematic review and meta-analysis. Int J Clin Oncol. 2014 Oct;19(5):829-41.

Postoperative exercise

1. De Groef A, Van Kampen M, Dieltjens E, et al. Effectiveness of postoperative physical therapy for upper-limb impairments after breast cancer treatment: a systematic review. Arch Phys Med Rehabil. 2015 Jun;96(6):1140-53. doi: 10.1016/j.apmr.2015.01.006. Epub 2015 Jan 13. Review.
2. McNeely ML, Campbell K, Ospina M et al.: Exercise interventions for upper-limb dysfunction due to breast cancer treatment. Cochrane Database of Systematic Reviews 2010, Issue 6. Art. No.: CD005211. DOI: 10.1002/14651858.CD005211.pub2.
3. Cavanaugh KM.: Effects of Early Exercise on the Development of Lymphedema in Patients With Breast Cancer Treated With Axillary Lymph Node Dissection. J Oncol Pract. 2011 March; 7(2): 89–93.
4. Anderson RT, Kimmick GG, McCoy TP, et al. A randomized trial of exercise on well-being and function following breast cancer surgery: the RESTORE trial. J Cancer Surv 2012;6(2):172-81

Prophylactic lymph drainage

1. Devoogdt N, Christiaens MR, Geraerts I, et al: Effect of manual lymph drainage in addition to guidelines and exercise therapy on arm lymphoedema related to breast cancer: randomised controlled trial. BMJ 2011;343:d5326 doi: 10.1136/bmj.d5326
2. Li L, Yuan L, Chen X: Current Treatments for Breast Cancer-Related Lymphoedema: A Systematic Review. Asian Pac J Cancer Prev. 2016 Nov 1;17(11):4875-4883.



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2018.1D

www.ago-online.de
**FORSCHEN
LEHREN
HEILEN**

Komplementäre Therapien

Behandlungsphase – Einfluss auf Toxizität I

Bei laufender onkologischer Standardtherapie:
CAVE: Medikamenten-Interaktionen!

	Oxford		
	LoE	GR	AGO
▪ Mistellektine (Viscum album) zur Reduktion von therapieassoziierten Nebenwirkungen	1a	B	+/-
▪ Thymuspeptide verringern Risiko schwerer Infektionen	2a	B	+/-
▪ Ginseng verringert Krebs-assoziierte Fatigue; inhibiert u.a. Cytochrom P Enzyme, z.B. CYP3A4)	2b	C	-
▪ Ganoderma Lucidum Besserung der Fatigue; Cave: inhibiert P Enzyme, z.B. CYP3A4)	2b	C	-
▪ L-Carnitin Prävention der Toxizität, Verbesserung peripherer Neuropathie Verbesserung der Fatigue	1b	B	--
▪ Curcumin um Radiodermatitis zu vermindern	1b	B	+/-
▪ Ingwer komplementär zu Leitlinien-gerechter Medikation gegen Chemotherapie induzierte Übelkeit/Erbrechen; Cave: Wechselwirkungen	1b	C	+/-

General

- Li Y, Wang J, Lin F: A Methodology for Cancer Therapeutics by Systems Pharmacology-Based Analysis: A Case Study on Breast Cancer-Related Traditional Chinese Medicines. *PLoS One*. 2017 Jan 9;12(1):e0169363.
- Farahmand L, Darvishi B, Majidzadeh-A K: Naturally occurring compounds acting as potent anti-metastatic agents and their suppressing effects on Hedgehog and WNT/β-catenin signalling pathways. *Cell Prolif*. 2017 Feb;50(1). doi: 10.1111/cpr.12299.
- Cramer H, Lauche R, Klose P: Yoga for improving health-related quality of life, mental health and cancer-related symptoms in women diagnosed with breast cancer. *Cochrane Database Syst Rev*. 2017 Jan 3;1:CD010802.
- Neuhouser ML, Smith AW, George SM: Use of complementary and alternative medicine and breast cancer survival in the Health, Eating, Activity, and Lifestyle Study. *Breast Cancer Res Treat*. 2016 Dec;160(3):539-546.

Mistletoe

- Marvibaigi M, Supriyanto E, Amini N, et al: Preclinical and clinical effects of mistletoe against breast cancer. *BioMed research international* 2014;2014:785479.
- Shneerson C, Taskila T, Gale N, et al: The effect of complementary and alternative medicine on the quality of life of cancer survivors: A systematic review and meta-analyses. *Complementary therapies in medicine* 2013;21:417-429.
- Troger W, Zdrale Z, Stankovic N, et al: Five-year follow-up of patients with early stage breast cancer after a randomized study comparing additional treatment

- with viscum album (L.) extract to chemotherapy alone. Breast cancer : basic and clinical research 2012;6:173-180.
4. Troger W, Zdrale Z, Tisma N, et al: Additional therapy with a mistletoe product during adjuvant chemotherapy of breast cancer patients improves quality of life: An open randomized clinical pilot trial. Evidence-based complementary and alternative medicine : eCAM 2014;2014:430518.
 5. Weissenstein U, Kunz M, Urech K, et al: Interaction of standardized mistletoe (viscum album) extracts with chemotherapeutic drugs regarding cytostatic and cytotoxic effects in vitro. BMC complementary and alternative medicine 2014;14:6.

Thymus

1. Wolf E, Milazzo S, Boehm K, et al. Thymic peptides for treatment of cancer patients. Cochrane Database of Systematic Reviews 2012, Issue 2. Art. No.: CD003993. DOI: 10.1002/14651858.CD003993.pub3.

Ginseng, Ganoderma lucidum

1. Leggett S1, Koczwara B, Miller M. The impact of complementary and alternative medicines on cancer symptoms, treatment side effects, quality of life, and survival in women with breast cancer--a systematic review. Nutr Cancer. 2015;67(3):373-91.
2. Bao, P. P., W. Lu, Y. Cui, Y et al(2012). "Ginseng and Ganoderma lucidum use after breast cancer diagnosis and quality of life: a report from the Shanghai Breast Cancer Survival Study." PLoS One 7(6): e39343.
3. Jin X, Ruiz Beguerie J, Sze Daniel M-y et al: Ganoderma lucidum (reishi mushroom) for cancer treatment. Cochrane Database of Systematic Reviews 2012
4. JL Ryan et al.: Ginger (Zingiberofficinale) reduces acute chemotherapy-induced nausea: a URCC CCOP study of 576 patients. Support Care Cancer 2012;20(7):1479-89
5. Karimi N, Roshan VD: Change in adiponectin and oxidative stress after modifiable lifestyle interventions in breast cancer cases. Asian Pacific journal of cancer prevention : APJCP 2013;14:2845-2850.

L-Carnitine

1. Hershman DL, Unger JM, Crew K et al: Randomized double-blind placebo-controlled trial of acetyl-L-carnitine for the prevention of taxane-induced neuropathy in women undergoing adjuvant breast cancer therapy. J Clin Oncol. 2013 Jul 10;31(20):2627-33
2. Cruciani RA, Zhang JJ, Manola J et al. L-carnitine supplementation for the management of fatigue in patients with cancer: an eastern cooperative oncology group phase III, randomized, double-blind, placebo-controlled trial. J Clin Oncol. 2012 Nov 1;30(31):3864-9

Curcumin

1. Ryan, J. L., C. E. Heckler, M. Ling, et al (2013). "Curcumin for radiation dermatitis: a randomized, double-blind, placebo-controlled clinical trial of thirty breast

cancer patients." Radiat Res 180(1): 34-43.

2. Kumar P, Kadakol A, Shasthrula P, et al: Curcumin as an adjuvant to breast cancer treatment. Anti-cancer agents in medicinal chemistry 2015
3. Bandyopadhyay D: Farmer to pharmacist: Curcumin as an anti-invasive and antimetastatic agent for the treatment of cancer. Frontiers in chemistry 2014;2:113.

Ingwer

1. Thamlikitkul L, Srimuninnimit V, Akewanlop C, et al. Efficacy of ginger for prophylaxis of chemotherapy-induced nausea and vomiting in breast cancer patients receiving adriamycin-cyclophosphamide regimen: a randomized, double-blind, placebo-controlled, crossover study. Support Care Cancer. 2017 Feb;25(2):459-464. doi: 10.1007/s00520-016-3423-8. Epub 2016 Oct 6.
2. Sanaati F, Najafi S, Kashaninia Z, et al. Effect of Ginger and Chamomile on Nausea and Vomiting Caused by Chemotherapy in Iranian Women with Breast Cancer. Asian Pac J Cancer Prev. 2016;17(8):4125-9.



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2018.1D

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

Komplementäre Therapien Behandlungsphase – Einfluss auf Toxizität II

	Oxford		
	LoE	GR	AGO
▪ Antioxidanzien (Suppl.)	1b	B	-
▪ verschied. Antioxidative Extrakte zur Minderung anthrazyklinbedingter Cardiotoxizität	2b	B	+/-
▪ Hochdosiert Vitamin C *	1b	C	-
▪ Vitamin E	2b	D	-
▪ Selen zur Linderung der Nebenwirkungen	1b	B	-
▪ Co-Enzym Q 10 (Fatigue, Lebensqualität)	1b	B	-
▪ Proteolytische Enzyme (gegen Chemotherapie-induzierte Toxizität)	3b	B	-
▪ Verbesserung der Wundheilung durch Chinesische Medizin	1b	B	-*inf
▪ Sauerstoff- und Ozon-Therapie	5	D	--

Infusion in Dtd. nicht geprüfter Substanzen

* Studienteilnahme empfohlen

General

1. Zhu L, Li L, Li Y: Chinese Herbal Medicine as an Adjunctive Therapy for Breast Cancer: A Systematic Review and Meta-Analysis. Evid Based Complement Alternat Med. 2016;2016:9469276. doi: 10.1155/2016/9469276.
2. McPherson L, Cochrane S, Zhu X: Current Usage of Traditional Chinese Medicine in the Management of Breast Cancer: A Practitioner's Perspective. Integr Cancer Ther. 2016 Sep;15(3):335-42. doi: 10.1177/1534735415607656.

Antioxidant supplements

1. Greenlee H, Hershman DL, Jacobson JS: Use of antioxidant supplements during breast cancer treatment: a comprehensive review. Breast Cancer Res Treat. 2009 Jun;115(3):437-52.
2. S.-K. Myung, Y. Kim, W. Ju et al.: Effects of antioxidant supplements on cancer prevention: meta-analysis of randomized controlled trials. Annals of Oncology 21: 166–179, 2010.
3. Harvie M: Nutritional supplements and cancer: Potential benefits and proven harms. American Society of Clinical Oncology educational book / ASCO American Society of Clinical Oncology Meeting 2014:e478-486.
4. van Dalen EC, Caron HN, Dickinson HO, et al: Cardioprotective interventions for cancer patients receiving anthracyclines. Cochrane Database Syst Rev 2011:Cd003917.
5. The Protective Role of Phenolic Compounds Against Doxorubicin-induced Cardiotoxicity: A Comprehensive Review. Razavi-Azarkhiavi K, Iranshahy M, Sahebkar A, et al. Nutr Cancer. 2016 Aug-Sep;68(6):892-917. doi:

- 10.1080/01635581.2016.1187280. Epub 2016 Jun 24. Review
6. Drugs R D. 2012 Jun 1;12(2):101-6. doi: 10.2165/11632530-000000000-00000. Protective effects of salidroside on epirubicin-induced early left ventricular regional systolic dysfunction in patients with breast cancer.
 7. Zhang H1, Shen WS, Gao CH, et al.

Vitamin C

1. Ohno S, Ohno Y, Suzuki N et al.: High-dose Vitamin C (Ascorbic Acid) Therapy in the Treatment of Patients with Advanced Cancer. Anticancer Res. 2009 Mar; 29(3):809-15. Review.
2. Heaney M, Gardner J, Karasavvas N et al.: Vitamin C antagonizes the cytotoxic effects of antineoplastic drugs. Cancer Res. 2008 Oct 1;68(19):8031-8.
3. PLoS One. 2015 Apr 7;10(4):e0120228. doi: 10.1371/journal.pone.0120228. eCollection 2015. High-dose intravenous vitamin C combined with cytotoxic chemotherapy in patients with advanced cancer: a phase I-II clinical trial. Hoffer LJ1, Robitaille L2, Zakarian R3, et al..
4. Sci Transl Med. 2014 Feb 5;6(222):222ra18. doi: 10.1126/scitranslmed.3007154. High-dose parenteral ascorbate enhanced chemosensitivity of ovarian cancer and reduced toxicity of chemotherapy. Ma Y1, Chapman J, Levine M, et al.

Selen

1. Dennert G, Horneber M. Selenium for alleviating the side effects of chemotherapy, radiotherapy and surgery in cancer patients. Cochrane Database of Systematic Reviews 2010, Issue 11. Art. No.: CD005037. DOI: 10.1002/14651858.CD005037.pub2.
2. Greenlee H, Hershman DL, Jacobson JS: Use of antioxidant supplements during breast cancer treatment: a comprehensive review. Breast Cancer Res Treat. 2009 Jun;115(3):437-52.

Coenzym Q10

1. Lesser GJ, Case D, Stark N, et al. A randomized, double-blind, placebo-controlled study of oral coenzyme Q10 to relieve self-reported treatment-related fatigue in newly diagnosed patients with breast cancer. J Support Oncol 2013;11(1):31-42
2. Lockwood K et al.: Partial and complete regression of breast cancer in patients in relation to dosage of coenzyme Q10. Biochem Biophys Res Comm 1994;199:1504-8.
3. Lockwood K et al.: Progress on therapy of breast cancer with vitamin Q10 and the regression of metastasis. Biochem Biophys Res Comm 1995;212:172-7.
4. Lund EL, Quistorff B, Spang-Thomsen M et al.: Effect of radiation therapy on small-cell lung cancer is reduced by ubiquinone intake. Folia Microbiol 1998;4:505-6.
5. Ann Oncol. 2017 Mar 1;28(3):628-633. doi: 10.1093/annonc/mdw671. Interventions for preventing cardiomyopathy due to anthracyclines: a Bayesian network meta-analysis. Abdel-Qadir H1,2,3,4, Ong G4, Fazelzad R5, Amir E1,5,6, Lee DS1,2,6,7, Thavendiranathan P6,7, Tomlinson G1,6.

Proteolytic enzymes and toxicity of chemotherapy

1. Petru U, Stranz B, Petru C: Effects of proteolytic enzyme therapy with Wobe Mугос against chemotherapy-induced toxicity in breast cancer patients - results of a pilot study Wien Med Wochenschr. 2010 Nov;160(19-20):513-6.

Bromelain

1. Hidaka M, Nagata M, Kawano Y, et al.: Inhibitory effects of fruit juices on cytochrome P450 2C9 activity in vitro. Biosci Biotechnol Biochem. Feb 2008;72(2):406-411.

Chinese herbal medicine and wound healing

1. Chen J, Lv Q, Yu M et al.: Randomized clinical trial of Chinese herbal medications to reduce wound complications after mastectomy for breast carcinoma. Br J Surg. 2010 Dec;97(12):1798-804



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2018.1D

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

Komplementäre Therapien unter (Chemo)therapie Behandlung von Nebenwirkungen

	Oxford		
	LoE	GR	AGO
▪ Chinesische Kräutermedizin zur Behandlung chemotherapiebedingter Nebenwirkungen	1b	B	-
▪ Homöopathische Medizin gegen therapiebedingte Nebenwirkungen → Topische Kalendula (>=20% Kalendulaanteil) zur Prophylaxe einer akuten Dermatitis unter Strahlentherapie → Traumeel S® Mundspülung bei chemotherapieinduzierter Stomatitis	1b	B	+/-
▪ Topische Anwendung Silymarin (Silybin) (Mariendistelextrakt)	3a	B	+/-
▪ Massage zur Verbesserung von Fatigue, Schmerzen, Angst, Übelkeit	1b	C	+/-
▪ Transkutane elektrische Nervenstimulation (TENS) bei Karzinomschmerzen	2b	D	+/-

1. Chinese medicinal herbs

Zhu L, Li L, Li Y: Chinese Herbal Medicine as an Adjunctive Therapy for Breast Cancer: A Systematic Review and Meta-Analysis. Evid Based Complement Alternat Med. 2016;2016:9469276. doi: 10.1155/2016/9469276.

2. Zhang M, Liu X, Li J, et al. Chinese medicinal herbs to treat the side-effects of chemotherapy in breast cancer patients. Cochrane Database of Systematic Reviews 2007, Issue 2. Art. No.: CD004921. DOI: 10.1002/14651858.CD004921.pub2

1. Homeopathic medicines for adverse effects of cancer treatments

Kassab S, Cummings M, Berkovitz S, et al. Homeopathic medicines for adverse effects of cancer treatments. Cochrane Database of Systematic Reviews 2012, Issue 8. Art. No.: CD004845. DOI: 10.1002/14651858.CD004845.pub2.

Topical use of Silymarin

1. M. Becker-Schiebe et al.:Topical Use of a Silymarin-Based Preparation to Prevent Radiodermatitis. Strahlenther Onkol 2011;187:485–91.
2. Lazzeroni M, Guerrieri-Gonzaga A, Gandini S: A Presurgical Study of Oral Silybin-Phosphatidylcholine in Patients with Early Breast Cancer. Cancer Prev Res (Phila). 2016 Jan;9(1):89-95. doi: 10.1158/1940-6207.

Massage

1. Shin ES, Seo KH, Lee SH, et al. Massage with or without aromatherapy for symptom relief in people with cancer. Cochrane Database of Systematic Reviews

2016, Issue 6. Art. No.: CD009873. DOI: 10.1002/14651858.CD009873.pub3.

2. Robison JG, Smith CL. Therapeutic Massage During Chemotherapy and/or Biotherapy Infusions: Patient Perceptions of Pain, Fatigue, Nausea, Anxiety, and Satisfaction. *Clin J Oncol Nurs.* 2016 Apr;20(2):E34-40. doi: 10.1188/16.CJON.E34-E40.

Transcutaneous electric nerve stimulation (TENS) for cancer pain in adults: (von Slide 16 übertragen)

1. Hurlow A, Bennett MI, Robb KA, et al. Transcutaneous electric nerve stimulation (TENS) for cancer pain in adults. *Cochrane Database of Systematic Reviews* 2012, Issue 3. Art. No.: CD006276. DOI: 10.1002/14651858.CD006276.pub3.
2. Paley CA, Johnson MI, Tashani O et al. Acupuncture for cancer pain in adults. *Cochrane Database of Systematic Reviews* 2011, Issue 1. Art. No.: CD007753. DOI: 10.1002/14651858.CD007753.pub2.
3. Hurlow A, Bennett MI, Robb KA, et al. Transcutaneous electric nerve stimulation (TENS) for cancer pain in adults. *Cochrane Database of Systematic Reviews* 2012, Issue 3. Art. No.: CD006276. DOI: 10.1002/14651858.CD006276.pub3.



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2018.1D

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

Komplementäre Therapien unter (Chemo)therapie Behandlung von Nebenwirkungen

		Oxford		
		LoE	GR	AGO
▪ Akupunktur zur Verbesserung von:				
• Elektro-Akupunktur als Ergänzung zu antiemetischer Therapie bei Chemotherapie-induzierter Übelkeit und Erbrechen	1b	B	+	
▪ Schmerzen				
• Krebs-Schmerzen	1b	B	+	
• AI-induzierte Arthralgie	1a	B	+	
• TENS - transkutane elektrische Nervenstimulation bei Krebs-Schmerzen	2b	D	+/-	
▪ Fatigue	1a	B	+	
▪ Akupressur zur Verbesserung non				
• Übelkeit, Erbrechen ergänzend zu Antiemetika	1b	B	+	
• Fatigue	1b	B	+	
▪ Kognitiver Dysfunktion	5	D	+/-	
▪ Menopausensyndrom bei Patientinnen mit Mammakarzinom	1b	B	+	
• zur Verbesserung v. Häufigkeit und Schwere d. Hitzewallungen	1b	B	+/-	
• Elektroakupunktur zur Verbesserung des Schlafs bei Hitzewallungen	2a	B	+	
▪ Leukopenie (Moxibustion)	2b	B	+/-	
▪ Chemotherapie-induzierter Polyneuropathie				
• als Prophylaxe	1b	B	-	
• als Therapie	2b	B	+/-	
▪ Chronisches Lymphödem nach MaCa Therapie	2b	B	+/-	

Acupuncture

1. Wu X, Chung VCh, Hui EP, et al: Effectiveness of acupuncture and related therapies for palliative care of cancer: overview of systematic reviews. *Sci Rep.* 2015 Nov 26;5:16776.
2. Tao WW, Jiang H, Tao XM, et al. Effects of Acupuncture, Tuina, Tai Chi, Qigong, and Traditional Chinese Medicine Five-Element Music Therapy on Symptom Management and Quality of Life for Cancer Patients: A Meta-Analysis. *J Pain Symptom Manage.* 2016 Apr;51(4):728-47. doi: 10.1016/j.jpainsymman.2015.11.027. Epub 2016 Feb 12. Review.
3. Lau CH, Wu X, Chung VC, et al. Acupuncture and Related Therapies for Symptom Management in Palliative Cancer Care: Systematic Review and Meta-Analysis. *Medicine (Baltimore).* 2016 Mar;95(9):e2901. doi: 10.1097/MD.0000000000002901.

Menopausensyndrom

1. Salehi A, Marzban M, Zadeh AR: Acupuncture for treating hot flashes in breast cancer patients: an updated meta-analysis. *Support Care Cancer.* 2016 Dec;24(12):4895-4899.
2. Lesi G, Razzini G, Musti MA: Acupuncture As an Integrative Approach for the Treatment of Hot Flashes in Women With Breast Cancer: A Prospective Multicenter Randomized Controlled Trial (AcCliMaT). *J Clin Oncol.* 2016 May 20;34(15):1795-802. doi: 10.1200/JCO.2015.63.2893.
3. Chiu HY, Shyu YK, Chang PC: Effects of Acupuncture on Menopause-Related Symptoms in Breast Cancer Survivors: A Meta-analysis of Randomized Controlled

- Trials. Cancer Nurs. 2016 May-Jun;39(3):228-37. doi: 10.1097/NCC.0000000000000278.
4. Tao WW, Jiang H, Tao XM: Effects of Acupuncture, Tuina, Tai Chi, Qigong, and Traditional Chinese Medicine Five-Element Music Therapy on Symptom Management and Quality of Life for Cancer Patients: A Meta-Analysis. J Pain Symptom Manage. 2016 Apr;51(4):728-47. doi: 10.1016/j.jpainsymman.2015.11.027.
 5. Chien TJ, Hsu CH, Liu CY, et al. Effect of acupuncture on hot flush and menopause symptoms in breast cancer- A systematic review and meta-analysis. PLoS One. 2017 Aug 22;12(8):e0180918. doi: 10.1371/journal.pone.0180918. eCollection 2017. Review.
 6. Tao WW, Tao XM, Song CL. Effects of non-pharmacological supportive care for hot flushes in breast cancer: a meta-analysis. Support Care Cancer. 2017 Jul;25(7):2335-2347. doi: 10.1007/s00520-017-3691-y. Epub 2017 Apr 11.

Chemotherapy-induced Nausea and Vomiting

1. Ezzo J, Richardson MA, Vickers A et al.: Acupuncture-point stimulation for chemotherapy-induced nausea or vomiting (Review). The Cochrane Library 2010, Issue 1.
2. Fonnebo V et al.: Acupuncture and acupressure in the treatment of chemotherapy-associated nausea and vomiting. www.cam-cancer.org, Updated May 21, 2009.
3. Streitberger K, Ezzo J, Schneider A: Acupuncture for nausea and vomiting: an update of clinical and experimental studies. Autonomic Neuroscience: Basic and clinical 129 (2006) 107-117.
4. Beith JM, Oh B, Chateld MD, et al. Electroacupuncture for nausea, vomiting, and myelosuppression in women receiving adjuvant chemo- therapy for early breast cancer: a randomized controlled pilot trial. *Medical Acupuncture*. 2012;24:241–248.
5. Shen J, Wenger N, Glaspy J, et al. Electroacupuncture for control of myeloblative chemotherapy-induced emesis: a randomized controlled trial. *JAMA*. 2000;284(21):2755–2761.
6. Dibble SL, Luce J, Cooper BA, et al. Acupressure for chemotherapy- induced nausea and vomiting: a randomized clinical trial. *Oncol Nurs Forum*. 2007;34(4):813–820.
7. Eghbali M et al.. The effect of auricular acupressure on nausea and vomiting caused by chemotherapy among breast cancer patients. *Complement Ther Clin Pract*. 2016 Aug;24:189-94. doi: 10.1016/j.ctcp.2016.06.006. Epub 2016 Jul 5.

Cognitive dysfunction

1. Johnston MF, Yang C, Hui KK et al.: Acupuncture for Chemotherapy-Associated Cognitive Dysfunction: A Hypothesis-Generating Literature Review to Inform Clinical Advice. *Integr Cancer Ther* 2007; 6; 36.

Fatigue

1. Ling WM, Lui LY, So WK, et al: Effects of acupuncture and acupressure on cancer-

- related fatigue: a systematic review. *Oncol Nurs Forum*. 2014 Nov 1;41(6):581-92.
2. Mao JJ, Farrar JT, Bruner D et al. Electroacupuncture for fatigue, sleep, and psychological distress in breast cancer patients with aromatase inhibitor-related arthralgia: a randomized trial. *Cancer*. 2014;120(23):3744-51.
 3. Smith C, Carmady B, Thornton C et al. The effect of acupuncture on post-cancer fatigue and well-being for women recovering from breast cancer: a pilot randomised controlled trial. *Complement Ther Clin Pract*. 2013;19(1):32-5.
 4. Molassiotis et al (2012): Acupuncture for Cancer-Related Fatigue in Patients With Breast Cancer: A pragmatic Randomized Controlled Trial. *J Clin Oncol* published Ahead of Print on October 29.
 5. Zhang Y, Lin L, Li H, et al. Effects of acupuncture on cancer-related fatigue: a meta-analysis. *Support Care Cancer*. 2017 Nov 11. doi: 10.1007/s00520-017-3955-6.
 6. Zick SM, Sen A, Wyatt GK, et al. Investigation of 2 Types of Self-administered Acupressure for Persistent Cancer-Related Fatigue in Breast Cancer Survivors: A Randomized Clinical Trial. *JAMA Oncol*. 2016 Nov 1;2(11):1470-1476. doi: 10.1001/jamaoncol.2016.1867.
 7. Zhang B, Dong JN, Sun P, et al. Effect of therapeutic care for treating fatigue in patients with breast cancer receiving chemotherapy. *Medicine (Baltimore)*. 2017 Aug;96(33):e7750. doi: 10.1097/MD.0000000000007750.

Pain

1. Alimi D et al.: Analgesic effect of auricular acupuncture for cancer pain: A randomized, blinded, controlled trial. *J Clin Oncol* 2003;21(22):4120-26.
2. Garcia MK, McQuade J, Haddad R, et al. Systematic review of acupuncture in cancer care: a synthesis of the evidence. *J Clin Oncol*. 2013 Mar 1;31(7):952-60. doi: 10.1200/JCO.2012.43.5818. Epub 2013 Jan 22. (von Slide 16 übertragen)
3. Bae K, Yoo HS, Lamoury G, et al. Acupuncture for Aromatase Inhibitor-Induced Arthralgia: A Systematic Review. *Integr Cancer Ther*. 2015 Nov;14(6):496-502. doi: 10.1177/1534735415596573. Epub 2015 Jul 28. (von Slide 16 übertragen)
4. Paley CA1, Johnson MI, Tashani OA, et al. Acupuncture for cancer pain in adults. *Cochrane Database Syst Rev*. 2015 Oct 15;10:CD007753. doi: 10.1002/14651858.CD007753.pub3. (von Slide 16 übertragen)
5. TY Choi et al (2012): Acupuncture for the treatment of cancer pain: a systematic review of randomized clinical trials. *Support Care Cancer* 20:1147-1158. (von Slide 16 übertragen)
6. Chen L1, Lin CC1, Huang TW2, et al. Effect of acupuncture on aromatase inhibitor-induced arthralgia in patients with breast cancer: A meta-analysis of randomized controlled trials. *Breast*. 2017 Jun;33:132-138. doi: 10.1016/j.breast.2017.03.015. Epub 2017 Apr 4.
7. Chien TJ1, Liu CY, Chang YF, et al. Acupuncture for treating aromatase inhibitor-related arthralgia in breast cancer: a systematic review and meta-analysis. *J Altern Complement Med*. 2015 May;21(5):251-60. doi: 10.1089/acm.2014.0083. Epub 2015 Apr 27.
8. Yang GS1, Kim HJ, Griffith K et al. Interventions for the Treatment of Aromatase Inhibitor-Associated Arthralgia in Breast Cancer Survivors: A Systematic Review

- and Meta-analysis. *Cancer Nurs.* 2017 Jul/Aug;40(4):E26-E41. doi: 10.1097/NCC.0000000000000409
9. Hershman DL, Unger JM, Greenlee H, et al. Randomized blinded sham- and waitlist-controlled trial of acupuncture for joint symptoms related to aromatase inhibitors in women with early stage breast cancer (S1200). Presented at: 2017 San Antonio Breast Cancer Symposium; Dec. 5-9, 2017; San Antonio, TX. Abstract GS4-04. abstracts2view.com/sabcs/view.php?nu=SABCS17L_503&terms=
 10. Chiu HY, Hsieh YJ, Tsai PS. Systematic review and meta-analysis of acupuncture to reduce cancer-related pain. *Eur J Cancer Care (Engl)*. 2017 Mar;26(2). doi: 10.1111/ecc.12457. Epub 2016 Feb 7.

Leucopenia

1. Choi TY, Lee MS, Ernst E: Moxibustion for the treatment of chemotherapy-induced leukopenia: a systematic review of randomized clinical trials. *Support Care Cancer.* 2015 Jun;23(6):1819-26.
2. Lu W et al.: Acupuncture for chemotherapy-induced neutropenia in patients with gynecologic malignancies: a pilot randomized, sham-controlled clinical trial. *J Altern Complement Med.* 2009 Jul;15(7):745-53.

Chemotherapy induced peripheral neuropathy

1. Greenlee H, Crew KD, Capodice J et al. Randomized sham-controlled pilot trial of weekly electro-acupuncture for the prevention of taxane-induced peripheral neuropathy in women with early stage breast cancer. *Breast Cancer Res Treat.* 2016 Apr;156(3):453-464. doi: 10.1007/s10549-016-3759-2. Epub 2016 Mar 25.
2. Ben-Horin I, Kahan P, Ryvo L et al. Acupuncture and Reflexology for Chemotherapy-Induced Peripheral Neuropathy in Breast Cancer. *Integr Cancer Ther.* 2017 Sep;16(3):258-262. doi: 10.1177/1534735417690254. Epub 2017 Feb 2. (Haben wir die?)
3. Crit Rev Oncol Hematol. 2016 Feb;98:325-34. doi: 10.1016/j.critrevonc.2015.11.014. Epub 2015 Nov 23.
4. Natural products and complementary therapies for chemotherapy-induced peripheral neuropathy: A systematic review.
5. Brami C1, Bao T2, Deng G3.
6. Ligibel et al. San Antonio Breast Cancer Symposium (SABCS) 2016. Abstract PD4-01. Presented December 8, 2016.

Chronisches Lymphödem

1. Curr Oncol. 2016 Feb;23(1):e27-34. doi: 10.3747/co.23.2788. Epub 2016 Feb 18. Effects of warm acupuncture on breast cancer-related chronic lymphedema: a randomized controlled trial. Yao C1, Xu Y1, Chen L1, Jiang H1, Ki CS2, Byun JS3, Bian W1.



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2018.1D

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

Komplementäre Therapien Behandlungsphase – Mind-Body Medizin I

Oxford

LoE GR AGO

MBSR (Mindfulness-Based Stress Reduction –
dt. Achtsamkeits-basierte Stressbewältigung) Programm
verbessert Lebensqualität, Bewältigungsstrategien,
Achtsamkeit, vermindert Stress, Angst, Depression,
Fatigue und Schlafstörung

1a A +

Körperliches Training / Sport

mind. 150 Min. moderates Ausdauertraining pro Woche
in Kombination mit kräftigendem Gerätetraining (2x /Wo.)
verbessern Lebensqualität, kardiorespiratorische Fitness,
körperliche Leistungsfähigkeit, Schlaf und Fatigue, kein
höheres Risiko für die Entstehung von Lymphödemen

1a A ++

Mind-Body Medicine (MBM)

1. A general overview of mind/body therapies in cancer survivorship Elkins G1, Fisher W, Johnson A. Mind-body therapies in integrative oncology. Curr Treat Options Oncol. 2010 Dec;11(3-4):128-40.
2. Monti D, Sufian M, Peterson C (2008): Potential Role of Mind-Body Therapies in Cancer Survivorship. Cancer (supplement),112,11:2607-2616. DOI 101.1002/CNCR.23443.
3. Charalambous A, Giannakopoulou M, Bozas E, et al: Guided Imagery And Progressive Muscle Relaxation as a Cluster of Symptoms Management Intervention in Patients Receiving Chemotherapy: A Randomized Control Trial. PLoS One. 2016 Jun 24;11(6):e0156911. doi: 10.1371/journal.pone.0156911. eCollection 2016.

MBSR

1. Haller H, Winkler MM, Klose P et al: Mindfulness-based interventions for women with breast cancer: an updated systematic review and meta-analysis. Acta oncologica (Stockholm, Sweden). 2017 Jul 07:1-12. PubMed PMID: 28686520. Epub 2017/07/08. eng.
2. Lengacher CA, Reich RR, Paterson CL, et al: Examination of Broad Symptom Improvement Resulting From Mindfulness-Based Stress Reduction in Breast Cancer Survivors: A Randomized Controlled Trial. J Clin Oncol. 2016 Aug 20;34(24):2827-34. doi: 10.1200/JCO.2015.65.7874. Epub 2016 May 31.
3. Huang HP, He M, Wang HY, et al. A meta-analysis of the benefits of mindfulness-based stress reduction (MBSR) on psychological function among breast cancer

- (BC) survivors. *Breast Cancer*. 2015 Mar 28. [Epub ahead of print]
4. Lengacher CA, Reich RR, Paterson CL: Examination of Broad Symptom Improvement Resulting From Mindfulness-Based Stress Reduction in Breast Cancer Survivors: A Randomized Controlled Trial. *J Clin Oncol*. 2016 Aug 20;34(24):2827-34. doi: 10.1200/JCO.2015.65.7874.
 5. Huang HP, He M, Wang HY: A meta-analysis of the benefits of mindfulness-based stress reduction (MBSR) on psychological function among breast cancer (BC) survivors. *Breast Cancer*. 2016 Jul;23(4):568-76. doi: 10.1007/s12282-015-0604-0.
 6. Reich RR, Lengacher CA, Alinat CB: Mindfulness-Based Stress Reduction in Post-treatment Breast Cancer Patients: Immediate and Sustained Effects Across Multiple Symptom Clusters. *J Pain Symptom Manage*. 2017 Jan;53(1):85-95. doi: 10.1016/j.jpainsymman.2016.08.005.
 7. Schmidt ME, Wiskemann J, Ulrich CM: Self-reported physical activity behavior of breast cancer survivors during and after adjuvant therapy: 12 months follow-up of two randomized exercise intervention trials. *Acta Oncol*. 2017 Jan 13:1-10. doi: 10.1080/0284186X.2016.1275776.

Physical exercise

1. Furmaniak AC, Menig M, Markes MH. Exercise for women receiving adjuvant therapy for breast cancer. *Cochrane Database Syst Rev*. 2016 Sep 21;9:CD005001. [Epub ahead of print]
2. Mercier J, Savard J, Bernard P: Exercise interventions to improve sleep in cancer patients: A systematic review and meta-analysis. *Sleep Med Rev*. 2017 Dec;36:43-56. doi: 10.1016/j.smrv.2016.11.001. Epub 2016 Nov 10.
3. Steindorf K; Wiskemann J; Ulrich CM; et al: Effects of exercise on sleep problems in breast cancer patients receiving radiotherapy: a randomized clinical trial. *Breast Cancer Res Treat*. 2017; 162(3):489-499 (ISSN: 1573-7217)
4. Schmitz KH, Courneya KS, Matthews C et al (2010) American College of Sports Medicine roundtable on exercise guidelines for cancer survivors. *Med Sci Sports Exerc* 42(7):1409–1426
5. Loprinzi PD, Cardinal BJ (2012): Effects of physical activity on common side effects of breast cancer treatment. *Breast Cancer*. 2012 Jan;19(1):4-10. doi: 10.1007/s12282-011-0292-3. Epub 2011 Jul 2. Review.
6. Paramanandam VS, Roberts D. Weight training is not harmful for women with breast cancer-related lymphoedema: a systematic review. *J Physiother*. 2014 Sep;60(3):136-43.
7. Cormie P, Pumpa K, Galvao DA et al. Is it safe and efficacious for women with lymphedema secondary to breast cancer to lift heavy weights during exercise: a randomised controlled trial. *J Cancer Surviv*. 2013 Sep;7(3):413-24.
8. Kwan ML, Cohn JC, Armer JM,: Exercise in patients with lymphedema: a systematic review of the contemporary literature. *J Cancer Surviv*. 2011 Dec;5(4):320-36.

Statement on quality of life

1. Steindorf K, Schmidt ME, Klassen O et al. Randomized, controlled trial of resistance training in breast cancer patients receiving adjuvant radiotherapy:

- results on cancer-related fatigue and quality of life. Ann Oncol. 2014 Nov;25(11):2237-43.
2. Schmidt ME, Wiskemann J, Armbrust Pet al. Effects of resistance exercise on fatigue and quality of life in breast cancer patients undergoing adjuvant chemotherapy: A randomized controlled trial. Int J Cancer. 2014 Dec 6. doi: 10.1002/ijc.29383. [Epub ahead of print]
 3. Zeng Y, Huang M, Cheng AS et al. Meta-analysis of the effects of exercise intervention on quality of life in breast cancer survivors. Breast Cancer. 2014 May;21(3):262-74.
 4. Pastakia K, Kumar S (2011): Exercise Parameters in the Management of Breast Cancer: A Systematic Review of Randomized Controlled Trials. Physiother.Research. Res. Int. 16: 237-244.
 5. Duijts SF, Faber MM, Oldenburg HS, van Beurden M, Aaronson NK (2011): Effectiveness of behavioral techniques and physical exercise on psychosocial functioning and health-related quality of life in breast cancer patients and survivors-a meta-analysis. Psycho-Oncology 20: 115–126.
 6. Mishra SI, Scherer RW, Geigle PM, et al. (2012): Exercise interventions on health-related quality of life for cancer survivors. Cochrane Database Syst Rev., 15;8:CD007566. doi: 10.1002/14651858.CD007566.pub2.
 7. Shneerson C, Taskila T, Gale N, et al: The effect of complementary and alternative medicine on the quality of life of cancer survivors: a systematic review and meta-analyses.Complement Ther Med 2013 Aug;21(4):417-29

Cardio respiratory Fitness / Physical Functioning

1. Jones LW et al. (2016) Exercise and Risk of Cardiovascular Events in Women With Nonmetastatic Breast Cancer. J Clin Oncol. 10;34(23):2743-9. doi: 10.1200/JCO.2015.65.6603. Epub 2016 May 23.
2. Casla S, López-Tarruella S, Jerez Y, et al. Supervised physical exercise improves VO₂max, quality of life, and health in early stage breast cancer patients: a randomized controlled trial. Breast Cancer Res Treat. 2015 Sep;153(2):371-82.
3. Courneya KS, McKenzie DC, Mackey JR et al. Effects of exercise dose and type during breast cancer chemotherapy: multicenter randomized trial. J Natl Cancer Inst. 2013 Dec 4;105(23):1821-32.
4. Courneya KS, Segal RJ, Mackey JR et al. Effects of exercise dose and type on sleep quality in breast cancer patients receiving chemotherapy: a multicenter randomized trial. Breast Cancer Res Treat. 2014 Apr;144(2):361-9.
5. Courneya KS, McKenzie DC, Gelmon K et al. A multicenter randomized trial of the effects of exercise dose and type on psychosocial distress in breast cancer patients undergoing chemotherapy. Cancer Epidemiol Biomarkers Prev. 2014 May;23(5):857-64.
6. Courneya KS, McKenzie DC, Mackey JR et al. Subgroup effects in a randomised trial of different types and doses of exercise during breast cancer chemotherapy.Br J Cancer. 2014 Oct 28;111(9):1718-25.

Fatigue

1. Mustian KM, Alfano CM, Heckler C, et al. (2017) Comparison of Pharmaceutical,

- Psychological, and Exercise Treatments for Cancer-Related Fatigue: A Meta-analysis. *JAMA Oncol.* 1;3(7):961-968. doi: 10.1001/jamaoncol.2016.6914.
- 2. Hilfiker, R., Meichtry, A., Eicher, M et al. (2017). Exercise and other non-pharmaceutical interventions for cancer-related fatigue in patients during or after cancer treatment: a systematic review incorporating an indirect-comparisons meta-analysis. *Br J Sports Med*, bjsports-2016.
 - 3. Lipsett A, Barrett S, Haruna F et al. The impact of exercise during adjuvant radiotherapy for breast cancer on fatigue and quality of life: A systematic review and meta-analysis. *Breast*. 2017 Apr;32:144-155. doi: 10.1016/j.breast.2017.02.002. Epub 2017 Feb 9.
 - 4. Juvet LK, Thune I, Elvsaa IKØ, et al. The effect of exercise on fatigue and physical functioning in breast cancer patients during and after treatment and at 6 months follow-up: A meta-analysis. *Breast*. 2017 Jun;33:166-177. doi: 10.1016/j.breast.2017.04.003. Epub 2017 Apr 14.
 - 5. Meneses-Echávez JF, González-Jiménez E, Ramírez-Vélez R. Effects of supervised exercise on cancer-related fatigue in breast cancer survivors: a systematic review and meta-analysis. *BMC Cancer*. 2015 Feb 21;15:77. doi: 10.1186/s12885-015-1069-4.
 - 6. van Waart H, Stuiver MM, van Harten WH et al. Effect of Low-Intensity Physical Activity and Moderate- to High-Intensity Physical Exercise During Adjuvant Chemotherapy on Physical Fitness, Fatigue, and Chemotherapy Completion Rates: Results of the PACES Randomized Clinical Trial. *J Clin Oncol*. 2015 Jun 10;33(17):1918-27.
 - 7. Cramp F, Byron-Daniel J. Exercise for the management of cancer-related fatigue in adults. *Cochrane Database of Systematic Reviews* 2012 Nov 14;11:CD006145.
 - 8. Zou LY, Yang L, He XL et al. Effects of aerobic exercise on cancer-related fatigue in breast cancer patients receiving chemotherapy: a meta-analysis. *Tumour Biol*. 2014 Jun;35(6):5659-67.
 - 9. Brown JC, Huedo-Medina TB, Pescatello LS et al. Efficacy of exercise interventions in modulating cancer-related fatigue among adult cancer survivors: a meta-analysis. *Cancer Epidemiol Biomarkers Prev*. 2011 Jan;20(1):123-33.
 - 10. Steindorf K, Schmidt ME, Klassen O et al. Randomized, controlled trial of resistance training in breast cancer patients receiving adjuvant radiotherapy: results on cancer-related fatigue and quality of life. *Ann Oncol*. 2014 Nov;25(11):2237-43.
 - 11. Schmidt ME1, Wiskemann J, Armbrust P, et al.: Effects of resistance exercise on fatigue and quality of life in breast cancer patients undergoing adjuvant chemotherapy: A randomized controlled trial. *Int J Cancer*. 2014 Dec 6. doi: 10.1002/ijc.29383. [Epub ahead of print]



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2018.1D

www.ago-online.de
**FORSCHEN
LEHREN
HEILEN**

Komplementäre Therapien Behandlungsphase – Mind-Body Medizin II

	Oxford		
	LoE	GR	AGO
Yoga Verbesserung von Lebensqualität, Stress, Fatigue, Schlaf, Angst und Depression	1b	A	+
Qigong Verbesserung von Lebensqualität, Fatigue, Stimmung	2a	B	+/-
Tai-Chi Verbesserung von Lebensqualität, Muskelkraft, Schlaf	2a	B	+/-
Hypnose (in Kombination mit kognitiver Therapie) Verbesserung von Fatigue unter Radiotherapie, Reduktion von Distress	1b	A	+

General

1. Merckaert I, Lewis F, Delevallez F: Improving anxiety regulation in patients with breast cancer at the beginning of the survivorship period: A randomized clinical trial comparing the benefits of single-component and multi-component group interventions. *Psychooncology*. 2016 Oct 8. doi: 10.1002/pon.4294.
2. Montgomery GH, David D, Kangas M: Randomized controlled trial of a cognitive-behavioral therapy plus hypnosis intervention to control fatigue in patients undergoing radiotherapy for breast cancer. *J Clin Oncol*. 2014 Feb 20;32(6):557-63. doi: 10.1200/JCO.2013.49.3437.
3. Pan Y, Yang K, Wang Y, et al.: Could yoga practice improve treatment-related side effects and quality of life for women with breast cancer? A systematic review and meta-analysis. *Asia Pac J Clin Oncol*. 2015 Jan 6. doi: 10.1111/ajco.12329. [Epub ahead of print]

Yoga

1. Cramer H, Lauche R, Klose P, et al. Yoga for improving health-related quality of life, mental health and cancer-related symptoms in women diagnosed with breast cancer. *The Cochrane database of systematic reviews*. 2017 Jan 03;1:CD010802. PubMed PMID: 28045199. Epub 2017/01/04. eng.
2. Chakrabarty J, Vidyasagar MS, Fernandes D, et al: Effectiveness of pranayama on cancer-related fatigue in breast cancer patients undergoing radiation therapy: A randomized controlled trial, *Int J Yoga*. 2015 Jan-Jun; 8(1): 47–53. doi: 10.4103/0973-6131.146062 PMCID: PMC4278135
3. Derry et.al. Yoga and self-reported cognitive problems in breast cancer survivors:

- a randomized controlled trial. Psychooncology. 2015 Aug;24(8):958-66.
4. Sprod LK, Fernandez ID, Janelsins MC, et al: Effects of yoga on cancer-related fatigue and global side-effect burden in older cancer survivors. J Geriatr Oncol. 2015 Jan;6(1):8-14.
 5. Naciye Vardar Yagli*, Ozlem Ulger. The effects of yoga on the quality of life and depression in elderly breast cancer patients. Complement Ther Clin Pract. 2015 Feb;21(1):7-10. doi: 10.1016/j.ctcp.2015.01.002. Epub 2015 Jan 29.
 6. Vardar Yağlı N, Şener G, Arikán H et al. Do yoga and aerobic exercise training have impact on functional capacity, fatigue, peripheral muscle strength, and quality of life in breast cancer survivors? Integr Cancer Ther. 2015 Mar;14(2):125-32. doi: 10.1177/1534735414565699. Epub 2015 Jan 6.
 7. Rao RM, Raghuram N, Nagendra HR, et al. Effects of an integrated Yoga Program on Self-reported Depression Scores in Breast Cancer Patients Undergoing Conventional Treatment: A Randomized Controlled Trial. Indian J Palliat Care. 2015 May-Aug;21(2):174-81. doi: 10.4103/0973-1075.156486.
 8. Sadja J, Mills PJ. Effects of yoga interventions on fatigue in cancer patients and survivors: a systematic review of randomized controlled trials. Explore (NY). 2013 Jul-Aug;9(4):232-43.

Qigong

1. Fong SS, Choi AW, Luk WS: Bone Mineral Density, Balance Performance, Balance Self-Efficacy, and Falls in Breast Cancer Survivors With and Without Qigong Training. Integr Cancer Ther. 2016 Dec 1:1534735416686687. doi: 10.1177/1534735416686687.
2. Larkey LK, Roe DJ, Weihs KL et al. (2015) Randomized controlled trial of Qigong/Tai Chi Easy on cancer-related fatigue in breast cancer survivors. Annals of behavioral medicine : a publication of the Society of Behavioral Medicine. 49: 165-176 .
3. Zeng Y, Luo T, Xie H et al. Health benefits of qigong or tai chi for cancer patients: a systematic review and meta-analyses. Complement Ther Med. 2014 Feb;22(1):173-86.
4. Chan CL, Wang CW, Ho RT, et al. A systematic review of the effectiveness of qigong exercise in supportive cancer care. Support Care Cancer. 2012 Jun;20(6):1121-33. doi: 10.1007/s00520-011-1378-3.

Tai Chi

1. Irwin M R, Olmstead R, Carrillo C et al: Tai Chi Chih Compared With Cognitive Behavioral Therapy for the Treatment of Insomnia in Survivors of Breast Cancer: A Randomized, Partially Blinded, Noninferiority Trial . J Clin Oncol. 2017 Aug 10;35(23):2656-2665. doi: 10.1200/JCO.2016.71.0285. Epub 2017 May 10.
2. Larkey LK, Roe DJ, Weihs KL et al. (2015) Randomized controlled trial of Qigong/Tai Chi Easy on cancer-related fatigue in breast cancer survivors. Annals of behavioral medicine : a publication of the Society of Behavioral Medicine. 49: 165-176 .
3. Pan Y, Yang K, Shi X, et al: Tai chi chuan exercise for patients with breast cancer: a systematic review and meta-analysis. Evid Based Complement Alternat Med.

- 2015;2015:535237.
4. Zeng Y, Luo T, Xie H et al. Health benefits of qigong or tai chi for cancer patients: a systematic review and meta-analyses. Complement Ther Med. 2014 Feb;22(1):173-86.
 5. Yan JH, Pan L, Zhang XM et al. Lack of efficacy of Tai Chi in improving quality of life in breast cancer survivors: a systematic review and meta-analysis. Asian Pac J Cancer Prev. 2014;15(8):3715-20.
 6. Irwin MR, Olmstead R, Breen Ec et al., Tai Chi, Cellular Inflammation, and Transcriptome Dynamics in Breast Cancer Survivors With Insomnia: A Randomized Controlled Trial, J Natl Cancer Inst Monogr 2014 2014 (50) 295-301.

Hypnosis

1. Cramer H, Lauche R, Paul A et al. Hypnosis in Breast Cancer Care: A Systematic Review of Randomized Controlled Trials. Integr Cancer Ther. 2015 Jan;14(1):5-15. Epub 2014 Sep 18.
2. Montgomery GH, Schnur JB, Kravits K. Hypnosis for cancer care: Over 200 years young. CA Cancer J Clin. 2012 Nov 20. doi: 10.3322/caac.21165.



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2018.1D

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

Komplementäre Therapien

Rezidivprävention / Verbesserung Gesamtüberleben I

Beeinflussbare Lebensstilfaktoren – Sport - Genussmittel

	Oxford		
	LoE	GR	AGO
▪ Körperliches Training / Sport (das Äquivalent zu 3-5 Std. mäßiggradigem „Walking“ verbessert DFS und OS und kardiopulmonale Funktion)	2a	A	++
▪ Nikotinabusus	2b	A	+/-
▪ Alkohol (>6 g/die)	2b	A	+/-

Physical exercise

1. Nechuta S, Chen WY, Cai H: A pooled analysis of post-diagnosis lifestyle factors in association with late estrogen-receptor-positive breast cancer prognosis. *Int J Cancer*. 2016 May 1;138(9):2088-97. doi: 10.1002/ijc.29940.
2. de Glas NA, Fontein DB, Bastiaannet E, et al: Physical activity and survival of postmenopausal, hormone receptor-positive breast cancer patients: Results of the tamoxifen exemestane adjuvant multicenter lifestyle study. *Cancer* 2014;120:2847-2854.
3. Schmitz KH, Courneya KS, Matthews C et al (2010) American College of Sports Medicine roundtable on exercise guidelines for cancer survivors. *Med Sci Sports Exerc* 42(7):1409–1426
4. Lynch BM, Friedenreich CM, Winkler EA, et al: Associations of objectively assessed physical activity and sedentary time with biomarkers of breast cancer risk in postmenopausal women: findings from NHANES (2003-2006). *Breast Cancer Res Treat*. 2011 Nov;130(1):183-94.
5. Friedenreich CM, Neilson HK, Woolcott CG, et al: Inflammatory Marker Changes in a Yearlong Randomized Exercise Intervention Trial among Postmenopausal Women. *Cancer Prev Res (Phila)*. 2012 Jan;5(1):98-108.
6. Zeng H, Irwin ML, Lu L, et al: Physical activity and breast cancer survival: an epigenetic link through reduced methylation of a tumor suppressor gene L3MBTL1. *Breast Cancer Res Treat*. 2012 May;133(1):127-35. doi: 10.1007/s10549-011-1716-7. Epub 2011 Aug 12.
7. Rock CL, Doyle C, Demark-Wahnefried W, et al.: Nutrition and physical activity guidelines for cancer survivors. *CA Cancer J Clin*. 2012 Jul-Aug;62(4):243-74. doi:

10.3322/caac.21142. Epub 2012 Apr 26.

Improvements in DFS and OS, prevention of recurrence

1. Schmid D, Leitzmann MF. Physical exercise. Association between physical activity and mortality among breast cancer and colorectal cancer survivors: a systematic review and meta-analysis. *Ann Oncol*. 2014 Jul;25(7):1293-311. doi: 10.1093/annonc/mdu012. Epub 2014 Mar 18. Review.
2. Borch KB, Braaten T, Lund E et al. Physical activity before and after breast cancer diagnosis and survival - the Norwegian women and cancer cohort study. *BMC Cancer*. 2015 Dec 16;15(1):967. doi: 10.1186/s12885-015-1971-9.
3. Lahart IM, Metsios GS, Nevill AM, et al. Physical activity, risk of death and recurrence in breast cancer survivors: A systematic review and meta-analysis of epidemiological studies. *Acta Oncol*. 2015 May;54(5):635-54.
4. Zhong S, Jiang T, Ma T et al. Association between physical activity and mortality in breast cancer: a meta-analysis of cohort studies. *Eur J Epidemiol*. 2014 Jun;29(6):391-404.
5. Ibrahim EM, Al-Homaidh A (2010): Physical activity and survival after breast cancer diagnosis: meta-analysis of published studies. *Med Oncol*. 2011 Sep;28(3):753-65. doi: 10.1007/s12032-010-9536-x. Epub 2010 Apr 22.
6. Beasley JM, Kwan ML, Chen WY, et al: Meeting the physical activity guidelines and survival after breast cancer: findings from the after breast cancer pooling project. *Breast Cancer Res Treat*. 2012 Jan;131(2):637-43.
7. Schmidt ME, Chang-Claude J, Vrieling A et al. Association of pre-diagnosis physical activity with recurrence and mortality among women with breast cancer. *Int J Cancer*. 2013 Sep 15;133(6):1431-40.
8. de Glas NA, Fontein DB, Bastiaannet E et al. Physical activity and survival of postmenopausal, hormone receptor-positive breast cancer patients: results of the Tamoxifen Exemestane Adjuvant Multicenter Lifestyle study. *Cancer*. 2014 Sep 15;120(18):2847-54.
9. Ballard-Barbash R, Friedenreich CM, Courneya KS, et al: Physical activity, biomarkers, and disease outcomes in cancer survivors: a systematic review. *J Natl Cancer Inst*. 2012 Jun 6;104(11):815-40. doi: 10.1093/jnci/djs207. Epub 2012 May 8.

Smoking

1. Pierce JP, Patterson RE, Senger C et al: Lifetime cigarette smoking and breast cancer prognosis in the after breast cancer pooling project. *J Natl Cancer Inst* 2014;106:djt359.
2. Wang K, Li F, Zhang X: Smoking increases risks of all-cause and breast cancer specific mortality in breast cancer individuals: a dose-response meta-analysis of prospective cohort studies involving 39725 breast cancer cases. *Oncotarget*. 2016 Dec 13;7(50):83134-83147. doi: 10.18632/oncotarget.13366.
3. Pierce JP, Patterson RE, Senger CM: Lifetime cigarette smoking and breast cancer prognosis in the After Breast Cancer Pooling Project. *J Natl Cancer Inst*. 2014 Jan;106(1):djt359. doi: 10.1093/jnci/djt359.
4. Holmes MD, Murin S, Chen WY: Smoking and survival after breast cancer

- diagnosis. *Int J Cancer*. 2007 Jun 15;120(12):2672-7.
5. Bérubé S, Lemieux J, Moore L: Smoking at time of diagnosis and breast cancer-specific survival: new findings and systematic review with meta-analysis. *Breast Cancer Res*. 2014 Apr 19;16(2):R42. doi: 10.1186/bcr3646.

Alcohol

1. Larsen SB, Kroman N, Ibfelt EH: Influence of metabolic indicators, smoking, alcohol and socioeconomic position on mortality after breast cancer. *Acta Oncol*. 2015 May;54(5):780-8. doi: 10.3109/0284186X.2014.998774.
2. Bidstrup PE, Dalton SO, Christensen J: Changes in body mass index and alcohol and tobacco consumption among breast cancer survivors and cancer-free women: a prospective study in the Danish Diet, Cancer and Health Cohort. *Acta Oncol*. 2013 Feb;52(2):327-35. doi: 10.3109/0284186X.2012.746466.
3. Simapivapan P, Boltong A, Hodge A.:To what extent is alcohol consumption associated with breast cancer recurrence and second primary breast cancer?: A systematic review. *Cancer Treat Rev*. 2016 Nov;50:155-167. doi: 10.1016/j.ctrv.2016.09.010.
4. Katzke VA, Kaaks R, Kühn T. Lifestyle and cancer risk. *Cancer J*. 2015 Mar-Apr;21(2):104-10.
5. Jayasekara H, MacInnis RJ, Room R, English DR. Long-Term Alcohol Consumption and Breast, Upper Aero-Digestive Tract and Colorectal Cancer Risk: A Systematic Review and Meta-Analysis. *Alcohol Alcohol*. 2015 Sep 22. pii: agv110. [Epub ahead of print]
6. McDonald JA, Goyal A, Terry MB: Alcohol intake and breast cancer risk: Weighing the overall evidence. *Current breast cancer reports* 2013;5
7. Gou YJ, Xie DX, Yang K et al: Alcohol consumption and breast cancer survival: A meta- analysis of cohort studies. *Asian Pacific journal of cancer prevention : APJCP* 2013;14:4785-4790.
8. Kwan ML, Chen WY, Flatt SW et al. Postdiagnosis alcohol consumption and breast cancer prognosis in the after breast cancer pooling project. *Cancer Epidemiol Biomarkers Prev*. 2013 Jan;22(1):32-41. doi: 10.1158/1055-9965.EPI-12-1022. Epub 2012 Nov 13
9. Harris HR, Bergkvist L, Wolk A: Alcohol intake and mortality among women with invasive breast cancer. *Br J Cancer*. 2012 Jan 3.



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2018.1D

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

Komplementäre Therapien

Rezidivprävention / Verbesserung Gesamtüberleben II

Beeinflussbare Lebensstilfaktoren – Ernährung

	Oxford		
	LoE	GR	AGO
▪ Anstreben eines normalen BMI	1a	A	++
▪ Ernährung mit geringem Fettanteil (Ernährungsberatung empfohlen)	1a	B	+
▪ Ballaststoffhaltige Lebensmittel (u.a. Saaten z.B. Leinsamen)	2a	B	+
▪ Beachten genereller Ernährungsempfehlungen (z.B. von DGE, WCRF) im Sinne einer mediterranen (Vollwert-) Ernährung	2a	B	++
▪ Diät-Extreme	2a	B	--

Adherence to normal body weight/BMI

1. (Brenner DR, Brockton NT, Kotsopoulos J: Breast cancer survival among young women: a review of the role of modifiable lifestyle factors. *Cancer Causes Control.* 2016 Apr;27(4):459-72. doi: 10.1007/s10552-016-0726-5.)
2. (Dieli-Conwright CM, Lee K, Kiwata JL: Reducing the Risk of Breast Cancer Recurrence: an Evaluation of the Effects and Mechanisms of Diet and Exercise. *Curr Breast Cancer Rep.* 2016;8(3):139-150.)
3. (Schwingshackl L, Hoffmann G: Adherence to Mediterranean diet and risk of cancer: an updated systematic review and meta-analysis of observational studies. *Cancer Med.* 2015 Dec;4(12):1933-47. doi: 10.1002/cam4.539.)
4. (Ferrini K, Ghelfi F, Mannucci R: Lifestyle, nutrition and breast cancer: facts and presumptions for consideration. *Ecancermedicalscience.* 2015 Jul 23;9:557. doi: 10.3332/ecancer.2015.557. eCollection 2015.)
5. Rock CL, Flatt SW, Byers TE et al. Results of the Exercise and Nutrition to Enhance Recovery and Good Health for You (ENERGY) Trial: A Behavioral Weight Loss Intervention in Overweight or Obese Breast Cancer Survivors. *J Clin Oncol.* 2015 Oct 1;33(28):3169-76. doi: 10.1200/JCO.2015.61.1095. Epub 2015 Aug 17.
6. Ligibel JA, Cirrincione CT, Liu M2 et al. Body Mass Index, PAM50 Subtype, and Outcomes in Node-Positive Breast Cancer: CALGB 9741 (Alliance). *J Natl Cancer Inst.* 2015 Jun 25;107(9). pii: djv179. doi: 10.1093/jnci/djv179. Print 2015 Sep.
7. Azrad M, Demark-Wahnefried W: The association between adiposity and breast cancer recurrence and survival: A review of the recent literature. *Current nutrition reports* 2014;3:9-15.
8. Ligibel JA, Strickler HD: Obesity and its impact on breast cancer: Tumor incidence,

- recurrence, survival, and possible interventions. American Society of Clinical Oncology educational book / ASCO American Society of Clinical Oncology Meeting 2013:52-59.
9. Chlebowski RT: Nutrition and physical activity influence on breast cancer incidence and outcome. *Breast* (Edinburgh, Scotland) 2013;22 Suppl 2:S30-37.
 10. Chan DS, Vieira AR, Aune D, et al: Body mass index and survival in women with breast cancer-systematic literature review and meta-analysis of 82 follow-up studies. *Annals of oncology : official journal of the European Society for Medical Oncology / ESMO* 2014;25:1901-1914.
 11. Jiralerspong S, Goodwin PJ: Obesity and Breast Cancer Prognosis: Evidence, Challenges, and Opportunities. *J Clin Oncol*, 2016; 34:4203-4216
 12. Nechuta S, Chen WY, Cal H et al.: A pooled analysis of post-diagnosis lifestyle factors in association with late estrogen-receptor positive breast cancer prognosis. *Int J Cancer*. 2016 May 1; 138(9): 2088–2097. doi:10.1002/ijc.29940
 13. Lauby-Secretan, B., et al.: Body Fatness and Cancer--Viewpoint of the IARC Working Group. *N Engl J Med*, 2016. 375(8): p. 794-8.

Low-Fat Diet

1. Dieli-Conwright CM, Lee K, Kiwata JL: Reducing the Risk of Breast Cancer Recurrence: an Evaluation of the Effects and Mechanisms of Diet and Exercise. *Curr Breast Cancer Rep*. 2016;8(3):139-150.
2. Chlebowski RT, Blackburn GL, Thomson CA et al. Dietary fat reduction and breast cancer outcome: interim efficacy results from the Women's Intervention Nutrition Study. *J Natl Cancer Inst* 2006; 98:1767-76.
3. Chlebowski RT, Aragaki AK, Anderson GL: Low-Fat Dietary Pattern and Breast Cancer Mortality in the Women's Health Initiative Randomized Controlled Trial. *J Clin Oncol*. 2017 Sep 1;35(25):2919-2926. doi: 10.1200/JCO.2016.72.0326. Epub 2017 Jun 27.
4. Xing MY, Xu SZ, Shen P: Effect of low-fat diet on breast cancer survival: A meta-analysis. *Asian Pacific journal of cancer prevention : APJCP* 2014;15:1141-1144.

(Avoidance of high fat dairy products)

1. Dieli-Conwright CM, Lee K, Kiwata JL: Reducing the Risk of Breast Cancer Recurrence: an Evaluation of the Effects and Mechanisms of Diet and Exercise. *Curr Breast Cancer Rep*. 2016;8(3):139-150.
2. Zang J, Shen M, Du S, Chen T, Zou S. The Association between Dairy Intake and Breast Cancer in Western and Asian Populations: A Systematic Review and Meta-Analysis. *J Breast Cancer*. 2015 Dec;18(4):313-22. doi: 10.4048/jbc.2015.18.4.313. Epub 2015 Dec 23.
3. Xing MY, Xu SZ, Shen P: Effect of low-fat diet on breast cancer survival: A meta-analysis. *Asian Pacific journal of cancer prevention : APJCP* 2014;15:1141-1144.
4. Kroenke CH, Kwan ML, Sweeney C, et al.: High- and low-fat dairy intake, recurrence, and mortality after breast cancer diagnosis. *J Natl Cancer Inst* 2013;105:616-623.)

(Lignans/ flaxseed:

1. Schwingshackl L, Hoffmann G: Adherence to Mediterranean diet and risk of cancer: an updated systematic review and meta-analysis of observational studies. *Cancer Med.* 2015 Dec;4(12):1933-47. doi: 10.1002/cam4.539.
2. Ferrini K, Ghelfi F, Mannucci R: Lifestyle, nutrition and breast cancer: facts and presumptions for consideration. *Ecancermedicalscience.* 2015 Jul 23;9:557. doi: 10.3332/ecancer.2015.557. eCollection 2015.
3. Delman DM, Fabian CJ, Kimler BF et al: Effects of Flaxseed Lignan Secoisolariciresinol Diglucosideon Preneoplastic Biomarkers of Cancer Progression in a Model of Simultaneous Breast and Ovarian Cancer Development. *Nutr Cancer.* 2015;67(5):857-64. doi: 10.1080/01635581.2015.1042549. Epub 2015 May 26.
4. Flower, G., H. Fritz, L. G. Balneaves, et al (2013). "Flax and Breast Cancer: A Systematic Review." *Integr Cancer Ther* 13(3): 181-192.
5. JK, Thompson LU. Flaxseed and its lignan and oil components: can they play a role in reducing the risk of and improving the treatment of breast cancer? *Appl Physiol Nutr Metab.* 2014 Jun;39(6):663-78.
6. McCann SE, Edge SB, Hicks DG et al. A pilot study comparing the effect of flaxseed, aromatase inhibitor, and the combination on breast tumor biomarkers. *Nutr Cancer.* 2014;66(4):566-75.)

Adherence to general nutrition – guidelines:

1. World Cancer Research FundInternational/American Institute for Cancer Research, Continious Update Project: Diet, Nutrition, Physical Activity, and Breast Cancer Survivors. 2014. Available at: www.wcrf.org/sites/default/files/Breast-Cancer-Survivors-2014-Report.pdf
2. Oberritter, H., et al., The DGE nutrition circle—Presentation and basis of the food-related recommendations from the German Nutrition Society (DGE). *Ernährungs Umschau International,* 2013. 60(2): p. 24-29.
3. Ferrini K, Ghelfi F, Mannucci R: Lifestyle, nutrition and breast cancer: facts and presumptions for consideration. *Ecancermedicalscience.* 2015 Jul 23;9:557. doi: 10.3332/ecancer.2015.557. eCollection 2015.
4. Gonzales, J. F., N. D. Barnard, D. J. Jenkins, A. J. Lanou, B. Davis, G. Saxe and S. Levin (2014). "Applying the precautionary principle to nutrition and cancer." *J Am Coll Nutr* 33(3): 239-246.
5. Schwingshackl L, Schwedhelm C: Adherence to Mediterranean Diet and Risk of Cancer: An Updated Systematic Review and Meta-Analysis. *Nutrients* 2017 (9) 1063; doi:10.3390/nu9101063
6. Toledo, E., et al., Mediterranean Diet and Invasive Breast Cancer Risk Among Women at High Cardiovascular Risk in the PREDIMED Trial: A Randomized Clinical Trial. *JAMA Intern Med,* 2015. 175(11): p. 1752-60
7. Biasini C, di Nunzio C, Cordani MR, et al.: Mediterranean Diet influences breast cancer relapse: preliminary results of the SETA PROJECT. *Journal of Clinical Oncology,* 2016 ASCO Annual Meeting (June 3-7, 2016). Vol 34, No 15_suppl (May 20 Supplement), 2016: e13039

Diät Extreme:

1. Huebner J., Marienfeld S. et al.: Counseling Patients on Cancer Diets: A Review of the Literature and Recommendations for Clinical Practice. *Anticancer Res.* 2014 Jan; 34(1):39-48.
2. Erickson, N., Boscheri, A., Linke, B. et al.: Systematic review: isocaloric ketogenic dietary regimes for cancer patients. *Med Oncol* (2017) 34: 72.
<https://doi.org/10.1007/s12032-017-0930-5>



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2018.1D

www.ago-online.de
**FORSCHEN
LEHREN
HEILEN**

Komplementäre Therapien Rezidivprävention / Verbesserung Gesamtüberleben III Pflanzliche Therapieansätze - Nahrungsergänzung

- Nach Systemtherapie – Vitamine/Antioxidanzien scheinen nicht mit einem erhöhtem Rezidivrisiko assoziiert
- Raucher haben ein höheres Bronchial-Ca-Risiko unter Antioxidanzien

Oxford		
LoE	GR	AGO
2b	B	
1b	A	

Prävention eines brustkrebsassoz. Rezidivs

- Antioxidanzien
- Orthomolekulare Substanzen (Selen, Zink ...)
- Vitamine (zusätzlich zu ausgewogener Ernährung; Vit. C, E, D)
- Karotenoide erscheinen mit schlechterem Ergebnis assoziiert
- Proteolytische Enzyme (Papain, Trypsin, Chymotrypsin)
- Sojaprodukte (Phytoöstrogene)
 - Konzentration $\geq 100 \text{ mg Isoflavone pro Tag}$
- Traubensilberkerze (*Cimicifuga racemosa*)
- Mistelklette (*Viscum album*)
- Thymuspeptide (Einfluss auf Überleben)
- Sauerstoff- und Ozon-Therapie
- Antioxidative Supplamente nach Beendigung der Radiotherapie
- Laetrile (Aprikosenkernerktract)
- Methadon
- Cancer bush (*Sutherlandia frutescens*),
Devil's claw (*Harpagophytum procumbens*),
Rooibos Tee (*Aspalathus linearis*),
Bambara-Erdnuss (*Vignea subterranean*)

2a	B	+/-
5	D	-
2a	B	+/-
2b	B	-
3b	B	-
2a	B	+/-
2a	B	-
3b	C	+/-
1b	C	-
2a	B	-
5	D	--
2b	B	+/-
1c	D	--
5	D	--
5	D	-

General

1. Hervik JB, Stub T: Adverse effects of non-hormonal pharmacological interventions in breast cancer survivors, suffering from hot flashes: A systematic review and meta-analysis. *Breast Cancer Res Treat.* 2016 Nov;160(2):223-236.

Post treatment vitamin and/or antioxidant supplements

1. Drewe J, Bucher KA, Zahner C. A systematic review of non-hormonal treatments of vasomotor symptoms in climacteric and cancer patients. *Springerplus.* 2015 Feb 10;4:65. doi: 10.1186/s40064-015-0808-y. eCollection 2015.
2. Sodde VK, Lobo R, Kumar N, et al. Cytotoxic activity of Macrosolen parasiticus (L.) Danser on the growth of breast cancer cell line (MCF-7). *Pharmacogn Mag.* 2015 May;11(Suppl 1):S156-60. doi: 10.4103/0973-1296.157719.
3. Salarabadi A, Bidgoli SA, Madani SH. Roles of Kermanshahi Oil, Animal Fat, Dietary and Non- Dietary Vitamin D and other Nutrients in Increased Risk of Premenopausal Breast Cancer: A Case Control Study in Kermanshah, Iran. *Asian Pac J Cancer Prev.* 2015;16(17):7473-8.
4. Karimi Z, Bahadoran Z, Abedini S, et al. Dietary total antioxidant capacity and the risk of breast cancer: a case-control study. *East Mediterr Health J.* 2015 Sep 28;21(8):564-71.
5. Carioca AA, Verde SM, Luzia LA, et al. Association of oxidative stress biomarkers with adiposity and clinical staging in women with breast cancer. *Eur J Clin Nutr.* 2015 Nov;69(11):1256-61. doi: 10.1038/ejcn.2015.84. Epub 2015 Jun 3.
6. Martin-Herranz A, Salinas-Hernández P. Vitamin D supplementation review and recommendations for women diagnosed with breast or ovary cancer in the

- context of bone health and cancer prognosis/risk. Crit Rev Oncol Hematol. 2015 Oct;96(1):91-9. doi: 10.1016/j.critrevonc.2015.05.006. Epub 2015 May 19.
7. Poole EM, Shu X, Caan BJ, et al: Postdiagnosis supplement use and breast cancer prognosis in the after breast cancer pooling project. Breast cancer research and treatment 2013;139:529-537.
 8. Harris HR, Orsini N, Wolk A: Vitamin c and survival among women with breast cancer: A meta-analysis. European journal of cancer 2014;50:1223-1231.
 9. Mohr SB, Gorham ED, Kim J, et al: Meta-analysis of vitamin d sufficiency for improving survival of patients with breast cancer. Anticancer Res 2014;34:1163-1166.
 10. Greenlee H, Kwan ML, Kushi LH, et al.: Antioxidant supplement use after breast cancer diagnosis and mortality in the life after cancer epidemiology (lace) cohort. Cancer 2012;118:2048-2058.

Soy as normal part of the diet/soy concentrates

1. Wu AH, Spicer D, Garcia A, et al. Double-Blind Randomized 12-Month Soy Intervention Had No Effects on Breast MRI Fibroglandular Tissue Density or Mammographic Density. Cancer Prev Res (Phila). 2015 Oct;8(10):942-51. doi: 10.1158/1940-6207.CAPR-15-0125. Epub 2015 Aug 14.
2. Fritz H, Seely D, Flower G, et al.: Soy, red clover, and isoflavones and breast cancer: A systematic review. PloS one 2013;8:e81968.
3. Messina M, Caan BJ, Abrams DI, et al: It's time for clinicians to reconsider their proscription against the use of soyfoods by breast cancer patients. Oncology (Williston Park, NY) 2013;27

Black cohosh

1. Fritz H, Seely D, McGowan J, et al: Black cohosh and breast cancer: A systematic review. Integrative cancer therapies 2014;13:12-29.
2. Rostock M, Fischer J, Mumm A, et al.: Black cohosh (*Cimicifuga racemosa*) in tamoxifen-treated breast cancer patients with climacteric complaints - a prospective observational study. Gynecol Endocrinol. 2011 Oct;27(10):844-8.
3. Villaseca P. Non-estrogen conventional and phytochemical treatments for vasomotor symptoms: what needs to be known for practice. Climacteric. 2012 Apr;15(2):115-24. doi: 10.3109/13697137.2011.624214. Epub 2011 Dec 8.
4. Geller SE, Shulman LP, van Breemen RB et al.: Safety and efficacy of black cohosh and red clover for the management of vasomotor symptoms: a randomized controlled trial. Menopause. 2009;16(6):1156–1166.

Laetrile treatment for cancer

1. Milazzo S, Ernst E, Lejeune S, et al.: Laetrile treatment for cancer. Cochrane Database of Systematic Reviews 2011, Issue 11. Art. No.: CD005476. DOI: 10.1002/14651858.CD005476.pub3.

St John's Wort

1. Caraci F, Crupi R, Drago F, et al. Metabolic drug interactions between antidepressants and anticancer drugs: focus on selective serotonin reuptake

- inhibitors and hypericum extract. *Curr Drug Metab.* 2011 Jul 1;12(6):570-7.
2. Schellander R, Donnerer J: Antidepressants: clinically relevant drug interactions to be considered. *Pharmacology.* 2010;86(4):203-15. Epub 2010 Sep 8.
 3. Nahrstedt A, Butterweck V: Lessons learned from herbal medicinal products: the example of St. John's Wort (perpendicular). *J Nat Prod.* 2010 May 28;73(5):1015-21.

Red clover

1. Fritz H, Seely D, Flower G et al. Soy, red clover, and isoflavones and breast cancer: A systematic review. *PLoS One.* 2013 Nov 28;8(11):e81968.
2. Geller SE, Shulman LP, van Breemen RB et al.: Safety and efficacy of black cohosh and red clover for the management of vasomotor symptoms: a randomized controlled trial. *Menopause.* 2009;16(6):1156–1166.
3. Booth NL et al.: Clinical studies of red clover (*Trifolium pratense*) dietary supplements in menopause: a literature review. *Menopause,* Vol. 13, No. 2, 2006.
4. Kronenberg F, Fugh-Berman A: Complementary and alternative medicine for menopausal symptoms: a review of randomized, controlled trials. *Ann Intern Med.* 2002;137(10):805–813.

Dong Quai

1. Zhuang SR, Chiu HF, Chen SL, Effects of a Chinese medical herbs complex on cellular immunity and toxicity-related conditions of breast cancer patients. *Br J Nutr.* 2011 Aug 25:1-7.
2. Rotem C, Kaplan B: Phyto-Female Complex for the relief of hot flushes, night sweats and quality of sleep: randomized, controlled, double-blind pilot study. *Gynecol Endocrinol.* 2007;23(2):117-122.
3. Lau CB, Ho TC, Chan TW et al.: Use of dong quai (*Angelica sinensis*) to treat peri- or postmenopausal symptoms in women with breast cancer: is it appropriate? *Menopause.* 2005;12(6):734–740.

Ginseng root

1. Yamada N, Araki H, Yoshimura H: Identification of antidepressant-like ingredients in ginseng root (*Panax ginseng* C.A. Meyer) using a menopausal depressive-like state in female mice: participation of 5-HT2A receptors. *Psychopharmacology (Berl).* 2011 Aug;216(4):589-99.
2. Peralta EA, Murphy LL, Minnis J, et al.: American Ginseng inhibits induced COX-2 and NFkB activation in breast cancer cells. *J Surg Res.* 2009 Dec;157(2):261-7.

Bromelain+Papain+Selen+Lektin bei AI-induced athralgia

1. Uhlenbruck B, Van Leendert R, Schneider B et al.: Reduced side-effects of adjuvant hormone therapy in breast cancer patients by complementary medicine. *In Vivo.* 2010 Sep-Oct;24(5):799-802.