

Diagnostik und Therapie früher und fortgeschrittener Mammakarzinome

Adjuvante endokrine Therapie bei prä- und postmenopausalen Patientinnen



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2020.1D

www.ago-online.de

FORSCHEN
LEHREN
HEILEN


Adjuvante endokrine Therapie bei prä- und postmenopausalen Patientinnen

■ Versionen 2002–2019:

Bauerfeind / Dall / Diel / Fersis / Fehm / Friedrichs / Gerber / Göring / Hanf / Harbeck / Huober / Jackisch / Lisboa / Lück / Lux / Maass / von Minckwitz / Möbus / Müller / Oberhoff / Schaller / Scharl / Schneeweiss / Schütz / Solomeyer / Stickeler / Thomssen / Untch/Fehm/Gerber

■ Version 2020:

Nitz / Huober



© AGO e. V.
in der DGOG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2020.1D

www.ago-online.de

**FORSCHEN
LEBEN
HEILEN**

Bestimmung des Steroid-Hormonrezeptorstatus

Oxford LoE: 1
GR: A
AGO: ++

**„Endokrines Ansprechen“ (früher rezeptorpositiv):
Immunhistologie (ER und / oder PgR)**

0% pos. Zellen:	endokrin nicht sensitiv
1–9% pos. Zellen:	endokrin fraglich sensitiv
≥ 10% pos. Zellen:	endokrin sensitiv

**Hormonrezeptor-Status
unbekannt:**

endokrin sensitiv

**Bei ER negativ / PR positiv (> = 10% Zellen): immunhistochemische Reevaluation
erforderlich**

Endocrine responsiveness:

1. Hammond ME , Hayes DF, Dowsett M et al. American Society of Clinical Oncology/College Of American Pathologists guideline recommendations for immunohistochemical testing of estrogen and progesterone receptors in breast cancer. J Clin Oncol. 2010 Jun 1;28(16):2784-95. Review. Erratum in: J Clin Oncol. 2010 Jul 20;28(21):3543.
2. Early Breast Cancer Trialists Collaborative Group EBCTCG. Effects of chemotherapy and hormonal therapy for early breast cancer on recurrence and 15-year survival: an overview of the randomised trials. Lancet. 2005;365(9472):1687–717.
3. Traub L, Thill M, Nitschmann S: 20-Jahres-Ergebnisse einer 5-jährigen Hormontherapie bei Mammakarzinom : Early Breast Cancer Trialists' Collaborative Group (EBCTCG). Internist (Berl). Springer Medizin 2018;59(4):410–2.
4. Pan H, Gray R, Braybrooke J et al. 20-Year Risks of Breast-Cancer Recurrence after Stopping Endocrine Therapy at 5 Years. N Engl J Med. 2017;377(19):1836–46.

In case of ER negative / PR positive (>10% cells): consider immunohistochemical re-evaluation:

1. Viale G, Regan MM, Maiorano E et al. Prognostic and predictive value of centrally reviewed expression of estrogen and progesterone receptors in a randomized trial comparing letrozole and tamoxifen adjuvant therapy for postmenopausal early

breast cancer: BIG 1-98. J Clin Oncol 2007;25:3846-52.

2. Cserni G, Fracz M, Kalman E et al. Estrogen receptor negative and progesterone receptor positive breast carcinomas-how frequent are they? Pathol Oncol Res 2011;17:663-8.
3. Hefti MM, Hu R, Kniblauch NW et al. Estrogen receptor negative/progesterone receptor positive breast cancer is not a reproducible subtype. Breast Cancer Res 2013;15:R68.
4. Yi M, Huo L, Koenig KB et al. Which threshold for ER positivity? a retrospective study based on 9639 patients. Ann Oncol 2014;25:1004-11.

Adjuvante endokrine Therapie

Bestimmung des Menopausenstatus

Bestimmung des Menopausenstatus:

- Menstruationsanamnese
- FSH, E2

Oxford		
LoE	GR	AGO

++

++

1. Ortmann O, Cufer T, Dixon JM et al. Adjuvant endocrine therapy for perimenopausal women with early breast cancer. Breast 2009;18(1):2-7.
2. Clemons M, Simmons C: Identifying menopause in breast cancer patients: considerations and implications. Breast Cancer Res Treat 2007;104(2):115-20.
3. Su HI, Sammel MD, Green J et al. Antimüllerian hormone and inhibin B are hormone measures of ovarian function in late reproductive-aged breast cancer survivors. Cancer 2010;116(3):592-9.
4. Partridge AH, Ruddy KJ, Gelber S et al. Ovarian reserve in women who remain premenopausal after chemotherapy for early stage breast cancer. Fertil Steril 2010;94(2):638-44.
5. Anders C, Marcom PK, Peterson B et al. A pilot study of predictive markers of chemotherapy-related amenorrhea among premenopausal women with early stage breast cancer. Cancer Invest 2008;26(3):286-95
6. Anderson RA, Cameron DA: Pretreatment serum anti-müllerian hormone predicts long-term ovarian function and bone mass after chemotherapy for early breast cancer. J Clin Endocrinol Metab 2011 96(5):1336-43.
7. Su HI, Chung K, Sammel MD et al. Antral follicle count provides additive information to hormone measures for determining ovarian function in breast cancer survivors. Fertil Steril 2011;95(5):1857-9.
8. Furlanetto J, Nekljudova V: Impact of chemotherapy-induced ovarian failure (CIOF) on disease-free survival (dfs) and overall survival (osl in young women with early breast cancer, ESMO 2019 180 PD



ARBEITSGEMEINSCHAFT
GYNAKOLOGISCHE
ONKOLOGIE e.V.



FORSCHEN
LEHREN
HEILEN

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2020.1D

Adjuvante endokrine Therapie

Endokrine Therapie:

- Endokrin sensitiv
- fraglich endokrin sensitiv
- Endokrine Therapie sequentiell:
nach einer adjuvanten Chemotherapie
- Nicht endokrin sensitiv:
keine endokrine Therapie

Oxford		
LoE	GR	AGO
1a	A	++
3b	D	+
1a	A	++
1a	A	++

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

1. Early Breast Cancer Trialists' Collaborative Group. Effects of chemotherapy and hormonal therapy for early breast cancer on recurrence and 15-year survival: an overview of randomised trials. Lancet 2005;365:1687-717.
2. Early Breast Cancer Trialists' Collaborative Group (EBCTCG). Relevance of breast cancer hormone receptors and other factors to the efficacy of adjuvant tamoxifen: patient-level meta-analysis of randomised trials. Lancet 2011;378(9793):771-84.
3. Hackshaw A, Roughton M, Forsyth S et al. Long-term benefits of 5 years of tamoxifen: 10-year follow-up of a large randomized trial in women at least 50 years of age with early breast cancer. J Clin Oncol 2011;29(13): 1657-63.
4. Albain KS, Barlow WE, Ravdin PM, et al. Breast Cancer Intergroup of North America. Adjuvant chemotherapy and timing of tamoxifen in postmenopausal patients with endocrine-responsive, node-positive breast cancer: a phase 3, open-label, randomised controlled trial. Lancet 2009;374(9707):2055-63.
5. Bedognetti D, Sertoli MR, Pronzato P, et al. Concurrent vs sequential adjuvant chemotherapy and hormone therapy in breast cancer: a multicenter randomized phase III trial. J Natl Cancer Inst 2011;103(20):1529-39.
6. Fujii T, Kogawa T, Dong W et al. Revisiting the definition of estrogen receptor positivity in Her2-negative primary breast cancer. Ann Oncol 2017;28:2420-2428
7. Curigliano G, Burstein HJ, P Winer E et al. De-Escalating and escalating treatment for early-stage breast cancer: the St. Gallen International Expert Consensus Conference on the Primary Therapy of Early Breast Cancer. Ann Oncol 2017;28:1700-1712 .
8. Regan MM, Walley BA, Francis PA et al. Concurrent and sequential initiation of ovarian function suppression with chemotherapy

in premenopausal women with endocrine-responsive early breast cancer: an exploratory analysis of TEXT and SOFT. *Ann Oncol* 2017;28:2225-2232.

9. Villegas S, Lederer B: Similarities between low hormone receptor positive and hormone receptor negative breast cancer: an analysis of 4366 patients from multicenter clinical trials, SABCS 2018 P2-08-10

Generelle Prinzipien der adjuvanten endokrinen Therapie AGO ++

- Die adjuvante endokrine Therapie wird in die initiale Therapie (Jahre 0–5) und die erweiterte adjuvante Therapie (EAT, Jahre 6–15) eingeteilt.
- Standard Therapiedauer der adjuvanten Therapie: 5 Jahre
- Erweiterte Therapiedauer nach individueller Nutzen-Risiko-Abwägung.
- Dauer, Wahl & Sequenz von AI oder Tam hängen v.a. von Menopausenstatus, Verträglichkeit und dem Rückfall-Risiko ab.
- Der Wechsel auf eine andere endokrine Therapie (Tam oder AI) ist besser als die Therapie zu stoppen.
- Beginn mit AI bei postmenopausalen Patientinnen insbesondere bei lobulären Karzinomen und erhöhtem Rückfall-Risiko.
- Es existiert kein ausreichend validierter Biomarker für einen frühen versus einen späten Rückfall.

1. Ingle JN: Overview of adjuvant trials of aromatase inhibitors in early breast cancer. Steroids 2011;76(8):765-7.
2. Higgins MJ, Liedke PE, Goss PE et al. Extended adjuvant endocrine therapy in hormone dependent breast cancer: the paradigm of the NCIC-CTG MA.17/BIG 1-97 trial. Crit Rev Oncol Hematol 2013;86(1):23-32.
3. Regan MM, Neven P, Giobbie-Hurder A et al. BIG 1-98 Collaborative Group; International Breast Cancer Study Group (IBCSG). Assessment of letrozole and tamoxifen alone and in sequence for postmenopausal women with steroid hormone receptor-positive breast cancer: the BIG 1-98 randomised clinical trial at 8.1 years median follow-up. Lancet Oncol 2011;12(12):1101-8.
4. Early Breast Cancer Trialists' Collaborative Group (EBCTCG): Aromatase inhibitors versus tamoxifen in early breast cancer: patient-level meta-analysis of the randomised trials. Lancet 2015;386(10001):1341-52.
5. Rydén L, Heibert Arnlind M, Vitols S et al. Aromatase inhibitors alone or sequentially combined with tamoxifen in postmenopausal early breast cancer compared with tamoxifen or placebo - Meta-analyses on efficacy and adverse events based on randomized clinical trials. Breast 2016;26:106-140.
6. Goss PE, Ingle JN, Pritchard KI et al. Extending aromatase-inhibitor adjuvant therapy to 10 years. N Engl J Med 2016;375(3):209.
7. Pan H, Gray R, Braybrooke J et al. 20-year risks of breast recurrence after stopping endocrine therapy at 5 years. N Engl J Med 2017;1836-49.
8. Burstein HJ, Lacchetti C, Anderson H et al. Adjuvant endocrine therapy for women with hormone receptor-positive breast cancer: ASCO clinical practice guideline focused update. J Clin Oncol 2018 Nov 19;JCO1801160. doi: 10.1200/JCO.18.01160

9. Strasser-Weippl K, Sudan G, Ramjeesingh R et al. Outcomes in women with invasive ductal or invasive lobular early stage breast cancer treated with anastrozole or exemestane in CCTG (NCIC CTG) MA.27. *Eur J Cancer* 2018;90:19-25.
10. Goldvaser H, Barnes TA, Šeruga B, et al. Toxicity of extended adjuvant therapy with aromatase inhibitors in early breast cancer: a systematic review and meta-analysis. *J Natl Cancer Inst.* 2018;110(1)djx141.
11. van Hellemond I, Geurts SME, Tjan-Heijnen VCG: Current status of extended adjuvant endocrine therapy in early stage breast cancer. *Curr Treat Options in Oncol* 2018;19:26.
12. Regan MM, Walley BA, Francis PA et al. Concurrent and sequential initiation of ovarian function suppression with chemotherapy in premenopausal women with endocrine-responsive early breast cancer: an exploratory analysis of TEXT and SOFT. *Ann Oncol* 2017;28:2225-2232.
13. Blok EJ, Kroep JR, Meershoek-Klein Kranenbarg E et al. Treatment decisions and the impact of adverse events before and during extended endocrine therapy in postmenopausal early breast cancer. *Eur J Cancer* 2018;95:59-67.
14. Blok EJ, Kroep JR, Meershoek-Klein Kranenbarg E et al: Relevant factors for the optimal duration of extended endocrine therapy in early breast cancer. *Breast Cancer Res Treat* 2018;168:413-420.
15. Clement Z, Kollias J, Bingham J et al: Extended duration of adjuvant aromatase inhibitor in breast cancer: a meta-analysis of randomized controlled trials. *Gland Surg* 2018;7:449-457.

Adjuvante endokrine Therapie bei prämenopausalen Patientinnen (Jahr 0–5)			
	Oxford		
	LoE	GR	AGO
■ Tamoxifen* 5–10 Jahre	1a	A	++
■ GnRHa Monotherapie (Bei relevanten Kontraindikationen für Tam, gegenüber keiner Therapie)	1a	B	+
■ Ohne Indikation zu neo-/adjuvanter Chemotherapie mit erhaltener Ovarialfunktion:			
■ Tamoxifen	1b	B	++
■ Tamoxifen + OFS	1b	B	+/-
■ AI + OFS	1b	B	+/-
■ Nach neo-/adjuvanter Chemotherapie mit erhaltener Ovarialfunktion **:			
■ Tamoxifen + OFS 5 Jahre	1b	B	+
→ Bei Patientinnen < 35 Jahre	1b	B	++
■ AI + OFS 5 Jahre	1b	B	+/-
→ Bei Patientinnen < 35 Jahre	1b	B	+

OFS: Ovarialfunktions-Suppression;
 * Behandlung nur solange sie tolerabel ist und die Pat. eindeutig prämenopausal ist
 ** Bei Wiedereintritt der Ovarialfunktion innerhalb von 24 Monaten

Tamoxifen 5-10 yrs:


1. Early Breast Cancer Trialists' Collaborative Group (EBCTCG). Effects of chemotherapy and hormonal therapy for early breast cancer on recurrence and 15-year survival: an overview of the randomised trials. Lancet 2005;365:1687-717.
2. Early Breast Cancer Trialists' Collaborative Group (EBCTCG). Relevance of breast cancer hormone receptors and other factors to the efficacy of adjuvant tamoxifen: patient-level meta-analysis of randomised trials. Lancet 2011;378:771-84.
3. Davies C, Pan H, Godwin J et al. Long-term effects of continuing adjuvant tamoxifen to 10 years versus stopping at 5 years after diagnosis of oestrogen receptor-positive breast cancer: ATLAS, a randomised trial. Lancet 2013;381:805-806.
4. Tormey DC, Gray R, Falkson HC: Postchemotherapy adjuvant tamoxifen therapy beyond five years in patients with lymph node-positive breast cancer. Eastern Cooperative Oncology Group. J Natl Cancer Inst 1996;88:1828-33.
5. Goel S, Sharma R, Hamilton A et al: LHRH agonists for adjuvant therapy of early breast cancer in premenopausal women. Cochrane Database Syst Rev. 20097;(4):CD004562.
6. Ruhstaller T, Giobbie-Hurder A: Adjuvant Letrozole and Tamoxifen Alone or Sequentially for Postmenopausal Women With Hormone Receptor-Positive Breast Cancer: Long-Term Follow-Up of the BIG 1-98 Trial JCO 37:105-114

GnRH as monotherapy:

1. Cuzick J, Ambroisine L, Davidson N et al: Use of luteinising-hormone-releasing hormone agonists as adjuvant treatment in premenopausal patients with hormone-receptor-positive breast cancer: a meta-analysis of individual patient data from randomised adjuvant trials. *Lancet* 2007; 369:1711-23.

Ovarian function suppression (OFS) with Tam/AI and Tam with or without OFS:

1. Goel S, Sharma R, Hamilton A et al: LHRH agonists for adjuvant therapy of early breast cancer in premenopausal women. *Cochrane Database Syst Rev* 2009;(4):CD004562.
2. Francis PA, Regan MM, Fleming GF et al. The SOFT Investigators and the International Breast Cancer Study Group. Adjuvant Ovarian Suppression in Premenopausal Breast Cancer. *N Engl J Med* 2015;372(5):436-46.
3. Pagani O, Regan MM, Walley BA et al. TEXT and SOFT Investigators; International Breast Cancer Study Group. Adjuvant exemestane with ovarian suppression in premenopausal breast cancer. *N Engl J Med* 2014;371(2):107-18.
4. Gnant M, Mlineritsch B, Schippinger W et al: Endocrine therapy plus zoledronic acid in premenopausal breast cancer. *N Engl J Med* 2009;360(7):679-91.
5. Shiba E, Yamashita H, Kurebayashi J et al. A randomized controlled study evaluating safety and efficacy of leuprorelin acetate every-3-months depot for 2 versus 3 or more years with tamoxifen for 5 years as adjuvant treatment in premenopausal patients with endocrine-responsive breast cancer. *Breast Cancer* 2016;23(3):499-509.
6. JSaha P, Regan MM, Pagani O: Treatment efficacy, adherence, and quality of life among younger than 35 years in the International Breast Cancer Study Group TEXT and SOFT adjuvant endocrine therapy trial. *J Clin Oncol* 2017;35:3113-3122.
7. Francis PA, Pagani O, Fleming GF et al. Tailoring adjuvant endocrine therapy for premenopausal breast cancer. *N Engl J Med* 2018; 379: 122-137
8. Kim HA, Lee JW, Nam SJ et al. Adding Ovarian Suppression to Tamoxifen for Premenopausal Breast Cancer: A Randomized Phase III Trial. *J Clin Oncol*. 2019, <https://doi.org/10.1200/JCO.19.0012>



1. Pagani O, Gelber S, Colleoni M, et al. Impact of SERM adherence on treatment effect: International Breast Cancer Study Group Trials 13-93 and 14-93. Breast Cancer Res Treat 2013;142(2):455-9.
2. Francis PA, Regan MM, Fleming GF et al. The SOFT Investigators and the International Breast Cancer Study Group. Adjuvant Ovarian Suppression in Premenopausal Breast Cancer. N Engl J Med 2015;372(5):436-46.

Adjuvante endokrine Therapie bei postmenopausalen Patientinnen (Jahre 0-5)			
	Oxford		
	LoE	GR	AGO
■ Aromatasehemmer für die ersten 5 Jahre			
■ Nicht-steroidaler AI bei lobulärem Karzinom	1a	A	++
■ Hohes Rezidivrisiko	2b	B	+
■ Sequentielle Therapie für die ersten 5 Jahre*			++
■ Tam (2–3 Jahre) gefolgt von AI bis zur Gesamtdauer von 5 Jahren	1a	A	
■ AI (2–3 Jahre) gefolgt von Tamoxifen bis zur Gesamtdauer von 5 Jahren	1b	C	
■ Tamoxifen 20 mg/d für die ersten 5 Jahre**	1a	A	+

* Die endokrine adjuvante Therapie postmenopausaler Patientinnen sollte in den ersten 5 Jahren für 2–3 Jahre einen Aromatasehemmer enthalten

** Eine Monotherapie mit Tamoxifen kann bei Patientinnen im Senium, bei niedrigem Risiko oder bei Kontraindikation für Aromatasehemmer eingesetzt werden

www.ago-online.de
FORSCHEN
LEHREN
HEILEN

AI for first 5 years:

1. Early Breast Cancer Trialists' Collaborative Group (EBCTCG): Aromatase inhibitors versus tamoxifen in early breast cancer: patient-level meta-analysis of the randomised trials. Lancet 2015;386(10001):1341-52.
2. Rydén L, Heibert Arnlin M, Vitols S et al. Aromatase inhibitors alone or sequentially combined with tamoxifen in postmenopausal early breast cancer compared with tamoxifen or placebo - Meta-analyses on efficacy and adverse events based on randomized clinical trials. Breast 2016;26:106-14.

-Especially in case of lobular cancer

3. Strasser-Weippl K et al. Outcomes in women with invasive ductal or invasive lobular early stage breast cancer treated with anastrozole or exemestane in CCTG (NCIC CTG) MA.27. Eur J Cancer 2018;90:19-25. doi: 10.1016/j.ejca.2017.11.014

-High risk of recurrence:

4. Early Breast Cancer Trialists' Collaborative Group (EBCTCG): Aromatase inhibitors versus tamoxifen in early breast cancer: patient-level meta-analysis of the randomised trials. Lancet 2015;386(10001):1341-52.

Sequential therapy for first 5 years:

Tam (2-3 yrs.) followed by AI to complete 5 years

AI (2-3 yrs.) followed by Tam to complete 5 years

1. Early Breast Cancer Trialists' Collaborative Group (EBCTCG): Aromatase inhibitors versus tamoxifen in early breast cancer: patient-level meta-analysis of the randomised trials. Lancet 2015;386(10001):1341-52.
2. Rydén L, Heibert Arnlin M, Vitols S et al. Aromatase inhibitors alone or sequentially combined with tamoxifen in postmenopausal early breast cancer compared with tamoxifen or placebo - Meta-analyses on efficacy and adverse events based on randomized clinical trials. Breast 2016;26:106-14.
3. Derks MGM, Blok EJ, Seynaeve C et al. Adjuvant tamoxifen and exemestane in women with postmenopausal early breast cancer (TEAM): 10-year follow-up of a multicentre, open-label, randomised, phase 3 trial. Lancet Oncol 2017;18:1211-1220.
4. Ruhstaller T, Giobbie-Hurder A, Colleoni M et al. Adjuvant letrozole and tamoxifen alone or sequentially for postmenopausal women with hormone receptor-positive breast cancer: long-term follow-up of the BIG 1-98 trial. J Clin Oncol 2019;37(2):105-114.


Tamoxifen 20 mg/d for first 5 yrs:

1. Early Breast Cancer Trialists' Collaborative Group (EBCTCG), et al. Relevance of breast cancer hormone receptors and other factors to the efficacy of adjuvant tamoxifen: patient-level meta-analysis of randomised trials. Lancet 378:771-84, 2011
2. Early Breast Cancer Trialists' Collaborative Group (EBCTCG) et al. Aromatase inhibitors versus tamoxifen in early breast cancer: patient-level meta-analysis of the randomised trials. Lancet 2015;386:1341-52.
3. Rydén L, Heibert Arnlin M, Vitols S et al. Aromatase inhibitors alone or sequentially combined with tamoxifen in postmenopausal early breast cancer compared with tamoxifen or placebo - Meta-analyses on efficacy and adverse events based on randomized clinical trials. Breast. 2016;26:106-14.

Patient care/ adherence and side effects


1. Inwa Id EC, Koller M, Klinkhammer-Schalke M et al. Adjuvant endocrine therapy in pre- versus postmenopausal patients with steroid hormone receptor-positive breast cancer: results from a large population-based cohort of a cancer registry. *J Cancer Res Clin Oncol* 2015;141(12):2229-40.
2. Markopoulos C, Koukouras D, Venizelos V et al. Impact of chemotherapy followed by aromatase inhibitors on bone health of women with ER-positive early breast cancer in real world clinical settings in Greece: Results of the POCHARBI trial conducted by the Hellenic Society of Breast Surgeons. *Breast* 2016 ;27:27-34.
3. Kesmodel SB, Goloubeva OG, Rosenblatt PY et al. Patient-reported adherence to adjuvant aromatase inhibitor therapy using the Morisky Medication Adherence Scale: An evaluation of predictors. *Am J Clin Oncol* 2018;41(5):508-512.

Aromatase Inhibitor vs. Tamoxifen vs. Sequentieller Therapie - 5 Jahre Upfront Therapie



Aromatase inhibitors versus tamoxifen in early breast cancer: patient-level meta-analysis of the randomised trials.
Early Breast Cancer Trialists' Collaborative Group (EBCTCG). Lancet. 2015 Oct 3;386(10001):1341-52.

1. Early Breast Cancer Trialists' Collaborative Group (EBCTCG) et al. Aromatase inhibitors versus tamoxifen in early breast cancer: patient-level meta-analysis of the randomised trials. Lancet. 2015;386(10001):1341-52.


	Erweiterte adjuvante endokrine Therapie (EAT) bei prämenopausalen Patientinnen (Jahre 6–10)		
© AGO e. V. in der DGGG e.V. sowie in der DKG e.V. Guidelines Breast Version 2020.1D	Bei erhöhtem Rückfallrisiko		
www.ago-online.de FORSCHEN LEHREN HEILEN		Oxford LoE GR AGO	
■ 5 Jahre Tamoxifen nach 5 Jahren Tamoxifen	1a	A	++
■ 2,5–5 Jahre AI nach 5 Jahren Tamoxifen prämenopausal, bei im Verlauf eindeutig nachgewiesener postmenopausaler Situation	1b	B	+
■ 5 Jahre Tamoxifen nach 5 Jahre endokriner Therapie + OFS	5	D	+

5 years Tamoxifen after 5 years Tamoxifen:

1. Davies C, Pan H, Godwin J et al. Adjuvant Tamoxifen: Longer Against Shorter (ATLAS) Collaborative Group. Long-term effects of continuing adjuvant tamoxifen to 10 years versus stopping at 5 years after diagnosis of oestrogen receptor-positive breast cancer: ATLAS, a randomised trial. Lancet 2013;381(9869):805-16. Erratum in: Lancet. 2013;381(9869):804.
2. Gray RG, Rea D, Handley K et al. ATTom: long-term effects of continuing adjuvant tamoxifen to 10 years versus stopping at 5 years in 6953 women with early breast cancer. J Clin Oncol 2013; 31 (18 suppl):5.
3. Petrelli F, Coinu A, Cabiddu M et al. Five or more years of adjuvant endocrine therapy in breast cancer: a meta-analysis of published randomised trials. Breast Cancer Res Treat 2013;140(2):233-40.
4. Burstein HJ, Temin S, Anderson H et al. Adjuvant endocrine therapy for women with hormone receptor-positive breast cancer: american society of clinical oncology clinical practice guideline focused update. J Clin Oncol 2014;32(21):2255-69.

2–5 years AI after 5 years Tamoxifen in initially premenopausal patients with validated postmenopausal status in the course of therapy:

1. Goss PE, Ingle JN, Martino S et al. Randomized trial of letrozole following tamoxifen as extended adjuvant therapy in receptor-positive breast cancer: updated findings from NCIC CTG MA.17. J Natl Cancer Inst 2005;97(17):1262-71.
2. Jin H, Tu D, Zhao N et al. Longer-term outcomes of letrozole versus placebo after 5 years of tamoxifen in the NCIC CTG MA.17 trial: analyses adjusting for treatment crossover. J Clin Oncol 2012;30(7):718-21
3. Burstein HJ, Temin S, Anderson H, et al. Adjuvant endocrine therapy for women with hormone receptor-positive breast cancer: american society of clinical oncology clinical practice guideline focused update. J Clin Oncol. 2014;32(21):2255-69.



ARBEITSGEMEINSCHAFT
GYNAEKOLOGISCHE
ONKOLOGIE e.V.

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2020.1D

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

Erweiterte adjuvante endokrine Therapie (EAT) bei postmenopausalen Patientinnen (Jahre 6–10)

Bei erhöhtem Rückfallrisiko:

	Oxford		
	LoE	GR	AGO
▪ Nach 5 Jahren Tamoxifen, Tamoxifen für 5 Jahre	1a	A	+
▪ Nach 5 Jahren Tamoxifen, AI für 2 bis 5 Jahre	1a	A	++
▪ Nach initialer AI-haltiger Therapie (upfront oder Switch) Verlängerung der endokrinen Therapie mit AI für 2 bis 5 * Jahre			
▪ höheres Rückfall-Risiko und bei guter Verträglichkeit des AIs	1a	A	+
▪ niedriges Rückfall-Risiko, schlechte Verträglichkeit des AIs	1a	A	-
▪ Therapiepause bis zu 3 Monaten unter kontinuierlicher EAT mit AI	1b	B	+/-

* Kein Einfluss auf das Gesamtüberleben (OS)

5 years Tamoxifen after 5 years Tamoxifen:

1. Davies C, Pan H, Godwin J et al. Adjuvant Tamoxifen: Longer Against Shorter (ATLAS) Collaborative Group. Long-term effects of continuing adjuvant tamoxifen to 10 years versus stopping at 5 years after diagnosis of oestrogen receptor-positive breast cancer: ATLAS, a randomised trial. Lancet 2013;381(9869):805-16. Erratum in: Lancet. 2013;381(9869):804.
2. Gray RG, Rea D, Handley K et al. ATTom: long-term effects of continuing adjuvant tamoxifen to 10 years versus stopping at 5 years in 6953 women with early breast cancer. J Clin Oncol 2013; 31 (18 suppl):5.
3. Petrelli F, Coinu A, Cabiddu M et al. Five or more years of adjuvant endocrine therapy in breast cancer: a meta-analysis of published randomised trials. Breast Cancer Res Treat 2013;140(2):233-40.
4. Burstein HJ, Lacchetti C, Anderson H et al. Adjuvant endocrine therapy for women with hormone receptor-positive breast cancer: ASCO clinical practice guideline focused update. J Clin Oncol. 2018 Nov 19;JCO1801160. doi: 10.1200/JCO.18.01160

2–5 years AI after 5 years Tamoxifen

1. Goss PE, Ingle JN, Martino S et al. Randomized trial of letrozole following tamoxifen as extended adjuvant therapy in receptor-positive breast cancer: updated findings from NCIC CTG MA.17. J Natl Cancer Inst 2005;97(17):1262-71.


2. Jin H, Tu D, Zhao N et al. Longer-term outcomes of letrozole versus placebo after 5 years of tamoxifen in the NCIC CTG MA.17 trial: analyses adjusting for treatment crossover. *J Clin Oncol* 2012;30(7):718-21.
3. Jakesz R, Greil R, Gnant M et al. Austrian Breast and Colorectal Cancer Study Group. Extended adjuvant therapy with anastrozole among postmenopausal breast cancer patients: results from the randomized Austrian Breast and Colorectal Cancer Study Group Trial 6a. *J Natl Cancer Inst.* 2007;99(24):1845-53. Erratum in: *J Natl Cancer Inst* 2008;100(3):226.
4. Mamounas EP, Jeong JH, Wickerham DL et al. Benefit from exemestane as extended adjuvant therapy after 5 years of adjuvant tamoxifen: intention-to-treat analysis of the National Surgical Adjuvant Breast And Bowel Project B-33 trial. *J Clin Oncol* 2008;26(12):1965-71.
5. Burstein HJ, Lacchetti C, Anderson H et al. Adjuvant endocrine therapy for women with hormone receptor–positive breast cancer: ASCO clinical practice guideline focused update. *J Clin Oncol.* 2018 Nov 19;JCO1801160. doi: 10.1200/JCO.18.01160
6. Gnant M, G Steger, R Greil, et al. A prospective randomized multi-center phase-III trial of additional 2 versus additional 5 years of Anastrozole after initial 5 years of adjuvant endocrine therapy - results from 3,484 postmenopausal women in the ABCSG-16 trial. *SABCS 2017; GS3-01*
7. Gray R (EBCTCG) et al. Extended aromatase inhibitor treatment following 5 or more years of endocrine therapy: a metaanalysis of 22192 women in 11 randomised trials. *SABCS 2018;GS3-03*
8. Zackariah C, Kollias J, Bingham J et al. Extended duration of adjuvant aromatase inhibitor in breast cancer: a meta-analysis of randomized controlled trials. *Gland Surg* 2018;7(5):449-457.
9. Mamounas EP, Bandos H, Lembersky BC et al. Use of letrozole after aromatase inhibitor-based therapy in postmenopausal breast cancer (NRG Oncology/NSABP B-42): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Oncol* 2019;20(1):88-99.
10. Del Mastro L, Masutti M: Benefit from letrozole as extended adjuvant therapy after sequential endocrine therapy: a randomized phase III trial of the Gruppo Italiano Mammella, *ASCO 2019, abstract 505*
11. Mamounas EP, Bandos H: Ten year results from NRG/NSABP – B42: a randomized , double blinded placebo controlle clinical trial of extended adjuvant endocrine therapy with letrozole in postmenopuasal women with hormone receptor + breast cancer who have completed prevoius adjuvant therapy with an aromatase inhibitor after initial AI containing therapy (upfront or switch) further prolongation of endocrine therapy with AI 2-5years. *SABCS 2019, GS4-01*

low risk, poor tolerability of the AI


1. Blok EJ, Kroep JR, Meershoek-Klein Kranenbarg E et al. Optimal Duration of Extended Adjuvant Endocrine Therapy for early breast cancer; results of the IDEAL trial (BOOG 2006-05). J Natl Cancer Inst 2018;110(1): dxx134
2. Mamounas EP, Bandos H, Lembersky BC et al. Use of letrozole after aromatase inhibitor-based therapy in postmenopausal breast cancer (NRG Oncology/NSABP B-42): a randomised, double-blind, placebo-controlled, phase 3 trial. Lancet Oncol 2019;20(1):88-99.
3. Tjan-Heijnen VCG, van Hellemond IEG, Peer PGM et al. Extended adjuvant aromatase inhibition after sequential endocrine therapy (DATA): a randomised, phase 3 trial. Lancet Oncol 2017;18(11):1502-1511.
4. Goss PE, Ingle JN, Pritchard KI et al. Extending Aromatase-Inhibitor Adjuvant Therapy to 10 Years. N Engl J Med. 2016;375(3):209-19.
5. Zdenkowski N, Forbes JF, Boyle FM et al. Australia and New Zealand Breast Cancer Trials Group. Observation versus late reintroduction of letrozole as adjuvant endocrine therapy for hormone receptor-positive breast cancer (ANZ0501 LATER): an open-label randomised, controlled trial. Ann Oncol 2016;27(5):806-12.
6. Gray R (EBCTCG) et al. Extended aromatase inhibitor treatment following 5 or more years of endocrine therapy: a metaanalysis of 22192 women in 11 randomised trials. SABCS 2018;GS3-03
7. Zackariah C, Kollias J, Bingham J et al. Extended duration of adjuvant aromatase inhibitor in breast cancer: a meta-analysis of randomized controlled trials. Gland Surg 2018;7(5):449-457.
8. Burstein HJ, Lacchetti C, Anderson H et al. Adjuvant endocrine therapy for women with hormone receptor-positive breast cancer: ASCO clinical practice guideline focused update. J Clin Oncol. 2018 Nov 19;JCO1801160. doi: 10.1200/JCO.18.01160

Interruption of endocrine treatment up to 3 months during EAT:

1. Colleoni M, Luo W, Karlsson P et al. Extended adjuvant intermittent letrozole versus continuous letrozole in postmenopausal women with breast cancer (SOLE): a multicentre, open-label, randomised, phase 3 trial. Lancet Oncol 2018; 19: 127–38.



1. Gray R (EBCTCG) et al. Extended aromatase inhibitor treatment following 5 or more years of endocrine therapy: a metaanalysis of 22192 women in 11 randomised trials. SABCS 2018;GS3-03



© AGO e. V.
in der DGOG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2020.1D

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

Entscheidungskriterien für die erweiterte Adjuvanz

Kriterien, die auf einen klinischen Benefit hinweisen:

- Alleinige adjuvante Therapie mit Tamoxifen
- Z. n. Chemotherapie (höheres Risiko)
- Positiver Lymphknotenstatus
- T2/T3-Tumoren
- hohes Rückfall-Risiko nach immunhistochemischen Kriterien oder Multi-Gen Assays
- Erhöhter CTS5-Score

Weitere Entscheidungsfaktoren:

- Patientenwunsch
- bisherige gute Verträglichkeit der AI-Therapie bzw. Nebenwirkungen
- Knochengesundheit
- jüngeres Alter
- Adhärenz

1. Gnant M, G Steger, R Greil, et al. A prospective randomized multi-center phase-III trial of additional 2 versus additional 5 years of Anastrozole after initial 5 years of adjuvant endocrine therapy - results from 3,484 postmenopausal women in the ABCSG-16 trial. SABCS 2017; GS3-01
2. Li L, Chang B, Jiang X et al. Clinical outcomes comparison of 10 years versus 5 years of adjuvant endocrine therapy in patients with early breast cancer. Clinical outcomes comparison of 10 years versus 5 years of adjuvant endocrine therapy in patients with early breast cancer. BMC Cancer 2018;18:977
3. Goldvaser H, Barnes TA, Šeruga B, et al. Toxicity of extended adjuvant therapy with aromatase inhibitors in early breast cancer: a systematic review and meta-analysis. J Natl Cancer Inst 2018;110(1)djx141
4. van Hellemond I, Geurts SME, Tjan-Heijnen VCG: Current status of extended adjuvant endocrine therapy in early stage breast cancer. Curr Treat Options in Oncol 2018;19:26.
5. Pan H, Gray R, Braybrooke J et al. 20-year risks of breast recurrence after stopping endocrine therapy at 5 years. N Engl J Med 2017;1836-49.
6. Munzone E, Colleoni M: Optimal management of luminal breast cancer: how much endocrine therapy is long enough? Ther Adv Med Oncol 2018;10: 1–11.
7. Dowsett M, Sestak I, Regan MM et al. Integration of clinical variables for the prediction of late distant recurrence in patients with estrogen receptor–positive breast cancer treated with 5 years of endocrine therapy: CTS5. J Clin Oncol 2018 : 36(19): 1941–1948.

8. Sestak I, Buus R, Cuzick J et al. Comparison of the performance of 6 prognostic signatures for estrogen receptor–positive breast cancer: a secondary analysis of a randomized clinical trial. *JAMA Oncol* 2018; 4(4): 545–553.
9. Burstein HJ, Lacchetti C, Anderson H et al. Adjuvant endocrine therapy for women with hormone receptor–positive breast cancer: ASCO clinical practice guideline focused update. *J Clin Oncol*. 2018 Nov 19;JCO1801160. doi: 10.1200/JCO.18.01160
10. Bartlett J, SgROI DTrans-aTTom: Breast Cancer Index predicts benefit of extended endocrine therapy in HR+ breast cancers treated in the adjuvant tamoxifen-to offer meore (aTTom) trial Abstract 505 ASCO 2019

	Oxford		
	LoE	GR	AGO
Prophylaxe des ovariellen Funktionsausfalls und Fertilitätserhaltung bei prämenopausalen Patientinnen mit (neo-)adjuvanter Chemotherapie (CT)			
■ Angebot zur Beratung über Fertilitätserhaltung inkl. assist. reprod. Therapie (Information: www.fertiprotekt.com)			++
■ CHT + GnRHa (zur Prophylaxe des ovariellen Funktionsausfalls) (GnRHa Applikation > 2 Wochen vor Chemotherapie, unabhängig vom Hormonrezeptorstatus)	1a	A	+
■ CHT + GnRHa (zur Erhöhung der Schwangerschaftsrate)	1b	A	+/-

© AGO e. V.
 in der DGGG e.V.
 sowie
 in der DKG e.V.
 Guidelines Breast
 Version 2020.1D

www.ago-online.de
 FORSCHEN
 LEHREN
 HEILEN

Ovarian function protection

1. Gerber B, von Minckwitz G, Stehle H et al.: Effect of luteinizing hormone-releasing hormone agonist on ovarian function after modern adjuvant breast cancer chemotherapy: the GBG 37 ZORO study. J Clin Oncol. 2011 Jun 10;29(17):2334-41.
2. Del Mastro L, Ceppi M, Poggio F et al.: Gonadotropin-releasing hormone analogues for the prevention of chemotherapy-induced premature ovarian failure in cancer women: systematic review and meta-analysis of randomized trials. Cancer Treat Rev. 2014 Jun;40(5):675-83.
3. Del Mastro L, Rossi G, Lambertini M et al.: New insights on the role of luteinizing hormone releasing hormone agonists in premenopausal early breast cancer patients. Cancer Treat Rev. 2016 Jan;42:18-23.
4. Munholz RR, et al: Gonadotropin-Releaseing hormone agonists for ovarian function preservation in premenopausal women undergoing chemotherapy for early stage breast cancer- A systematic Review and Meta Analysis. JAMA Oncol 2016;2:65-73
5. Munster PN, Moore AP, Ismail-Khan R et al.: Randomized Trial Using Gonadotropin-Releasing Hormone Agonist Triptorelin for the Preservation of Ovarian Function During (Neo)Adjuvant Chemotherapy for Breast Cancer. J Clin Oncol. 2012;30(5):533–8.
6. Lambertini M, Boni L, Michelotti A et al.: Ovarian Suppression With Triptorelin During Adjuvant Breast Cancer Chemotherapy and Long-term Ovarian Function, Pregnancies, and Disease-Free Survival: A Randomized Clinical Trial. JAMA. 2015 Dec 22-

29;314(24):2632-40. doi: 10.1001/jama.2015.17291.

7. Lambertini M, Moore HCF, Leonard RCF et al.: Gonadotropin-Releasing Hormone Agonists During Chemotherapy for Preservation of Ovarian Function and Fertility in Premenopausal Patients With Early Breast Cancer: A Systematic Review and Meta-Analysis of Individual Patient-Level Data. *J Clin Oncol* 2018, 36:1981-90
8. Elgindy E, Sibai H, Abdelghani A et al.: Protecting Ovaries During Chemotherapy Through Gonad Suppression: A Systematic Review and Meta-analysis. *Obstet Gynecol.* 2015;126(1):187–95.
9. Sun X, Dongol S, Jiang J et al.: Protection of ovarian function by GnRH agonists during chemotherapy: a meta-analysis. *Int J Oncol.* 2014;44(4):1335–40.

Pregnancy rates


1. Lambertini M, Ceppi M, Poggio F et al.: Ovarian suppression using luteinizing hormone-releasing hormone agonists during chemotherapy to preserve ovarian function and fertility of breast cancer patients: a meta-analysis of randomized studies. *Ann Oncol* 2015; 26(12):2408-19.
2. Moore HCF, Unger JM, Phillips K-A et al. Goserelin for ovarian protection during breast-cancer adjuvant chemotherapy. *N Engl J Med.* 2015;372(10):923–32.
3. Lambertini M, Boni L, Michelotti A et al. Ovarian suppression with triptorelin during adjuvant breast cancer chemotherapy and long-term ovarian function, pregnancies, and disease-free survival. A randomized clinical trial. *JAMA.* 2015;314(24):2632-40.

Fertility preservation counselling


1. Loren AW, Mangu PB, Beck LN et al. Fertility Preservation for Patients With Cancer: American Society of Clinical Oncology Clinical Practice Guideline Update. *J Clin Oncol.* 2013;31(19):2500–10.
2. Peccatori FA, Azim Jr HA, Orecchia R et al. Cancer, pregnancy and fertility: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol.* 2013;24 Suppl 6:vi160–70.
3. Abe A, Kuwahara A, Iwasa T et al.: A survey on fertility management in young women of reproductive age treated with chemotherapy. *Int J Clin Oncol.* 2016 Dec;21(6):1183-1190.

Fertility preservation with assisted reproduction therapy

1. Dittrich R, Hackl J, Lotz L et al.: Pregnancies and live births after 20 transplantations of cryopreserved ovarian tissue in a single center. *Fertil Steril*. 2015 Feb;103(2):462-8.
2. Gamzatova Z, Komlichenko E, Kostareva A et al.: Autotransplantation of cryopreserved ovarian tissue--effective method of fertility preservation in cancer patients. *Gynecol Endocrinol*. 2014 Oct;30 Suppl 1:43-7.
3. Goldrat O, Kroman N, Peccatori FA et al.: Pregnancy following breast cancer using assisted reproduction and its effect on long-term outcome. *Eur J Cancer*. 2015;51(12):1490–6.
4. Lambertini M, Fontana V, Massarotti C et al.: Prospective study to optimize care and improve knowledge on ovarian function and/or fertility preservation in young breast cancer patients: Results of the pilot phase of the PREgnancy and FERTility (PREFER) study. *Breast* 2018, 41:51-6.
5. Luke B, Brown MB, Missmer SA et al.: Assisted reproductive technology use and outcomes among women with a history of cancer. *Hum Reprod*. 2016 Jan;31(1):183-9.
6. Oktay K, Turan V, Bedoschi G et al.: Fertility Preservation Success Subsequent to Concurrent Aromatase Inhibitor Treatment and Ovarian Stimulation in Women With Breast Cancer. *J Clin Oncol*. 2015;33(22):2424–9.



ARBEITSGEMEINSCHAFT
GYNAKOLOGISCHE
ONKOLOGIE e.V.



MAMMIA

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2020.1D

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

Gonadotropin-Releasing Hormone Agonists During Chemotherapy for Preservation of Ovarian Function and Fertility in Premenopausal Patients With Early Breast Cancer: A Systematic Review and Meta-Analysis of Individual Patient-Level Data

N = 837 patients from 5 trial, median follow-up time 5.0 years (IQR, 3.0–6.3 years)

	Control	GnRH	HR (95%-CI)	P-value
POI ^{1,2}	30.9%	14.1%	0.38; 0.26 to 0.57	< 0.001
Pregnancy ³	5.5%	10.3%	1.83; 1.06 to 3.15;	0.03

¹premature ovarian insufficiency, ² different definitions and time points were used

³in most trials POI and not pregnancy was defined as the primary endpoint

No significant differences in disease-free survival and overall survival were observed between groups.

Lambertini M et al. J Clin Oncol 2018

1. Lambertini M, Moore HCF, Leonard RCF et al. Gonadotropin-Releasing hormone agonists during chemotherapy for preservation of ovarian function and fertility in premenopausal patients with early breast cancer: A systematic review and meta-analysis of individual patient-level data. J Clin Oncol 2018;36(19):1981-1990. doi: 10.1200/JCO.2018.78.0858.

Adjuvante endokrine Therapie im Überblick

Studie	Therapien															De-facto- Vergleiche (Jahre)	HR für DFS	AI-Therapie Jahre 0-5 (%)
Jahre nach Diagnose	1	2	3	4	5	6	7	8	9	10	15							
Studien mit Tamoxifen nach 5 Jahren Tamoxifen																		
ATLAS					*											5 vs 10	0,75 – 0,99 †	0
ATTOM					*											5 vs 10	0,75 – 0,99 †	0
Studien mit AI nach 5 Jahren Tamoxifen																		
MA. 17					*											5 vs 10	0,57	0
NSABP B-33					*											5 vs 10	0,68	0
ABCSG 6a					*											5 vs 8	0,62	0
Studien mit erweiterter AI-Th. nach 5 Jahren endokrin inkl. AI																		
DATA				*												6 vs 9	0,79	100
NSABP B-42					*											5 vs 10	0,85	100
MA.17R											§					10 vs 15	0,66	100
Studien bzgl. optimaler Dauer in Jahr 5-10																		
BOOG 2006-05 IDEAL					*											7,5 vs 10	0,92	88
ABCSG 16					*											7 vs 10	1,007	49

Braun: Tamoxifen,

Grün: Tamoxifen oder AI,

Blau: AI

Gestreift: Zeit der
randomisierten
Intervention vs keine
Therapie od. Placebo,

*: Randomisierungs-
zeitpunkt,

§ : MA17R nach 5 Jahren
AI mit /ohne Tam zuvor