

Diagnosis and Treatment of Patients with early and advanced Breast Cancer

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

Adjuvant Cytotoxic and Targeted Therapy

Adjuvant Cytotoxic and Targeted Therapy

© AGO e. V.
in der DGGO e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

- **Versionen 2002 – 2020:**

**Dall / Fehm / Harbeck / Jackisch / Janni / Loibl / Lux/
von Minckwitz / Möbus / Müller / Nitz / Schmidt /
Schneeweiss / Simon / Schütz / Solomayer /
Stickeler / Thill / Thomssen / Untch**

- **Version 2021:**
Albert / Kümmel

Subtype-specific Strategies for Systemic Treatment

AGO

© AGO e. V.
in der DGGO e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

If chemotherapy is indicated systemic treatment before surgery (neoadjuvant) should be preferred

HR+/HER2- and „low-risk“

- Endocrine therapy without chemotherapy

++

HR+/HER2- and „high-risk“

- Conventionally dosed AT- based chemotherapy (q3w)
- Dose dense chemotherapy (including weekly schedule)
- Followed by endocrine therapy

+

++

++

HER2+

- Trastuzumab (plus Pertuzumab in N+ or NACT)
 - Sequential AT-based chemotherapy with concurrent T + anti-HER2 therapy
 - Anthracycline-free chemotherapy + anti-HER2 therapy

++

++

++

Triple-negative (TNBC)

- Conventionally-dosed AT-based chemotherapy
- Dose-dense chemotherapy (AT-based including weekly schedule)
- Neoadjuvant platinum-containing chemotherapy
- Neoadjuvant chemotherapy + ICPI (immune checkpoint-inhibitors)

+

++

+

+/-*

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

* Study participation recommended

Adjuvant Cytotoxic and Targeted Therapy

Adjuvant Chemotherapy: TNBC

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

■ Indication for chemotherapy in node-negative disease

- > 10 mm
- > 5–10 mm
- ≤ 5 mm

Oxford

LoE	GR	AGO
2b	B	++
2b	B	+
2b	B	-

Adjuvant Chemotherapy without Trastuzumab: Overview

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

	Oxford		
	LoE	GR	AGO
■ Dose-dense anthracycline / taxane based (incl. weekly) chemotherapy	1a	A	++
■ Conventional anthracycline-/taxane-based (q3w)	1a	A	+
■ „Tailored“ anthracycline-/taxane-based	1b	B	+/-
■ If anthracyclines cannot be given			
■ Docetaxel plus cyclophosphamide	1b	B	+
■ Paclitaxel mono weekly	1b	B	+/-
■ CMF	1a	A	+/-
■ Low-dose maintenance chemo	1b	B	-

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

Gray R et al., Lancet 2019

© AGO e. V.
in der DGGO e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

Early Breast Cancer Trialists' Cooperative Group (EBCTCG)

Increasing the dose-density of adjuvant chemotherapy: an EBCTCG meta-analysis

Same chemotherapy drugs and doses (**n = 10,004**)

Recurrence-free survival: 10-y Gain 4.3% (95%-C.I. 2.2 – 6.5)

(RR = 0.83; 95%-C.I. 0.76 – 0.91; p<0.0001)

Overall survival: 10-y Gain 2.8% (95%-C.I. 0.8 – 4.8)

(RR = 0.86; 95%-C.I. 0.77 – 0.96; p=0.0054)

ER negative: **10-y Gain 4.7%** (95%-C.I. 2.3 – 7.1)

ER positive: **10-y Gain 3.1%** (95%-C.I. 1.5 – 4.7)

Recommended Dose-dense and / or Dose-escalated, Sequential Adjuvant Chemotherapy

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

Dose-dense regimen

- $A_{60} \times 4 \rightarrow Pac_{175} \times 4 \rightarrow C_{600} \times 4 \text{ q2w}$
- $A_{60}C \text{ q2w } \times 4 \rightarrow Pac_{175} \text{ q2w } \times 4$
- $E_{90}C \text{ q2w } \times 4 \rightarrow Pac_{175} \text{ q2w } \times 4$
- $E_{90}C \text{ q2w } \times 4 \rightarrow Pac_{80} \text{ q1w } \times 12$
- $NabPac_{125} \times 8-12 \rightarrow E_{90}C \text{ q2(3)w } \times 4$

Dose-dense and dose-escalated regimen ($N \geq 4+$)

- $E_{150} \rightarrow Pac_{225} \rightarrow C_{2000} \text{ q2w}$

Oxford		
LoE	GR	AGO
1b	A	++
1b	B	++
1b	A	++
1b	B	++
1b	B	+
1b	A	++

Recommended Conventional Regimens for Adjuvant Chemotherapy

* Extrapolation from doxorubicin trials

Oxford

LoE GR AGO

Anthrazycline-/ taxane-based regimen

- *EC q3w x 4 → Pac q1w x 12
 - AC q3w x 4 → Pac q1w x 12
 - AC → D qw3
 - *EC → D qw3
 - DAC
- $A_{60}C$ q3w x 4 → D_{100} x 4
- $E_{90}C$ q3w x 4 → D_{100} x 4
- $D_{75}A_{50}C$ q3w x 6

2b	B	++
1b	A	++
1b	A	+
1b	B	+
1b	A	+

Anthrazycline-free regimen

- DC similar efficacy as EC → D
 - DC >> 4 x AC
 - Pac mono
 - CMF
- $D_{75}C_{600}$ x 6
- $D_{75}C_{600}$ x 4
- P_{80} q1w x 12

1b	B	+
1b	B	+
1b	B	+/-
1a	A	+/-

Taxane-free regimen (if pN0)

- $FE_{100}C$ x 6
- $F_{500}E_{100}C_{500}$ x 6

2b ^(a)	B	+
-------------------	---	---

Adjuvant Cytotoxic and Targeted Therapy

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

Adjuvant Chemotherapy

Other Drugs

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

- **Capecitabine-containing regimen in TNBC**
 - adjuvant/neoadjuvant
 - postneoadjuvant in non-pCR patients*
- **Platinum-containing regimen**
- **Anthracycline-free adjuvant therapy in TNBC**
- **Anthracycline-based adjuvant therapy in TNBC**
- **5- fluorouracile added to EC/AC**

Oxford		
LoE	GR	AGO
1a	B	+/-
1a ^a	A	+/-
1a ^a	A	+
1b	B	+
5	D	+/-
1b	A	--

*no platinum pretreatment

Van Mackelenbergh M et al., SABCS 2019, abstr. GS1-07

Effects of capecitabine as part of neo-/adjuvant chemotherapy

Meta-analysis of individual patient data from 12 randomized trials (n=15,457)

HR for DFS overall 0.952 (95%-C.I. 0.895-1.012, p=0.115)

X add. 0.888 (95%-C.I. 0.817-0.965, p=0.005)

X instead 1.035 (95%-C.I. 0.945-1.134, p=0.455)

HR for OS overall 0.892 (95%-C.I. 0.824-0.965, p=0.005)

X add. 0.837 (95%-C.I. 0.751-0.933, p=0.001)

X instead 0.957 (95%-C.I. 0.853-1.073, p=0.450)

Significance only for TNBC overall DFS 0.886 (95%-C.I. 0.789-0.994, p=0.040)

OS 0.828 (95%-C.I. 0.720-0.952, p=0.008)

X add.: DFS 0.818 (95%-C.I. 0.713-0.938, p=0.004)

OS 0.778 (95%-C.I. 0.657-0.921, p=0.004)

Adjuvant Treatment with Trastuzumab +/- Pertuzumab

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

Oxford		
LoE	GR	AGO
<hr/>		
1b ^a	B	+
1b ^a	B	+/-
1a	A	++
2b	B	+
2b	B	+/-

- **Trastuzumab + Pertuzumab**
 - pN+
 - pN-
- **Trastuzumab in node-negative disease
(if chemotherapy is indicated)**
 - > 10 mm
 - > 5–10 mm
 - ≤ 5 mm

Adjuvant treatment with Trastuzumab/Pertuzumab

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

Oxford		
LoE	GR	AGO

Start of treatment

■ Simultaneously with taxanes	1a	A	++
■ Sequentially up to 3 months after chemotherapy	1b	B	+
■ s.c. = i.v.	1a	A	++

Duration

■ For 1 year	1a	A	++
■ For 0.5 years (Trastuzumab)	1a	A	+
■ For 2 years	1b	A	-

Adjuvant Treatment with Trastuzumab +/- Pertuzumab: Chemotherapy regimen

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

Oxford		
LoE	GR	AGO
<hr/>		
1a	A	++
2b	B	+
1b	A	+
1b	B	++
1b	B	+
2b	B	+
2b	B	+

Trastuzumab simultaneously with

- paclitaxel / docetaxel after AC / EC
- P q1w 12 x in pT < 2 cm, pN0
- docetaxel and carboplatin

Trastuzumab + Pertuzumab simultaneously with

- paclitaxel q1w (or docetaxel q3w) after EC/AC
- docetaxel+ carboplatin
- taxanes dose-dense

Radiotherapy concurrently with Trastuzumab/Pertuzumab

Adjuvant Therapy With Other Targeted Agents

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

- **Lapatinib**
 - (delayed adjuvant treatment)
- **Lapatinib + Trastuzumab**
- **Neratinib* (one year) after completing a year of adjuvant trastuzumab (if HR-positive)**
- **Bevacizumab**

Oxford		
LoE	GR	AGO
1b ^a	B	-
1b	B	-
1b ^a	B	-
1b	B	+
1b	B	--

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

* In addition to standard endocrine treatment

Post-neoadjuvant therapy: HER2-negative

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

Oxford		
LoE	GR	AGO

HR positiv (pCR and non-pCR)

■ Endocrine therapy according to menopausal status (s. chap. 10)	1a	A	++
■ Capecitabine (in case of non-pCR)	3b	C	+/-
■ Endocrine therapy + Abemaciclib	2b	B	+/-*
■ Endocrine therapy + Palbociclib	1b ^a	B	-*

Triple negative (TNBC) (in case of non-pCR)

■ Capecitabine (up to 8 cycles)**	1b	B	+
■ Experimental post-neoadjuvant therapies within clinical trials	5	D	+*

* Study participation recommended

** without prior platinum-based therapy

Adjuvant Cytotoxic and Targeted Therapy

Post-neoadjuvant therapy: HER2-positive

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

pCR

- Low risk: Trastuzumab (to complete 12 mths)
- High risk (cN+): Trastuzumab + Pertuzumab (to complete 12 mths)
- Neratinib after 1 year Trastuzumab (HR-positive)*

non-pCR

- T-DM1
- Neratinib after 1 year* Trastuzumab (HR-positive)*
- Trastuzumab + Pertuzumab (to complete 12 mths)

	Oxford		
	LoE	GR	AGO
Low risk: Trastuzumab (to complete 12 mths)	2a	C	++
High risk (cN+): Trastuzumab + Pertuzumab (to complete 12 mths)	2b	C	+
Neratinib after 1 year Trastuzumab (HR-positive)*	2b	B	-
T-DM1	1b	B	+
Neratinib after 1 year* Trastuzumab (HR-positive)*	2b	B	+/-
Trastuzumab + Pertuzumab (to complete 12 mths)	2b	C	+/-

* In combination with standard endocrine treatment

Adjuvant Cytotoxic and Targeted Therapy

Biosimilars

General Considerations

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

Biosimilars that are used for treatment (i.e. trastuzumab) and supportive care of breast cancer (i.e G-CSF) must be approved by the respective regulatory authorities (EMA, FDA) after passing the stringent development and validation processes required before being used in daily practise.*