



Diagnosis and Treatment of Patients with early and advanced Breast Cancer

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

Adjuvant Radiotherapy

FORSCHEN
LEHREN
HEILEN

Adjuvant Radiotherapy (RT)

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

- **Versions 2002 – 2020:**
Blohmer / Budach / Friedrichs / Göhring / Huober/ Janni / Kühn / Möbus / Rody / Scharl / Seegenschmiedt / Souchon / Thomssen / Untch / Wenz
- **Version 2021:**
Budach / Friedrich / Krug



Preliminary Note

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

- **The recommendations on adjuvant radiotherapy for breast cancer are based on a consensus discussion between AGO and DEGRO experts**
- **For technical radiotherapy details, we refer to the corresponding updated DEGRO practice guidelines**

www.ago-online.de

**FORSCHEN
LEHREN
HEILEN**

Radiotherapy (RT) after Breast Conserving Surgery (Invasive Cancer): Whole Breast Irradiation

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

	Oxford		
	LoE	GR	AGO
▪ Radiotherapy of the affected breast	1a	A	++
▪ Hypofractionated radiotherapy (total dose approx. 40 Gy in 15-16 fractions within 3-5 weeks)	1a	A	++
▪ Conventionally fractionated radiotherapy (total dose about 50 Gy in approx. 25-28 fractions in 5-6 weeks)	1a	B	+
▪ Ultra-hypofractionated RT (total dose 26 or 28,5 Gy in 5 fractions in 1 or 5 weeks)	1b	B	+/-
▪ In case of life expectancy <10 years and pT1, pN0, R0, ER/PR-positive, HER2-negative, endocrine therapy (all criteria), radiotherapy can be omitted after individual counseling, resulting in an increased risk for in-breast recurrence	1a	B	+

FAST / FAST-Forward

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

	FAST	FAST Forward
Timeframe	2004-2007	2011-2014
Sample size	915	4096
Dosse/Fractionation	50 Gy/2 Gy/5 weeks 30 Gy/6 Gy/5 weeks 28,5 Gy/5,7 Gy/5 weeks	40 Gy/2,67 Gy/3 weeks 27 Gy/5,4 Gy/1 weeks 26 Gy/5,2 Gy/1 weeks
Median follow-up	119.8 months	71.5 months
Primary endpoint	change in photographic breast appearance	Ipsilateral breast tumor recurrence (non-inferiority margin 1,6%)
Inclusion criteria	pT1-2 (< 3 cm) pN0 Age ≥50 years Breast conserving surgery No chemotherapy	pT1-3 pN0-1 Age ≥18 years Breast-conserving surgery or mastectomy Approx. 25% adj. chemotherapy
Boost	No	Approx. 25%, 5-8x 2 Gy

www.ago-online.de

FAST / FAST-Forward

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

	FAST (10 year-data)			FAST Forward (5 year-data)		
	Dose	Frequency	Hazard ratio (95%-CI)	Dose	Frequency	Hazard ratio (95%-CI)
Ipsilateral in-breast recurrence	50 Gy	0.7%	-	40 Gy	2.1%	-
	30 Gy	1.4%	HR 1.36 (0.3-6.06)	27 Gy	1.7%	HR 0.86 (0.51-1.44)
	28.5 Gy	1.7%	HR 1.35 (0.3-6.05)	26 Gy	1.4%	HR 0.67 (0.38-1.16)
Moderate/marked normal tissue effects breast/chestwall	50 Gy	33.6%	-	40 Gy	26.8%	-
	30 Gy	50.4%	HR 1.79 (1.37-2.34)	27 Gy	35.1%	HR 1.41 (1.23-1.61)
	28.5 Gy	47.6%	HR 1.45 (1.10-1.91)	26 Gy	28.5%	HR 1.09 (0.95-1.27)

www.ago-online.de

BCS $\geq 70y$ $< 4cm$ cN0 : Tamoxifen vs. Tamoxifen + RT

Time: 1994-1999, since 8/1996 only pT1cN0 ER/PR+ or unknown allowed

@10 yrs (95% C.I.)	Tamoxifen	Tamoxifen plus Radiotherapy	Hazard Ratio
Local recurrence-free ($\Delta=8\%$)	90% (85%-93%)	98% (96%-99%)	HR=0.18 (95% CI, 0.07 to 0.42; P < .001)
Mastectomy-free	96% (93% - 98%)	98% (96% - 99%)	HR=0.50 (95% CI, 0.17 to 1.48; n.s.)
Distant metastasis-free	95% (91% - 97%)	95% (92% - 97%)	HR=1.20 (95% CI, 0.63 to 2.32; n.s)
Overall survival	66% (61% - 71%)	67% (62% - 72%)	HR=0.95 (95% CI, 0.77 to 1.18; n.s.)

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

Radiotherapy (RT) after Breast Conserving Surgery (Invasive Cancer) – Boost Irradiation

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

Oxford

LoE GR AGO

	LoE	GR	AGO
<ul style="list-style-type: none"> Boost-RT (improves local control, no survival benefit) <ul style="list-style-type: none"> Premenopausal Postmenopausal, if >T1*, G3, HER2-positive, triple negative, EIC (at least 1 factor) 	1b	B	++
	2b	B	+
<ul style="list-style-type: none"> Techniques <ul style="list-style-type: none"> Percutaneous boost (photons, electrons) as sequential boost Multicatheter brachytherapy-boost Percutaneous boost as simultaneous integrated boost (with conventionally fractionated whole-breast irradiation) Percutaneous boost as simultaneous integrated boost (with hypofractionated whole-breast irradiation) Intraoperative boost irradiation (followed by whole-breast irradiation) 	1a	A	++
	1a	A	++
	1b	B	+
	2b	B	+/-
	2b	B	+

* continuous parameter with regard to risk of relapse

EORTC 22881-10882: Boost vs no Boost (Endpoint: Ipsilateral Breast Recurrence)

@20 yrs (95% C.I.)	Boost (n=2.661)	No boost (n=2.657)	Hazard Ratio (95% C.I.)
<u>Overall Survival</u> (Δ =-1.4%)	59.7% (56.3–63.0)	61.1% (57.6–64.3)	HR 1.05 (0.92–1.19) n.s.
<u>Cumulative Risk of Ipsilateral Breast Tumour Recurrence</u>			
All patients	12.0% (9.8–14.4)	16.4% (14.1–18.8)	HR=0.65 (0.52–0.81); p<0.0001
≤40 years (Δ =11.6%)	24.4% (14.9–33.8)	36.0% (25.8–46.2)	HR=0.56 (0.34–0.92); p=0.003
41–50 years (Δ =5.9%)	13.5% (9.5–17.5)	19.4% (14.7–24.1%)	HR=0.66 (0.45–0.98); p=0.007
51–60 years (Δ =2.96%)	10.3% (6.3–14.3)	13.2% (9.8–16.7)	HR=0.69 (0.46–1.04); p=0.020
>60 years (Δ =3.0%)	9.7% (5.0–14.4)	12.7% (7.4–18.0)	HR=0.66 (0.42–1.04); p=0.019

(Median F/U 17.2 y)

acc. to: Bartelink et al. Lancet Oncol 2015; 16: 47–56

EORTC 22881-10882: Boost vs no Boost (Endpoint: Any First Recurrence)

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

@15 yrs/20 yrs (95% C.I.)	Boost (n=2.661)		No boost (n=2.657)	Hazard Ratio (95% C.I.)
Overall Survival (Δ = - 1.4%)	59.7% (56.3–63.0)		61.1% (57.6–64.3)	HR 1.05 (0.92–1.19) n.s.
Cumulative Risk of Any First Recurrence				
All patients ($\Delta \geq 4\%$)	@15y @20y	28.1% 32,8%	32.1% 38.7%	HR=0.92 (0.81-1.04), n.s.
≤ 40 years ($\Delta > 6\%$)	@15y @20y	41.5% 49.5%	48.1% 56.8%	HR=0.80 (0.56-1.15), n.s.
41–50 years	@15y @20y	34.0% 38.6%	35.6% 44.2%	HR=0.91 (0.71-1.16), n.s.
51–60 years	@15y @20y	28.5% 34.7%	28.7% 36.2%	HR=0.96 (0.76-1.21), n.s.
>60 years	@15y @20y	27.4% 32.1%	29.1% 32.8%	HR=0.94 (0.74-1.19), n.s.

(Median F/U 17.2 y)

acc. Bartelink et al. Lancet Oncol 2015; 16: 47–56. Suppl.

Radiotherapy (RT) after Breast Conserving Surgery (Invasive Cancer) – Partial Breast Irradiation (PBI)

Oxford

LoE GR AGO

■ Intraoperative Radiotherapy (low-risk)*

■ As sole radiotherapy, during first breast surgery (IORT 50 kV, IOERT)

- >50 years
- >70 years

1b A +/-

1b A +

■ Postoperative partial breast irradiation (low-risk)*

- Interstitial Multicatheter-Brachytherapy
- Intracavitary balloon-technique
- Intensity-modulated radiotherapy (IMRT) (5x6 Gy in 2 weeks)
- 3D-conformal radiotherapy (15x2.67 Gy in 3 weeks)
- 3D-conformal radiotherapy (10x3.8 Gy in 2 weeks)
- 3D-conformal radiotherapy (10x3.85 Gy in 1 week)

1b A +

2b B -

1b A +

1b A +

2b B +/-

1b A +/-

For definition of target volume and practical conduct see DEGRO practice guidelines

* only for pT1 pN0 R0 G1-2, HR+, non-lobular, >50 years, no extensive DCIS

Data on partial breast irradiation

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

NSABP B-39/RTOG 0413 (Vicini FA et al. *Lancet*. 2019 Dec 14;394(10215):2155-2164.)

- Randomised phase III equivalence trial, 4216 pat., 2005-2013, DCIS or invasive carcinoma \leq 3 cm, 0-3 involved lymph nodes, age >18 y
- 50 Gy/25 fr. +/- boost vs. APBI with
 - 38.5 Gy/10 fr. in one week (external beam irradiation)
 - 34 Gy/10 fr. in one week (Multicatheter- or Single lumen-Brachytherapy)
- **“We observed an HR of 1.22 with a 90% CI of 0.94–1.58, which did not meet the equivalence criteria and favoured whole-breast irradiation. The 10-year cumulative incidence of IBTR was 3.9% (95% CI 3.1–5.0) in the whole-breast irradiation group and 4.6% (3.7–5.7) in the APBI group for an absolute difference of 0.7%.”**
- **“Significantly more evaluable patients in the APBI group had recurrence-free interval events than patients in the whole-breast irradiation group (figure 3). The 10-year point estimate of recurrence-free interval for the whole breast irradiation group was 93.4% (95% CI 92.1–94.6), and in the APBI group it was 91.8% (90.4–93.0; figure 3)”.**
- **“Our findings support whole-breast irradiation but the absolute outcome difference compared with APBI is small, so partial breast irradiation might also be an acceptable treatment for some patients. “**

Data on partial breast irradiation

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

RAPID (Whelan TJ et al. Lancet. 2019 Dec 14;394(10215):2165-2172.)

- Randomised phase III non-inferiority trial, 2135 pat., 2006-2011, DCIS or invasive carcinoma \leq 3 cm, pN0, age \geq 40 y., no ILC
- 42.56/16 fr. or 50 Gy/25 fr. +/- Boost vs. APBI 38.5 Gy/10 fr. in one week (external beam irradiation)
- “In patients treated with **APBI**, the **5 year cumulative rate of IBTR was 2.3%** (95% CI 1.4–3.2) and the **8 year cumulative rate was 3.0%** (1.9–4.0). In patients treated with **whole breast irradiation**, the **5 year cumulative rate of IBTR was 1.7%** (0.9–2.5) and the **8 year cumulative rate was 2.8%** (1.8–3.9; figure 2). The HR for APBI versus whole breast irradiation was 1.27 (90% CI 0.84–1.91). Thus, **the upper bound of the estimated 90% CI did not exceed the non-inferiority margin of 2.02.**”
- “**Late radiation toxicity (grade \geq 2 [...]) was more common in patients treated with APBI** (346 [32%] of 1070 patients) than whole breast irradiation (142 [13%] of 1065 patients; $p < 0.0001$). **Adverse cosmesis [...]** was more common in patients treated with APBI than in those treated by whole breast irradiation at 3 years (absolute difference, 11.3%, 95% CI 7.5–15.0), 5 years (16.5%, 12.5–20.4), and 7 years (17.7%, 12.9–22.3).”

Postmastectomy Radiotherapy (PMRT)* to the Chest Wall

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

- **> 3 tumor infiltrated lymph nodes (LN)**
- **1–3 tumor infiltrated LN (high-risk)**
- **1–3 tumor infiltrated LN (low-risk*)**
- **T3 / T4**
 - pT3 pN0 R0 (and no additional risk factors)
- **If R0 is impossible to reach (for invasive tumor)**
- **In young pts with high-risk features**

The indications for PMRT and regional RT are independent of adjuvant systemic treatment

Oxford		
LoE	GR	AGO
1a	A	++
1a	A	+
5	D	+/-
1a	A	++
2b	B	+/-
1a	A	++
2b	B	++
1a	A	

* For definition of low-risk, see next slide **Radiotherapy of the Chest Wall After Mastectomy (PMRT)**

Radiotherapy of the Chest Wall After Mastectomy (PMRT) in Case of 1-3 Axillary Lymph Node Metastases

© AGO e. V.
 in der DGGG e.V.
 sowie
 in der DKG e.V.
 Guidelines Breast
 Version 2021.1E

PMRT can be omitted LoE 3b B AGO +	PMRT to be discussed LoE 3b B AGO +/-	PMRT recommended LoE 3b B AGO +
<div style="background-color: #008000; color: white; padding: 5px; border: 1px solid #000;"> ER pos, G1, HER2 neg, pT1 (at least 3 criteria present) </div>	<div style="border: 1px solid #008000; padding: 10px;"> <p>Patients, who don't fulfill the mentioned criteria for high or low risk</p> </div>	<div style="border: 1px solid #000; padding: 5px;"> <p>≥45 y. AND >25% pos. ax. Lnn in case of axillary dissection OR <45 y. AND (ER neg. OR >25% pos. ax. Lnn in case of axillary dissection OR medial tumor location)</p> </div>
<p>Kyndi et al. 2009</p>		<div style="border: 1px solid #000; padding: 5px;"> <p><40 y. OR HER2 pos. OR lymphovascular invasion</p> </div>
		<div style="border: 1px solid #000; padding: 5px;"> <p>G3 OR lymphovascular invasion OR triple negative</p> </div>

Truong et al. 2005

Shen H et al. 2015

Different publications

Comment: In case of an indication for radiotherapy of regional lymph nodes, radiotherapy of the chest wall should also be administered

Boost in PMRT

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

- **An additional boost irradiation to a part of the chest wall has not been shown to improve DSS and overall survival**
- **An additional boost irradiation to a part of the chest wall should be given in case of of R1/R2-resection, if secondary resection is not feasible**
- **In case of tumor extention to the pectoral resection margin, but no clinical signs of extention beyond the fascia, the resection margin should be regarded as R0 (provided, that the pectoral fascia was resected). A boost radiotherapy is not required in this situation**

Oxford		
LoE	GR	AGO
2a	B	
5	D	++
5	D	++

Radiotherapy of axillary lymph nodes in patients with positive sentinel-lymph nodes**, who did not undergo axillary dissection



© AGO e. V.
 in der DGGG e.V.
 sowie
 in der DKG e.V.
 Guidelines Breast
 Version 2021.1E

	Oxford		
	LoE	GR	AGO
BCS and ACOSOG Z0011-criteria⁺ met	2b	B	+*
<ul style="list-style-type: none"> Radiotherapy of the breast including LN level 1 + 2 to 5 mm below the axillary vein (PTV) 			
BCS and ACOSOG Z0011-criteria⁺ <u>not</u> met	1b	B	++*
<ul style="list-style-type: none"> Radiotherapy of the axillary lymph nodes (analog AMAROS) 			
ME and chest wall RT indicated and ACOSOG Z011-criteria⁺ <u>not</u> met or ME and chest wall RT <u>not planned</u>			
<ul style="list-style-type: none"> Radiotherapy of the axillary lymph nodes (analog AMAROS) 	1b	B	++
<u>>=3 pos. SLN</u>			
<ul style="list-style-type: none"> Radiotherapy of the axillary lymph nodes (analog AMAROS) 	1b	B	+

* Study participation recommended
 ** Macrometastases
 + <T3, no palpable LN, R0, 1-2 positive SN, no extracapsular extension, no NACT
 Adjuvant Radiotherapy

<u>Additional RT of the axilla after primary surgery</u>	Oxford		
	LoE	GR	AGO
(in case of an indication for RT of the breast/chest wall¹ +/- supra-/infraclavicular and internal mammary node RT²)			
Expansion of the PTV (planning target volume) to level I-II³			
pN-status			
pN0(sn)/pN1mic(sn)	1b	B	--
pN0/+ after ALND	1a	A	--
pN+(sn) in analogy to ACOSOG Z0011 (no ALND)	2b	B	+
pN+(sn) not fitting ACOSOG Z0011-criteria → RT in analogy to AMAROS⁴ (no ALND)	1b	B	++
R2-situation in the axilla	5	D	++

¹Incidental dose to parts of level i/II is inevitable. ²The indication for supra-/infraclavicular and internal mammary node RT has to be assessed separately. ³Cranial border 5 mm below the axillary vein. ⁴ ≤ T3, cN0, no NACT, always in conjunction with supra-/infraclavicular RT

Additional RT of the axilla after neoadjuvant therapy

(in case of an indication for RT of the breast/chest wall¹ +/- supra-/infraclavicular and internal mammary node RT²)

Expansion of the PTV (planning target volume) to level I-II³

Oxford

LoE

GR

AGO


		LoE	GR	AGO
N-status pre/post NACT	pN-status			
cN0 / ycN0	ypN0(sn)	5	D	-
cN0 / ycN0	ypN1mic(sn)/ypN+(sn) (no ALND)	5	D	+⁴
pN⁺_{CNB} / ycN0	ypN0(sn/TAD)	5	D	+/-⁴
pN⁺_{CNB} / ycN0	ypN1mic(sn/TAD)/ypN+(sn/TAD) (no ALND)	5	D	+⁴
cN0/cN+	ypN0/+ after ALND	2b	B	-
	R2-situation in the axilla	5	D	++

¹Incidental dose to parts of level i/II is inevitable. ²The indication for supra-/infraclavicular and internal mammary node RT has to be assessed separately. ³Cranial border 5 mm below the axillary vein. ⁴Study participation recommended.

Dose in the axillary LN-levels I + II using different RT-techniques

ACOSOG Z0011 Trial
45% micrometast. in the exp. arm

RT-volume
% of patientis



LN level 1	mean dose*	encompassed volume**
AMAROS	>95%	>95%
high tangent	86%	79%
standard tangent	66%	51%
IMRT ⁺	29%	1%
LN-level 2		
AMAROS	>95%	>95%
high tangent	71%	51%
standard tangent	44%	26%
IMRT ⁺	7%	0%

* in relation to the prescribed dose in the breast
 ** % volume receiving the prescribed dose
 + Lee et al. Medicine 2016 (3)

Data from 228/856 pat.

Jagsi (2): “The results of Z0011 should not be extrapolated to patients who receive RT using partial-breast or prone techniques, in which substantially less of the axilla is included”

Radiotherapy (RT) of Other Locoregional Lymph Node Areas (SCG/ICG)



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

RT to supra-/infraclavicular lymphatic regions

- ≥ 4 positive axillary lymph nodes (LN) or involved LN in level III or in supra-/infraclavicular LN
- 1–3 positive axillary lymph nodes¹ in case of
 - central or medial tumor and G2-3 or ER/PR-negative
 - premenopausal patient and G2-3 or ER/PR-negative
- pN0 with central or medial tumors, if premenopausal and G2-3 and ER/PR-negative

Oxford		
LoE	GR	AGO
1b	A	++
2a	B	+
2a	B	+/-

¹ not applicable for micrometastases

Radiotherapy (RT) of Other Locoregional Lymph Node Areas (IMN)



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

Internal mammary lymph node region (IMN)

- pN0 high-risk with central or medial tumor
and premenopausal and G2-3 and ER/PR-negative
- 1–3 positive axillary lymph nodes¹ in case of
 - central or medial tumor and G2-3 or ER/PR-negative
 - premenopausal patient and G2-3 or ER/PR-negative
- ≥ 4 positive axillary lymph nodes
- involved internal mammary lymph nodes
- In case of cardiac risk factors or if trastuzumab is given

	Oxford		
	LoE	GR	AGO
1b	B	+/-	
2a	B	+	
2a	B	+	
2a	B	+	
2b	A	--	

¹ not applicable for micrometastases

Fractionation of Radiotherapy in Case of Regional Nodal Irradiation

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

- **Conventionally fractionated radiotherapy**
(total dose about 50 Gy in approx. 25-28 fractions within 5–6 weeks)
- **Hypofractionated radiotherapy**
(total dose approx. 40–43.5 Gy in 15-16 fractions within 3–5 weeks)

Oxford		
LoE	GR	AGO
1a	A	++
2b	B	+/-

Hypofractionated post-mastectomy radiotherapy and regional nodal irradiation



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

Wang et al. Lancet Oncol. 2019 Mar;20(3):352-360.

- Randomised phase III non-inferiority trial, 820 pat., 2008-2016, T3/4 and/or ≥ 4 involved lymph nodes, 50 Gy/25 fr. vs. 43.5 Gy/15 fr.
- 98% 2D-planned radiotherapy, no treatment of the internal mammary lymph nodes
- “The 5-year cumulative incidence of locoregional recurrence was 8.3% (90% CI 5.8–10.7) in the hypo- fractionated radiotherapy group compared with 8.1% (90% CI 5.4–10.6) in the conventional fractionated radiotherapy group (absolute difference 0.2%, 90% CI –3.0 to 2.6; HR 1.10, 90% CI 0.72 to 1.69; figure 2).
- ”In conclusion, this study provides high-level evidence for the clinical use of hypofractionated postmastectomy radiotherapy for patients with high-risk breast cancer. It can be recommended in clinical practice to patients who do not plan breast reconstruction and will not receive internal mammary node irradiation.”

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

Multivariate Analysis of Overall Survival: Effect of Radiotherapy of the Internal Mammaria Lymph Nodes

(median follow-up 10.9 yrs)

Adjuvant treatment	n*	Hazard ratio (95%CI)
No adjuvant reported	625	0.91 (0.59 - 1.39)
Chemotherapy	954	1.05 (0.84 - 1.32)
Endocrine therapy	1185	0.82 (0.63 - 1.06)
Both (endocrine th. and chemotherapy)	1200	0.72 (0.55 - 0.94)
Total	4004	0.88 (0.76 - 1.01)

* missing data on 40 patients

Poortmans et al. ECCO Amsterdam 2013

Adjuvant Radiotherapy

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

Radiotherapy following NACT

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

Pretreatment	Posttreatment	RT-BCS	PMRT	RT-RN
Locally advanced	pCR / no pCR	yes	yes	yes
cT1/2 cN1+*	ypT1+ o. ypN1 + (no pCR)	yes	yes	yes
cT1/2 cN1+*	ypT0/is ypN0	yes	Increased risk of relapse ¹	
cT1/2 cN0 (Sonogr.bligmat)	ypT0/is ypN0	Ja	nein	nein

Oxford		
LoE	GR	AGO
1a/1a/1a	A/A/A	++/+/++
1a/2b/2b	A/B/B	++/+/+
2b/2b/2b	B/B/B	+/+/+
2b/2b/2b	A/B/B	+/-/-

Locally advanced: T3-4 or cN2-N3,

¹ Criteria for increased risk of relapse:

- pN0 premenopausal high risk: central or medium tumor localization, and (G2-3 and ER/PR-neg.)
- pretreatment pN1a/ cN+* high risk: central or medium tumor localization and (G2-3 or ER/PR-neg.) or premenopausal, lateral tumor localization and (G2-3 or ER/PR-neg.)

* Regarding coverage of axilla level I/II please also see slides „Additional RT of the axilla after primary surgery“ and „Additional RT of the axilla after neoadjuvant therapy“

** = confirmed by core biopsy

Molecular predictors and use of radiotherapy

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

- Results of gene expression profiling should not be used for indication of radiotherapy

Oxford		
LoE	GR	AGO
2b	B	++

Use of concomitant systemic therapy with adjuvant locoregional radiotherapy

Oxford

- Trastuzumab/Pertuzumab*
- T-DM1
- Tamoxifen
- Aromatase inhibitors
- Checkpoint inhibitors
- Capecitabine
- CDK4/6-inhibitors

LoE	GR	AGO
1a	A	++
1b	A	+
2b	B	+
2b	B	+
2b	C	+
2b	B	+**
4	C	+/-***

* concurrent Trastuzumab/Pertuzumab and parasternal radiotherapy should be avoided

** with hypofractionated RT approx. 40 Gy, consider dose reduction of Capecitabine, Pat. at high-risk for locoregional recurrence

*** In currently available phase III-trials (monarchE, PALLAS, Penelope-B) RT was given before initiation of CDK4/6-inhibitors. No definitive signs of significantly increased toxicity with concomitant RT in the palliative setting.

Adjuvant Radiotherapy



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

Simultaneous Capecitabine with locoregional radiotherapy

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

Woodward et al. Int J Radiat Oncol Biol Phys. 2017 Nov 15;99(4):777-783

- Prospective phase trial, 32 pat. with LABC, sim. def./neoadj. chemoradiotherapy, median total dose 66 Gy
- “The first 9 patients analyzed [...] received CAP 825 mg/m² twice daily continuously beginning on the first day of RT. **Because of observed excess grade 3 toxicity the protocol was amended,** and subsequent patients received CAP only on RT days (5 days per week).”
- “Noncontinuous CAP dosing was much better tolerated than continuous dosing. **Thirteen of 26 patients (50%) had grade ≥3 and higher treatment-related dermatologic toxicity.** “

Alhanafy et al. Menoufia Medical Journal 2015, 28:325-332

- Randomised phase II-trial, 100 pat., adj. Radiotherapy 40 Gy/15 fr. +/- CAP 825 mg/m² Mo-Fr, LABC
- “ [...] **concurrent capecitabine was feasible with a high percent of patients (96%),** [...] only two out of 50 (4%) patients had capecitabine dose modification ...”.
- “**All early toxicities were GI/GII.** Radiation dermatitis had a peak incidence in the last few fractions of the radiation therapy and the week after radiotherapy; no treatment interruption was needed and the incidence was close in both groups”.
- Radiation dermatitis grade I 14% vs. 18%; grade 2 4% vs. 4%

Smoking and Risk of secondary lung cancer

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1E

- Increased risk of lung cancer secondary to breast cancer radiotherapy in smokers
- Inform patients about risk
- Recommend to stop smoking

Oxford		
LoE	GR	AGO
1a	A	
		++
		++