



© AGO e. V.
in der DGGO e.V.
sowie
in der DKG e.V.


Guidelines Breast
Version 2023.1E

FORSCHEN
LEHREN
HEILEN

Diagnosis and Treatment of Patients with early and advanced Breast Cancer

Prognostic and Predictive Factors

Prognostic and Predictive Factors



- **Versions 2002–2022:**

Costa / Fasching / Fersis / Friedrichs / Gerber / Göhring / Harbeck /
Jackisch / Janni / Kolberg-Liedtke / Kreipe / Loibl / Lück / Mundhenke /
Nitz / Rody / Schaller / Schmidt / Schmutzler / Schneeweiss / Simon /
Solomayer / Thill / Thomssen / Untch / Witzel / Wöckel

- **Version 2023:**

Gluz / Witzel

Data bases screened

Pubmed 2008 - 2022, ASCO 2017-2022, SABCS 2003 – 2022, ESMO 2022, Cochrane data base (n.d.)


Definition

A **Prognostic Factors** is associated with the probability of the course of the disease (e.g. disease-free or progression-free survival, overall survival). The probability can be influenced by therapy.

A **Predictive Factor** is associated with the probability of the effect of a given therapy.

Definition of Prognosis and Prediction

1. Hayes DF, Bast RC, Desch CE et al.: Tumor marker utility grading system: a framework to evaluate clinical utility of tumor markers. J Natl Cancer Inst. 1996 Oct 16;88(20):1456-66.
2. McGuire WL, Clark GM. Prognostic factors and treatment decisions in axillary-node-negative breast cancer. N Engl J Med. 1992 Jun 25;326(26):1756-61.



© AGO e. V.
in der DGOG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2023.1E


www.ago-online.de

FORSCHEN
LEHREN
HEILEN

“Low absolute risk implies low absolute benefit”

Early Breast Cancer Trialists' Collaborative Group (EBCTCG), Lancet 379: 432-444, 2012

1. Early Breast Cancer Trialists' Collaborative Group (EBCTCG), Lancet 379: 432-444, 2012
2. Peto, R., Davies, C., Godwin, J., et al. 2012. Comparisons between different polychemotherapy regimens for early breast cancer: meta-analyses of long-term outcome among 100,000 women in 123 randomised trials. Lancet 379, 432–444.
3. Nielsen T, Jensen B, et al High risk premenopausal luminal A breast cancer patients derive no benefit from adjuvant chemotherapy: results from DBCG77B, SABCS 2015S1-08



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2023.1E

www.ago-online.de

FORSCHEN
LEBEN
HEILEN

Quality Criteria

- **Biological hypothesis**
- **Simple and standardized assessment method, quality assurance (QA) of the test**
- **Prospectively planned statistical evaluation (primary goal)**
- **Validation of clinical significance according to**
 - „Oxford Level of Evidence (LoEOx2001)“ criteria and „Grades of Recommendation (GR)“
 - „Grades of Recommendation (GR)“ as well as modified LoE criteria for the use in archived specimen (LoE2009) and category of tumor marker study (CTS)
- **Clinical relevance for treatment decisions**

1. Febbo PG, Ladanyi M, Aldape KD, et al. (2011) NCCN Task Force report: Evaluating the clinical utility of tumor markers in oncology. J Natl Compr Canc Netw 9 Suppl 5: S1-32; quiz S33.
2. Hayes DF, Bast RC, Desch CE et al. (1996) Tumor marker utility grading system: a framework to evaluate clinical utility of tumor markers. J. Natl. Cancer Inst. 88 (20): 1456–1466.
3. Jeremy Howick, Iain Chalmers, Paul Glasziou, et al. Explanation of the 2011 Oxford Centre for Evidence-Based Medicine (OCEBM) Levels of Evidence (Background Document). Oxford Centre for Evidence-Based Medicine.
4. McShane LM, Altman DG, Sauerbrei W et al. (2005) Reporting recommendations for tumor marker prognostic studies. J. Clin. Oncol. 23 (36): 9067–9072.
5. McShane LM, Hayes DF (2012) Publication of tumor marker research results: the necessity for complete and transparent reporting. J. Clin. Oncol. 30 (34): 4223–4232.
6. Simon RM, Paik S, Hayes DF (2009) Use of archived specimens in evaluation of prognostic and predictive biomarkers. J. Natl. Cancer Inst. 101 (21): 1446–1452.

Early Breast Cancer (M0) – eBC Prognostic Factors I

Factor	Oxford		
	LoE _{Ox2001}	GR	AGO
▪ Tumor size - pT	1a	A	++
▪ Axillary lymph node status - pN	1a	A	++
▪ Histological tumor type (mucinous, tubular etc.)	2b	B	++
▪ Grade (Elston & Ellis) - G	2a	B	++
▪ Age	2a	B	++
▪ Histologically proven peritumoral lymphatic vessel and vascular invasion (L1 V1)	1b	B	++
▪ pCR after NACT* in (luminal-B-like, HER2+, TN)	1a	A	++
▪ Increased risk of recurrence in invasive-lobular BC, cT3/4, N+	2a	B	+/-
▪ Obesity (BMI > 30 kg/m ²)	1b	B	+
▪ Margins (resection status) - R0 / R1	1a	A	+

* NACT = Neoadjuvant Chemotherapy

General references

1. Balic M, Thomssen C, Würtle R et al. St. Gallen/Vienna 2019: A Brief Summary of the Consensus Discussion on the Optimal Primary Breast Cancer Treatment. Breast Care (Basel). 2019 Apr;14(2):103-110.
2. Harris LN, Ismaila N, McShane LM et al.: Use of Biomarkers to Guide Decisions on Adjuvant Systemic Therapy for Women With Early-Stage Invasive Breast Cancer: American Society of Clinical Oncology Clinical Practice Guideline. J Clin Oncol. 2016 Apr 1;34(10):1134-50.

Tumor size

1. Coates AS, Winer EP, Goldhirsch A, et al. Tailoring therapies--improving the management of early breast cancer: St Gallen International Expert Consensus on the Primary Therapy of Early Breast Cancer 2015. Ann Oncol. 2015 Aug;26(8):1533-46.
2. Balic M, Thomssen C, Würtle R et al. St. Gallen/Vienna 2019: A Brief Summary of the Consensus Discussion on the Optimal Primary Breast Cancer Treatment. Breast Care (Basel). 2019 Apr;14(2):103-110.

Lymph node status

1. Coates AS, Winer EP, Goldhirsch A, et al. Tailoring therapies--improving the management of early breast cancer: St Gallen International Expert Consensus on the Primary Therapy of Early Breast Cancer 2015. Ann Oncol. 2015 Aug;26(8):1533-46.

2. Balic M, Thomssen C, Würstlein R et al. St. Gallen/Vienna 2019: A Brief Summary of the Consensus Discussion on the Optimal Primary Breast Cancer Treatment. *Breast Care (Basel)*. 2019 Apr;14(2):103-110.

Histological type (mucinous, tubular etc.)

1. Dieci MV, Orvieto E, Dominici M. Rare breast cancer subtypes: histological, molecular, and clinical peculiarities. *Oncologist*. 2014 Aug;19(8):805-13.
2. Horlings HM, Weigelt B, Anderson EM et al. Genomic profiling of histological special types of breast cancer. *Breast Cancer Res Treat*. 2013 Nov;142(2):257-69.
3. Weigelt B, Geyer FC, Reis-Filho JS. Histological types of breast cancer: how special are they? *Mol Oncol*. 2010 Jun;4(3):192-208.
4. Mouabbi JA, Hassan A, Lim B, Hortobagyi GN et al. *Invasive lobular carcinoma: an understudied emergent subtype of breast cancer. Breast Cancer Res Treat*. 2022 Jun;193(2):253-264. doi: 10.1007/s10549-022-06572-w. Epub 2022 Mar 26. PMID: 35347549.

Tumor grade (Elston & Ellis)

1. Balic M, Thomssen C, Würstlein R et al. St. Gallen/Vienna 2019: A Brief Summary of the Consensus Discussion on the Optimal Primary Breast Cancer Treatment. *Breast Care (Basel)*. 2019 Apr;14(2):103-110.
2. Coates AS, Winer EP, Goldhirsch A, et al. Tailoring therapies--improving the management of early breast cancer: St Gallen International Expert Consensus on the Primary Therapy of Early Breast Cancer 2015. *Ann Oncol*. 2015 Aug;26(8):1533-46.
3. Rakha EA, Aleskandarani M, Toss MS et al. Breast cancer histologic grading using digital microscopy: concordance and outcome association. *J Clin Pathol*. 2018 Aug;71(8):680-686.

Age

1. Brandt J, Garne JP, Tengrup I et al. Age at diagnosis in relation to survival following breast cancer: a cohort study. *World J Surg Oncol*. 2015 Feb 7;13:33.
2. Johnson HM, Irish W, Muzaffar M et al. Quantifying the relationship between age at diagnosis and breast cancer-specific mortality. *Breast Cancer Res Treat*. 2019 Oct;177(3):713-722
3. Liu YR, Jiang YZ, Yu KD et al. Different patterns in the prognostic value of age for breast cancer-specific mortality depending on hormone receptor status: a SEER population-based analysis. *Ann Surg Oncol*. 2015 Apr;22(4):1102-10.

Histologically proven lymph and/or blood vessel invasion

1. Ryu YJ, Kang SJ, Cho JS et al. Lymphovascular invasion can be better than pathologic complete response to predict prognosis in breast cancer treated with neoadjuvant chemotherapy. *Medicine (Baltimore)*. 2018 Jul;97(30):e11647

pCR after NACT* in Luminal B-like, HER2 and TN Breast Cancer

1. Cortazar P, Geyer CE Jr. Pathological complete response in neoadjuvant treatment of breast cancer. *Ann Surg Oncol*. 2015 May;22(5):1441-6.
2. Cortazar P, Zhang L, Untch M et al. Pathological complete response and long-term clinical benefit in breast cancer: the CTNeoBC pooled analysis. *Lancet*. 2014 Jul 12;384(9938):164-72.
3. Nekljudova V, Loibl S, von Minckwitz G et al. Trial-level prediction of long-term outcome based on pathologic complete response (pCR) after neoadjuvant chemotherapy for early-stage breast cancer (EBC). *Contemp Clin Trials*. 2018 Aug;71:194-198.

Increased risk of recurrence in invasive-lobular BC, cT3/4, N+

1. Thomas M, Kelly ED, Abraham J et al. Invasive lobular breast cancer: A review of pathogenesis, diagnosis, management, and future directions of early stage disease. *Semin Oncol*. 2019 Apr;46(2):121-132.
2. Van Baelen K, Geukens T, Maetens M et al. Current and future diagnostic and treatment strategies for patients with invasive lobular breast cancer. *Ann Oncol*. 2022 Aug;33(8):769-785. doi: 10.1016/j.annonc.2022.05.006. Epub 2022 May 21. Erratum in: *Ann Oncol*. 2022 Dec 16;; PMID: 35605746.

Obesity (BMI > 30 kg/m²)

1. Chan DSM et al. Body mass index and survival in women with breast cancer—systematic literature review and meta-analysis of 82 follow-up studies *Ann Oncol*. Oct 2014; 25(10): 1901–1914.
2. Houssami, N., et al., The association of surgical margins and local recurrence in women with early-stage invasive breast cancer treated with breast-conserving therapy: a meta-analysis. *Ann Surg Oncol*, 2014. 21(3): p. 717-30.

Resection status (R0 / R1)

1. Harris LN, Ismaila N, McShane LM et al.: Use of Biomarkers to Guide Decisions on Adjuvant Systemic Therapy for Women With Early-Stage Invasive Breast Cancer: American Society of Clinical Oncology Clinical Practice Guideline. *J Clin Oncol*. 2016 Apr 1;34(10):1134-50.
2. Febbo PG, Ladanyi M, Aldape KD, et al. (2011) NCCN Task Force report: Evaluating the clinical utility of tumor markers in oncology. *J*

Natl Compr Canc Netw 9 Suppl 5: S1-32; quiz S33.

Early Breast Cancer (M0) - eBC

Prognostic Factors II

Factor	Oxford		
	LoE	GR	AGO
▪ ER / PR	1a	A	++
▪ HER2 (IHC, ISH)	1a	A	++
▪ ER / PR / HER2/ Ki-67 to assess the intrinsic type with regards to tumor histology and biology	2b	B	++
▪ Proliferation markers			
▪ Ki-67 before, during, or after treatment	1a	B	+
▪ Ki-67 Re-Evaluation after short term preoperative endocrine therapy (2-4 weeks) (ypT and ypN)*	1a	B	+

* Biomarker and Multi Gene Expression test should be evaluated on core needle biopsy prior endocrine therapy

ER/PR

1. Allison KH, Hammond MEH, Dowsett M et al. Estrogen and Progesterone Receptor Testing in Breast Cancer: ASCO/CAP Guideline Update. J Clin Oncol. 2020 Jan 13;JCO1902309 (und: Arch Pathol Lab Med. 2020 Jan 13).
2. Jorns JM. Breast Cancer Biomarkers: Challenges in Routine Estrogen Receptor, Progesterone Receptor, and HER2/neu Evaluation. Arch Pathol Lab Med. 2019 Dec;143(12):1444-1449.

HER2


1. Ross, J.S., Slodkowska, E.A., Symmans, W.F., et al. 2009. The HER-2 receptor and breast cancer: ten years of targeted anti-HER-2 therapy and personalized medicine. Oncologist 14, 320–368.
2. Slamon, D.J., Clark, G.M., Wong, S.G. et al. 1987. Human breast cancer: correlation of relapse and survival with amplification of the HER-2/neu oncogene. Science 235, 177–182.
3. Wolff AC, Hammond ME, Hicks DG, et al.: Recommendations for human epidermal growth factor receptor 2 testing in breast cancer: American Society of Clinical Oncology/College of American Pathologists clinical practice guideline update. J Clin Oncol 2013, 31(31):3997-4013.

Ki-67

1. Denkert C, Budczies J, Regan MM, et al. Clinical and analytical validation of Ki-67 in 9069 patients from IBCSG VIII + IX, BIG1-98 and GeparTrio trial: systematic modulation of interobserver variance in a comprehensive in silico ring trial. *Breast Cancer Res Treat.* 2019 Aug;176(3):557-568. doi: 10.1007/s10549-018-05112-9. Epub 2019 May 7. PMID: 31065870.
2. Dowsett M, Ellis MJ, Dixon JM, et al. Evidence-based guidelines for managing patients with primary ER+ HER2- breast cancer deferred from surgery due to the COVID-19 pandemic. *Breast Cancer.* 2020 Jun 8;6:21.
3. Gluz O, Nitz UA, Christgen M, et al. Impact of age, recurrence score (RS) and ovarian function suppression (OFS) on endocrine response to short preoperative endocrine therapy (ET): Analysis of ADAPT and ADAPTCycle trials. *Annals of Oncology* (2022) 33 (suppl_7): S808-S869. 10.1016/annonc/annonc1089
4. Nielsen TO, Leung SCY, Rimm DL, et al. Assessment of Ki67 in Breast Cancer: Updated recommendations from the International Ki67 in Breast Cancer Working Group. *J Natl Cancer Inst.* 2020 Dec 28:djaa201.
5. Nitz U, Gluz O, Huober J et al. Final analysis of the prospective WSG-AGO EC-Doc versus FEC phase III trial in intermediate-risk (pN1) early breast cancer: efficacy and predictive value of Ki67 expression. *Ann Oncol.* 2017 Nov 1;28(11):2899.
6. Petrelli, F., et al., Prognostic value of different cut-off levels of Ki-67 in breast cancer: a systematic review and meta-analysis of 64,196 patients. *Breast Cancer Res Treat,* 2015. 153(3): p. 477-91.
7. Smith I, Robertson J, Kilburn L, et al. Long-term outcome and prognostic value of Ki67 after perioperative endocrine therapy in postmenopausal women with hormone-sensitive early breast cancer (POETIC): an open-label, multicentre, parallel-group, randomised, phase 3 trial. *Lancet Oncol.* 2020 Nov;21(11):1443-1454

Post-treatment Ki-67

1. Dowsett M, Smith IE, Ebbs SR, et al: Prognostic Value of Ki67 Expression After Short-Term Presurgical Endocrine Therapy for Primary Breast Cancer. *Journal of the National Cancer Institute* 99:167-170, 2007
2. DeCensi A, Guerrieri-Gonzaga A, Gandini S, et al: Prognostic significance of Ki-67 labeling index after short-term presurgical tamoxifen in women with ER-positive breast cancer. *Annals Oncol* 2011 Mar;22(3):582-7.
3. Gluz O, Nitz UA, Christgen M, et al. Impact of age, recurrence score (RS) and ovarian function suppression (OFS) on endocrine response to short preoperative endocrine therapy (ET): Analysis of ADAPT and ADAPTCycle trials. *Annals of Oncology* (2022) 33 (suppl_7): S808-S869. 10.1016/annonc/annonc1089



© AGO e. V.
in der DGOG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2023.1E

www.ago-online.de

FORSCHEN
LEBEN
HEILEN

Reproducibility – Quality Assurance is Key for Clinical Decision Making

- **ER / PR:** concordance central vs local is high (97%; Plan B, SABCS 2014)
- **Grade:** concordance central vs local is 68% (PlanB, JCO 2016)
- **HER2:** frequency of false-positive test results 6% (ASCO /CAP JCO 2013)
- **Impact of routine pathologic review in N0 BC:** 20% changes: grade 40%, LVI 26%, N 15%, margin 12% (JCO 2012)
- **pN0 from MIRROR study:** pN0 was upstaged in 22%, in central pathology review (Ann Oncol 2012)
- **Ki67:**
 - Inter- and intraobserver variability in measurement of Ki-67 is high (J Nat. Cancer Institute 2011)
 - High reproducibility for low and high Ki67 levels (J Pathol 2002)
 - Standardized methodology improves analytical validity (JNCI 2020)

1. Dowsett M, Nielsen TO, A'Hern R, et al: Assessment of Ki67 in breast cancer: recommendations from the International Ki67 in Breast Cancer working group. J Natl Cancer Inst 2011, 103(22):1656-1664.
2. Gluz O, Nitz UA, Christgen M, et al. West German Study Group Phase III PlanB Trial: First Prospective Outcome Data for the 21-Gene Recurrence Score Assay and Concordance of Prognostic Markers by Central and Local Pathology Assessment. J Clin Oncol. 2016 Jul 10;34(20):2341-9.
3. Hammond, M.E.H., Hayes, D.F., Dowsett, M., et al. 2010. American Society of Clinical Oncology/College Of American Pathologists guideline recommendations for immunohistochemical testing of estrogen and progesterone receptors in breast cancer. J. Clin. Oncol. 28, 2784–2795.
4. Sloane, J.P., Amendoeira, I., Apostolikas, N., et al. 1999. Consistency achieved by 23 European pathologists from 12 countries in diagnosing breast disease and reporting prognostic features of carcinomas. European Commission Working Group on Breast Screening Pathology. Virchows Arch. 434, 3–10.
5. Vestjens, J.H.M.J., Pepels, M.J., Boer, M. de, et al. 2012. Relevant impact of central pathology review on nodal classification in individual breast cancer patients. Ann. Oncol. 23, 2561–2566.
6. Kennecke, H.F., Speers, C.H., Ennis, C.A., et al. 2012. Impact of routine pathology review on treatment for node-negative breast cancer. J. Clin. Oncol. 30, 2227–2231.
7. Wolff AC, Hammond ME, Hicks DG, et al.: Recommendations for human epidermal growth factor receptor 2 testing in breast cancer:

American Society of Clinical Oncology/College of American Pathologists clinical practice guideline update. J Clin Oncol 2013, 31(31):3997-4013.

8. Mengel M, von Wasielewski R, Wiese B, et al. Inter-laboratory and inter-observer reproducibility of immunohistochemical assessment of the Ki67 labelling index in a large multi-centre trial. J Pathol. 2002 Nov;198(3):292-9.
9. Nielsen TO, Leung SCY, Rimm DL, et al. Assessment of Ki67 in Breast Cancer: Updated recommendations from the International Ki67 in Breast Cancer Working Group. J Natl Cancer Inst. 2020 Dec 28:djaa201.

Early Breast Cancer (M0) - eBC

Prognostic Factors III

Factor	Oxford		
	LoE	GR	AGO
■ Gene expression profiles (GEP, multigene assays, gene signatures)			
■ MammaPrint® (N0-1)	1b	A	+*
■ Oncotype DX® (N0-1, HR+ HER2-)	1b	A	+*
■ EndoPredict® (N0-1, HR+, HER2 -)	2b	B	+*
■ Prosigna® (N0-1, HR+, HER2 -)	2b	B	+*
■ Breast Cancer Index SM (N0-1, HR+ HER2-)**	2b	B	+/-*
■ IHC4 (ER / PR / HER2 / Ki-67) (validated for central testing)	2b	B	+/-
■ PREDICT® algorithm (https://breast.predict.nhs.uk/)	1b	A	+
■ Clinical-pathological score for lobular breast cancer (nodal status, tumor size, lymphovascular invasion LVI)	2b	B	+/-
■ CTS5 Clinical Treatment Score**	2b	B	+
■ CPS-EG Score	2b	B	+

* Should only be used in the context of clinical-pathological criteria (tumor size, nodal involvement, grade, Ki-67, ER, PR, HER2)
 ** Estimation of late recurrence

Gene expression profiles (GEP; Multigene Assays, Gene expression signatures)

(*Should only be used in the context of clinico-pathological criteria (e.g. tumor size, number involved lymph nodes, grade, Ki67) for therapeutic decision making)

1. Andre F, Ismaila N, Allison KH, et al. Biomarkers for Adjuvant Endocrine and Chemotherapy in Early-Stage Breast Cancer: ASCO Guideline Update. J Clin Oncol. 2022 Jun 1;40(16):1816-1837. doi: 10.1200/JCO.22.00069. Epub 2022 Apr 19. Erratum in: J Clin Oncol. 2022 Aug 1;40(22):2514. PMID: 35439025.

MammaPrint®

1. Cardoso F, van't Veer LJ et al.; MINDACT Investigators. 70-Gene Signature as an Aid to Treatment Decisions in Early-Stage Breast Cancer. N Engl J Med. 2016 Aug 25;375(8):717-29.
2. Cardoso F, van't Veer LJ, Bogaerts J, et al. 70-Gene Signature as an Aid to Treatment Decisions in Early-Stage Breast Cancer. N Engl J Med. 2016 Aug 25;375(8):717-29.
3. Mittempergher L, Delahaye LJM, Witteveen AT et al. MammaPrint and Blueprint Molecular Diagnostics Using Targeted RNA Next-Generation Sequencing Technology. J Mol Diagn. 2019 Sep;21(5):808-823
4. Piccart M, van 't Veer LJ, Poncet C, et al. 70-gene signature as an aid for treatment decisions in early breast cancer: updated results of

the phase 3 randomised MINDACT trial with an exploratory analysis by age. *Lancet Oncol.* 2021 Apr;22(4):476-488. doi: 10.1016/S1470-2045(21)00007-3. Epub 2021 Mar 12. PMID: 33721561.

5. Slembrouck L, Darrigues L, Laurent C et al. Decentralization of Next-Generation RNA Sequencing-Based MammaPrint® and Blueprint® Kit at University Hospitals Leuven and Curie Institute Paris. *Transl Oncol.* 2019 Dec;12(12):1557-1565.
6. Viale G, de Snoo FA et al.; MINDACT investigators. Immunohistochemical versus molecular (Blueprint and MammaPrint) subtyping of breast carcinoma. Outcome results from the EORTC 10041/BIG 3-04 MINDACT trial. *Breast Cancer Res Treat.* 2018 Jan;167(1):123-131

Oncotype DX®

1. Gluz O, Nitz UA, Christgen M, et al. West German Study Group Phase III PlanB Trial: First Prospective Outcome Data for the 21-Gene Recurrence Score Assay and Concordance of Prognostic Markers by Central and Local Pathology Assessment. *J Clin Oncol.* 2016 Jul 10;34(20):2341-9.
2. Kalinsky K, Barlow WE, Gralow JR et al. 21-Gene Assay to inform chemotherapy benefit in node-positive breast cancer. *N Engl J Med.* 2021 Dec 16;385(25):2336-2347. doi: 10.1056/NEJMoa2108873. Epub 2021 Dec 1.
3. Nitz U, Gluz O, Christgen M et al. Reducing chemotherapy use in clinically high-risk, genomically low-risk pN0 and pN1 early breast cancer patients: five-year data from the prospective, randomised phase 3 West German Study Group (WSG) PlanB trial. *Breast Cancer Res Treat.* 2017 Oct;165(3):573-583.
4. Nitz UA, Gluz O, Kümmel S, et al. Endocrine Therapy Response and 21-Gene Expression Assay for Therapy Guidance in HR+/HER2- Early Breast Cancer. *J Clin Oncol.* 2022 Aug 10;40(23):2557-2567. doi: 10.1200/JCO.21.02759. Epub 2022 Apr 11. PMID: 35404683.
5. Sparano JA, Gray RJ, Makower DF et al. Clinical Outcomes in Early Breast Cancer With a High 21-Gene Recurrence Score of 26 to 100 Assigned to Adjuvant Chemotherapy Plus Endocrine Therapy: A Secondary Analysis of the TAILORx Randomized Clinical Trial. *JAMA Oncol.* 2019 Sep 30.
6. Sparano JA, Gray RJ, Makower DF et al. Adjuvant Chemotherapy Guided by a 21-Gene Expression Assay in Breast Cancer. *N Engl J Med.* 2018 Jul 12;379(2):111-121.

EndoPredict®

1. Buus R, Sestak I, Kronenwett R et al. Comparison of EndoPredict and EPclin With Oncotype DX Recurrence Score for Prediction of Risk

- of Distant Recurrence After Endocrine Therapy. *J Natl Cancer Inst.* 2016 Jul 10;108(11).
2. Buus R, I. Sestak, R. Kronenwett, et al (2016). "Comparison of EndoPredict and EPclin With Oncotype DX Recurrence Score for Prediction of Risk of Distant Recurrence After Endocrine Therapy." *J Natl Cancer Inst* 108(11)
 3. Filipits M, Dubsky P, Rudas M et al. Prediction of Distant Recurrence Using EndoPredict Among Women with ER+, HER2- Node-Positive and Node-Negative Breast Cancer Treated with Endocrine Therapy Only. *Clin Cancer Res.* 2019 Jul 1;25(13):3865-3872.
 4. Martin M, Brase JC, Ruiz A et al. Prognostic ability of EndoPredict compared to research-based versions of the PAM50 risk of recurrence (ROR) scores in node-positive, estrogen receptor-positive, and HER2-negative breast cancer. A GEICAM/9906 sub-study. *Breast Cancer Res Treat.* 2016 Feb;156(1):81-9.
 5. Sestak I, Buus R, Cuzick J et al. Comparison of the Performance of 6 Prognostic Signatures for Estrogen Receptor-Positive Breast Cancer: A Secondary Analysis of a Randomized Clinical Trial. *JAMA Oncol.* 2018 Apr 1;4(4):545-553.
 6. Sestak I, Martín M, Dubsky P et al. Prediction of chemotherapy benefit by EndoPredict in patients with breast cancer who received adjuvant endocrine therapy plus chemotherapy or endocrine therapy alone. *Breast Cancer Res Treat.* 2019 Jul;176(2):377-386.

Prosigna®

1. Berchtold E, Vetter M, Gündert M et al. Comparison of six breast cancer classifiers using qPCR. *Bioinformatics.* 2019 Sep 15;35(18):3412-3420.
2. Buus R, Szijgyarto Z, Schuster EF, et al. Development and validation for research assessment of Oncotype DX® Breast Recurrence Score, EndoPredict® and Prosigna®. *NPJ Breast Cancer.* 2021 Feb 12;7(1):15. doi: 10.1038/s41523-021-00216-w.
3. Erber R, Angeloni M, Stöhr R, et al. Molecular Subtyping of Invasive Breast Cancer Using a PAM50-Based Multigene Expression Test-Comparison with Molecular-Like Subtyping by Tumor Grade/Immunohistochemistry and Influence on Oncologist's Decision on Systemic Therapy in a Real-World Setting. *Int J Mol Sci.* 2022 Aug 5;23(15):8716. doi: 10.3390/ijms23158716.
4. Fernandez-Martinez A, Pascual T, Perrone G et al. Limitations in predicting PAM50 intrinsic subtype and risk of relapse score with Ki67 in estrogen receptor-positive HER2-negative breast cancer. *Oncotarget.* 2017 Mar 28;8(13):21930-21937.
5. Jensen MB, Lænkholm AV, Nielsen TO et al. The Prosigna gene expression assay and responsiveness to adjuvant cyclophosphamide-based chemotherapy in premenopausal high-risk patients with breast cancer. *Breast Cancer Res.* 2018 Jul 27;20(1):79.
6. Lænkholm AV, Jensen MB, Eriksen JO et al. PAM50 Risk of Recurrence Score Predicts 10-Year Distant Recurrence in a Comprehensive Danish Cohort of Postmenopausal Women Allocated to 5 Years of Endocrine Therapy for Hormone Receptor-Positive Early Breast Cancer. *J Clin Oncol.* 2018 Mar 10;36(8):735-740.
7. Laenkholm AV, Jensen MB, Eriksen JO et al. The ability of PAM50 risk of recurrence score to predict 10-year distant recurrence in

- hormone receptor-positive postmenopausal women with special histological subtypes. *Acta Oncol.* 2018 Jan;57(1):44-50.
8. Ohnstad HO, Borgen E, Falk RS et al. Prognostic value of PAM50 and risk of recurrence score in patients with early-stage breast cancer with long-term follow-up. *Breast Cancer Res.* 2017 Nov 14;19(1):120.
 9. Ohara AM, Naoi Y, Shimazu K et al. PAM50 for prediction of response to neoadjuvant chemotherapy for ER-positive breast cancer. *Breast Cancer Res Treat.* 2019 Feb;173(3):533-543
 - Prat A, Lluch A, Turnbull AK et al. A PAM50-Based Chemoendocrine Score for Hormone Receptor-Positive Breast Cancer with an Intermediate Risk of Relapse. *Clin Cancer Res.* 2017 Jun 15;23(12):3035-3044.

Breast Cancer Index®

1. Bartlett JMS, Sgroi DC, Treuner K et al. Breast Cancer Index and prediction of benefit from extended endocrine therapy in breast cancer patients treated in the Adjuvant Tamoxifen-To Offer More? (aTTom) trial. *Ann Oncol.* 2019 Nov 1;30(11):1776-1783.
2. Noordhoek I, Treuner K, Putter H, et al. Breast Cancer Index Predicts Extended Endocrine Benefit to Individualize Selection of Patients with HR⁺ Early-stage Breast Cancer for 10 Years of Endocrine Therapy. *Clin Cancer Res.* 2021 Jan 1;27(1):311-319. doi: 10.1158/1078-0432.CCR-20-2737. Epub 2020 Oct 27. PMID: 33109739.
3. Sgroi DC, Treuner K, Zhang Y, et al. Correlative studies of the Breast Cancer Index (HOXB13/IL17BR) and ER, PR, AR, AR/ER ratio and Ki67 for prediction of extended endocrine therapy benefit: a Trans-aTTom study. *Breast Cancer Res.* 2022 Dec 16;24(1):90. doi: 10.1186/s13058-022-01589-x.

IHC-4 Score

1. Cuzick J, Dowsett M, Pineda S et al. Prognostic value of a combined estrogen receptor, progesterone receptor, Ki-67, and human epidermal growth factor receptor-2 immunohistochemical score and comparison with the Genomic Health recurrence score in early breast cancer. *J Clin Oncol* 2011;29(32):4273-4278.
2. Sheri A, Smith IE, Hills M et al. Relationship between IHC4 Score and response to neo-adjuvant chemotherapy in estrogen receptor positive breast cancer. *Breast Cancer Res Treat* 2017;164(2):395-400

PREDICT (<https://breast.predict.nhs.uk/>)

1. Candido Dos Reis FJ, Wishart GC, Dicks EM, et al. An updated PREDICT breast cancer prognostication and treatment benefit prediction model with independent validation. *Breast Cancer Res.* ; 2017;19(1):58.
2. Gunda A, Eshwaraiah MS, Gangappa K, et al. A comparative analysis of recurrence risk predictions in ER+/HER2- early breast cancer using NHS Nottingham Prognostic Index, PREDICT, and CanAssist Breast. *Breast Cancer Res Treat.* 2022 Nov;196(2):299-310. doi:

10.1007/s10549-022-06729-7. Epub 2022 Sep 10.

Lobular Score:

1. De Nonneville A, Jauffret C, Goncalves A. et al. Adjuvant chemotherapy in lobular carcinoma of the breast: a clinicopathological score identifies high-risk patient with survival benefit *Breast Cancer Res Treat.* 2019 Jun;175(2):379-387. : Points-based risk score: pN1 = 6 points, tumnor size > 2cm = 3 points, LVI = 2 points; low risk = < 5 points; high risk 5-11 points.
2. Fu R, Yang J, Wang H et al.: A nomogram for determining the disease-specific survival in invasive lobular carcinoma of the breast: A population study. *Medicine (Baltimore).* 2020 Oct 23;99(43):e22807.

CTS Clinical Treatment Score

1. Lakhanpal R, Sestak I, Shadbolt B, et al. IHC4 score plus clinical treatment score predicts locoregional recurrence in early breast cancer. *Breast.* 2016;29:147–152.
2. Richman J, Ring A, Dowsett M, Sestak I. Clinical validity of clinical treatment score 5 (CTS5) for estimating risk of late recurrence in unselected, non-trial patients with early oestrogen receptor-positive breast cancer. *Breast Cancer Res Treat.* 2020 Nov 21.
3. Wang C, Xu Y, Lin Y, et al. Comparison of CTS5 risk model and 21-gene recurrence score assay in large-scale breast cancer population and combination of CTS5 and recurrence score to develop a novel nomogram for prognosis prediction. *Breast.* 2022 Jun;63:61-70. doi: 10.1016/j.breast.2022.03.007. Epub 2022 Mar 16.

CPS-EG Score

1. Loibl S, Weber K, Huober J et al.: Risk Assessment after Neoadjuvant Chemotherapy in Luminal Breast Cancer Using a Clinicomolecular Predictor. *Cancer Res.* 2018 Jul 15;24(14):3358-3365.
2. Marmé F, Solbach C, Michel L, et al. Utility of the CPS + EG scoring system in triple-negative breast cancer treated with neoadjuvant chemotherapy. *Eur J Cancer.* 2021 Aug;153:203-212. doi: 10.1016/j.ejca.2021.05.027. Epub 2021 Jun 26. PMID: 34186505.
3. Mittendorf EA, Jeruss JS, Tucker SL et al. Validation of a novel staging system for disease-specific survival in patients with breast cancer treated with neoadjuvant chemotherapy. *J Clin Oncol.* 2011 May 20;29(15):1956-62.

Early Breast Cancer (M0) - eBC Prognostic Factors IV

Factor	Oxford		
	LoE	GR	AGO
■ Disseminated tumor cells (DTC, in bone marrow)	1a	A	+/-
■ Circulating tumor cells (CTC, in blood, Cell Search®)*	1b	A	+/-
■ CTC before NACT (regarding OS, DDFS, LRFI)	1b	B	+/-
■ Therapy decisions based on CTC phenotypes	3a	C	-
■ Cell-free DNA (cfDNA, in blood, for DFS, PFS, OS)	2a	B	+/-

* Validated clinical data only available for this assay

DTC

1. Hartkopf AD, Brucker SY, Taran FA, et al. Disseminated tumour cells from the bone marrow of early breast cancer patients: Results from an international pooled analysis. Eur J Cancer. 2021 Sep;154:128-137.
2. Janni, W., Vogl, F.D., Wiedswang, G. et al. Persistence of disseminated tumor cells in the bone marrow of breast cancer patients predicts increased risk for relapse--a European pooled analysis. Clin. Cancer Res. 2011; 17, 2967–2976.

DTC and radiation

1. Mignot F, Loirat D, Dureau S, et al. Disseminated Tumor Cells Predict Efficacy of Regional Nodal Irradiation in Early Stage Breast Cancer. Int J Radiat Oncol Biol Phys. 2019 Feb 1;103(2):389-396.
2. Goodman CR, Seagle BL, Friedl TWP, et al. Association of Circulating Tumor Cell Status With Benefit of Radiotherapy and Survival in Early-Stage Breast Cancer. JAMA Oncol. 2018 Aug 1;4(8):e180163.

CTC

1. Bidard FC, Michiels S, Riethdorf S, et al. Circulating Tumor Cells in Breast Cancer Patients Treated by Neoadjuvant Chemotherapy: A Meta-analysis. J Natl Cancer Inst. 2018 Jun 1;110(6):560-567.
2. Cristofanilli, M., Budd, G.T., Ellis, M.J., et al. 2004. Circulating tumor cells, disease progression, and survival in metastatic breast

cancer. N. Engl. J. Med. 351, 781–791. doi:10.1056/NEJMoa040766.

3. Garcia-Murillas I, Chopra N Comino-Méndez I et al. Assessment of molecular relapse detection in early stage breast cancer. JAMA Oncol 2019;5(10):1473-1478)
4. Trapp E, Janni W, Schindlbeck C, et al; SUCCESS Study Group. Presence of Circulating Tumor Cells in High-Risk Early Breast Cancer During Follow-Up and Prognosis. J Natl Cancer Inst. 2018 Oct 11.
5. Radovich M, Jiang G, Hancock BA, et al. Association of Circulating Tumor DNA and Circulating Tumor Cells After Neoadjuvant Chemotherapy With Disease Recurrence in Patients With Triple-Negative Breast Cancer: Preplanned Secondary Analysis of the BRE12-158 Randomized Clinical Trial. JAMA Oncol. 2020 Sep 1;6(9):1410-1415. doi: 10.1001/jamaoncol.2020.2295.

Therapy decision based on CTCs

1. Turner NC, Kingston B, Kilburn LS et al. Circulating tumor DNA analysis to direct therapy in advanced breast cancer(plasmaMATCH):a multicentre, multicohort, phase 2a, platform trial. Lancet Oncol 2020;21(10):1296-1308.

Cell-free DNA

1. Cullinane C, Fleming C, O'Leary DP, et al. Association of Circulating Tumor DNA With Disease-Free Survival in Breast Cancer: A Systematic Review and Meta-analysis. JAMA Netw Open. 2020 Nov 2;3(11):e2026921.

Commercially Available Molecular Tests					
	70 gene signature (MammaPrint®) §	21 gene Recurrence score (Oncotype DX®) §	8 gene signature (Endopredict®) §	PAM 50 (Prosigna®) §	Breast Cancer Index® (BCI) §
Provider	Agendia	Genomic Health	Sividon (Myriad)	NanoString	Biotheranostics
Type of assay	70-gene assay	21-gene recurrence score	11-gene assay	50-gene assay	5 + 2 (MGI+H/I)
Type of tissue	fresh frozen (technical validation for FFPE available)	FFPE	FFPE	FFPE	FFPE
Technique	Microarrays for RNA	qRT-PCR	q-RT-PCR	Direct hybridization (nCounter®)	q-RT-PCR
Central lab	yes	yes	no	no	yes
Indication and population studied	prognostic N-/+, < 70 Jahre	prognostic N-/+, ER+ endocrine treated	prognostic (pre-) postmenopausal N-/+, ER+ HER2- endocrine treated	prognostic postmenopausal N-/+, ER+ HER2- endocrine treated	Prognostic pT1-3pN0 – pN1 ER+ / HER2- Endocrine treated
Risk classes	Low – high	RS (Low – intermediate – high)	Low – high	ROR (Low – inter- mediate – high) molecular types	Low - high
Clinical Validation	Yes	yes	yes	yes	Yes
Registration	FDA clearance as "In Vitro Diagnostic Multivariate Index Assay (IVDMIA)®" CE-Mark (fresh tissue and FFPE)	Clinical Laboratory Improvement Amendments (CLIA) + College of American Pathologists (CAP) -accredited ref lab	CE-Mark	CE-Mark FDA 510(k) Clearance	Service Mark (SM)

§ Validated clinical data only available for this assay

Head to head comparisons

1. Varga Z, Sinn P, Seidman AD. Summary of head-to-head comparisons of patient risk classifications by the 21-gene Recurrence Score® (RS) assay and other genomic assays for early breast cancer. Int J Cancer. 2019 Aug 15;145(4):882-893.
2. Berchtold E, Vetter M, Gündert M et al. Comparison of six breast cancer classifiers using qPCR. Bioinformatics. 2019 Sep 15;35(18):3412-3420.
3. Sestak I, Buus R, Cuzick J, et al. Comparison of the Performance of 6 Prognostic Signatures for Estrogen Receptor-Positive Breast Cancer: A Secondary Analysis of a Randomized Clinical Trial. JAMA Oncol. 2018 Apr 1;4(4):545-553.

Endopredict

1. Blank PR, Filipits M, Dubsky P, et al.: Cost-effectiveness analysis of prognostic gene expression signature-based stratification of early breast cancer patients. Pharmacoeconomics. 2015 Feb;33(2):179-90.
2. Buus R, Sestak I, Kronenwett R, et al.: Comparison of EndoPredict and EPclin with Oncotype DX Recurrence Score for Prediction of Risk of Distant Recurrence after endocrine therapy J Natl Cancer Inst. 2016 Jul 10;108(11).

3. Denkert, C., Kronenwett, R., Schlake, W. et al. 2012. Decentral gene expression analysis for ER+/Her2- breast cancer: results of a proficiency testing program for the EndoPredict assay. *Virchows Arch.* 460, 251–259. doi:10.1007/s00428-012-1204-4.
4. Dubsky, P., Filipits, M., Jakesz, R. et al. 2012. EndoPredict improves the prognostic classification derived from common clinical guidelines in ER-positive, HER2-negative early breast cancer. *Ann Oncol.* 2013 Mar;24(3):640-7.
5. Dubsky P, Brase JC, Jakesz R et al.: Austrian Breast and Colorectal Cancer Study Group (ABCSG). T al.: The EndoPredict score provides prognostic information on late distant metastases in ER+/HER2- breast cancer patients. *Br J Cancer.* 2013 Dec 10;109(12):2959-64
6. Filipits, M., Rudas, M., Jakesz, R. et al. 2011. A new molecular predictor of distant recurrence in ER-positive, HER2-negative breast cancer adds independent information to conventional clinical risk factors. *Clin. Cancer Res.* 17, 6012–6020.
7. Kronenwett, R., Bohmann, K., Prinzler, J. et al. 2012. Decentral gene expression analysis: analytical validation of the Endopredict genomic multianalyte breast cancer prognosis test. *BMC Cancer* 12, 456.
8. Martin M, Brase JC, Calvo L, et al. Clinical validation of the EndoPredict test in node-positive, chemotherapy-treated ER+/HER2- breast cancer patients: results from the GEICAM 9906 trial. *Breast Cancer Res* 2014; 16(2): R38.

MammaPrint

1. Buyse, M., Loi, S., van't Veer, L., et al. 2006. Validation and clinical utility of a 70-gene prognostic signature for women with node-negative breast cancer. *J. Natl. Cancer Inst.* 98, 1183–1192.
2. Drukker CA, Elias SG, Nijenhuis MV, et al. Gene expression profiling to predict the risk of locoregional recurrence in breast cancer: a pooled analysis. *Breast Cancer Res Treat.* 2014 Dec;148(3):599-613.
3. Exner R, Bago-Horvath Z, Bartsch R et al. The multigene signature MammaPrint impacts on multidisciplinary team decisions in ER+, HER2- early breast cancer. *Br J Cancer.* 2014 Aug 26;111(5):837-42.
4. Jonsdottir K, Assmus J, Slewa A, et al. Prognostic value of gene signatures and proliferation in lymph-node-negative breast cancer. *PLoS One.* 2014 Mar 5;9(3):e90642.
5. Mook, S., Schmidt, M.K., Weigelt, B., et al. 2010. The 70-gene prognosis signature predicts early metastasis in breast cancer patients

between 55 and 70 years of age. *Ann. Oncol.* 21, 717–722.

6. Mook, S., Schmidt, M.K., Rutgers, E.J., et al. 2009a. Calibration and discriminatory accuracy of prognosis calculation for breast cancer with the online Adjuvant! program: a hospital-based retrospective cohort study. *Lancet Oncol.* 10, 1070–1076.
7. Mook, S., Schmidt, M.K., Viale, G. et al. 2009b. The 70-gene prognosis-signature predicts disease outcome in breast cancer patients with 1-3 positive lymph nodes in an independent validation study. *Breast Cancer Res. Treat.* 116, 295–302.
8. van de Vijver, M.J., He, Y.D., van't Veer, L.J., et al. 2002. A gene-expression signature as a predictor of survival in breast cancer. *N. Engl. J. Med.* 347, 1999–2009. doi:10.1056/NEJMoa021967.
9. van Veer, L.J. 't, Dai, H., van de Vijver, M.J., et al. 2002. Gene expression profiling predicts clinical outcome of breast cancer. *Nature* 415, 530–536. doi:10.1038/415530a.
10. Cardoso F, van't Veer LJ, Bogaerts J, et al. 70-Gene Signature as an Aid to Treatment Decisions in Early-Stage Breast Cancer. *N Engl J Med.* 2016 Aug 25;375(8):717-29.
11. Piccart M, van 't Veer LJ, Poncet C, et al. 70-gene signature as an aid for treatment decisions in early breast cancer: updated results of the phase 3 randomised MINDACT trial with an exploratory analysis by age. *Lancet Oncol.* 2021 Apr;22(4):476-488. doi: 10.1016/S1470-2045(21)00007-3. Epub 2021 Mar 12. PMID: 33721561.

Oncotype DX

1. Albain, K.S., Barlow, W.E., Shak, S. et al. 2010. Prognostic and predictive value of the 21-gene recurrence score assay in postmenopausal women with node-positive, oestrogen-receptor-positive breast cancer on chemotherapy: a retrospective analysis of a randomised trial. *Lancet Oncol.* 11, 55–65. Cronin, M., Sangli, C., Liu, M.-L., et al. 2007. Analytical validation of the Oncotype DX genomic diagnostic test for recurrence prognosis and therapeutic response prediction in node-negative, estrogen receptor-positive breast cancer. *Clin. Chem.* 53, 1084–1091.
2. Dowsett, M., Cuzick, J., Wale, C. et al. 2010. Prediction of risk of distant recurrence using the 21-gene recurrence score in node-negative and node-positive postmenopausal patients with breast cancer treated with anastrozole or tamoxifen: a TransATAC study. *J.*

Clin. Oncol. 28, 1829–1834.

3. Geyer CE Jr, Tang G, Mamounas EP, et al. 21-Gene assay as predictor of chemotherapy benefit in HER2-negative breast cancer. *NPI Breast Cancer*. 2018 Nov 14;4:37.
4. Gluz O, Nitz UA, Christgen M, et al.: West German Study Group Phase III PlanB Trial: First Prospective Outcome Data for the 21-Gene Recurrence Score Assay and Concordance of Prognostic Markers by Central and Local Pathology Assessment. *J Clin Oncol*. 2016 Jul 10;34(20):2341-9.
5. Kalinsky K, Barlow WE, Gralow JR et al. 21-Gene Assay to inform chemotherapy benefit from node positive breast cancer. *Engl J Med* DOI: 10.1056/NEJMoa2108873
6. Khan SS, Karn T, Symmans WF, et al. Genomic predictor of residual risk of recurrence after adjuvant chemotherapy and endocrine therapy in high risk estrogen receptor-positive breast cancers. *Breast Cancer Res Treat*. 2015 Feb 5.
7. Mamounas, E.P., Tang, G., Fisher, B., et al. 2010. Association between the 21-gene recurrence score assay and risk of locoregional recurrence in node-negative, estrogen receptor-positive breast cancer: results from NSABP B-14 and NSABP B-20. *J. Clin. Oncol*. 28, 1677–1683.
8. Nitz U, Gluz O, Christgen M, et al. Reducing chemotherapy use in clinically high-risk, genomically low-risk pN0 and pN1 early breast cancer patients: five-year data from the prospective, randomised phase 3 West German Study Group (WSG) PlanB trial. *Breast Cancer Res Treat*. 2017 Oct;165(3):573-583.
9. Nitz UA, Gluz O, Kümmel S et al. Endocrine Therapy Response and 21-Gene Expression Assay for Therapy Guidance in HR+/HER2- Early Breast Cancer. *J Clin Oncol*. 2022 Aug 10;40(23):2557-2567. doi: 10.1200/JCO.21.02759. Epub 2022 Apr 11.
10. Paik, S., Shak, S., Tang, G. et al. 2004. A multigene assay to predict recurrence of tamoxifen-treated, node-negative breast cancer. *N. Engl. J. Med*. 351, 2817–2826. doi:10.1056/NEJMoa041588.
11. Paik, S., Tang, G., Shak, S., et al. 2006. Gene expression and benefit of chemotherapy in women with node-negative, estrogen receptor-positive breast cancer. *J. Clin. Oncol*. 24, 3726–3734.
12. Penault-Llorca F, Filleron T, Asselain B, et al. The 21-gene Recurrence Score® assay predicts distant recurrence in lymph node-positive, hormone receptor-positive, breast cancer patients treated with adjuvant sequential epirubicin- and docetaxel-based or epirubicin-

based chemotherapy (PACS-01 trial). BMC Cancer. 2018 May 4;18(1):526.

13. Sparano JA, Gray RJ, Makower DF, et al: Prospective Validation of a 21-Gene Expression Assay in Breast Cancer. N Engl J Med 2015, 373(21):2005-2014.
14. Sparano JA, Gray R, Oktay MH, et al. A metastasis biomarker (MetaSite Breast™ Score) is associated with distant recurrence in hormone receptor-positive, HER2-negative early-stage breast cancer. NPJ Breast Cancer. 2017 Nov 8;3:42.
15. Sparano JA, Gray RJ, Makower DF, et al. Adjuvant Chemotherapy Guided by a 21-Gene Expression Assay in Breast Cancer. N Engl J Med. 2018 Jul 12;379(2):111-121.
16. Tang, G., Cuzick, J., Costantino, J.P., et al. 2011. Risk of recurrence and chemotherapy benefit for patients with node-negative, estrogen receptor-positive breast cancer: recurrence score alone and integrated with pathologic and clinical factors. J. Clin. Oncol. 29, 4365–4372.
17. Zhao X, Rødland EA, Sørli T, et al. Systematic assessment of prognostic gene signatures for breast cancer shows distinct influence of time and ER status. BMC Cancer. 2014 Mar 19;14:211.
18. Sparano J, Gray RJ, Makower D, Albain KS, Hayes DF, Geyer C, Dees E, Goetz MP, Olson Jr JA, Lively TG, Badve S, Saphner T, Wagner LI, Whelan T, Kaklamani V, Sledge Jr GW. Trial Assigning Individualized Options for Treatment (TAILORx): An update including 12-year event rates. SABCS 2022, GS 1-05

Prosigna (ROR / PAM50)

1. Chia, S.K., Bramwell, V.H., Tu, Det al. 2012. A 50-gene intrinsic subtype classifier for prognosis and prediction of benefit from adjuvant tamoxifen. Clin. Cancer Res. 18, 4465–4472.
2. Gnant M, Filipits M, Greil R, et al. Austrian Breast and Colorectal Cancer Study Group. Predicting distant recurrence in receptor-positive breast cancer patients with limited clinicopathological risk: using the PAM50 Risk of Recurrence score in 1478 postmenopausal patients of the ABCSG-8 trial treated with adjuvant endocrine therapy alone. Ann Oncol. 2014 Feb;25(2):339-45
3. Jensen MB, Lænkholm AV, Nielsen TO, et al. The Prosigna gene expression assay and responsiveness to adjuvant cyclophosphamide-

based chemotherapy in premenopausal high-risk patients with breast cancer. *Breast Cancer Res.* 2018 Jul 27;20(1):79.

4. Lænkholm AV, Jensen MB, Eriksen JO, et al. PAM50 Risk of Recurrence Score Predicts 10-Year Distant Recurrence in a Comprehensive Danish Cohort of Postmenopausal Women Allocated to 5 Years of Endocrine Therapy for Hormone Receptor-Positive Early Breast Cancer. *J Clin Oncol.* 2018 Mar 10;36(8):735-740.
5. Liu S, Chapman JA, Burnell MJ, et al. Prognostic and predictive investigation of PAM50 intrinsic subtypes in the NCIC CTG MA.21 phase III chemotherapy trial. *Breast Cancer Res Treat.* 2015 Jan;149(2):439-48
Nielsen, T.O., Parker, J.S., Leung, S., et al. 2010. A comparison of PAM50 intrinsic subtyping with immunohistochemistry and clinical prognostic factors in tamoxifen-treated estrogen receptor-positive breast cancer. *Clin. Cancer Res.* 16, 5222–5232.
6. Ohnstad HO, Borgen E, Falk RS, et al. Prognostic value of PAM50 and risk of recurrence score in patients with early-stage breast cancer with long-term follow-up. *Breast Cancer Res.* 2017 Nov 14;19(1):120.
7. Parker, J.S., Mullins, M., Cheang, M.C.U., et al. 2009. Supervised risk predictor of breast cancer based on intrinsic subtypes. *J. Clin. Oncol.* 27, 1160–1167.
8. Prat, A., Cheang, M.C.U., Martín, M., et al. 2012b. Prognostic Significance of Progesterone Receptor-Positive Tumor Cells Within Immunohistochemically Defined Luminal A Breast Cancer. *J. Clin. Oncol.*
9. Prat, A., Parker, J.S., Fan, C., et al. 2012a. Concordance among gene expression-based predictors for ER-positive breast cancer treated with adjuvant tamoxifen. *Ann. Oncol.* 23, 2866–2873.
10. Perou, C.M., Sørlie, T., Eisen, M.B. et al. 2000. Molecular portraits of human breast tumours. *Nature* 406, 747–752.
11. Pogue-Geile KL, Song N, Jeong JH, et al. Intrinsic Subtypes, PIK3CA Mutation, and the Degree of Benefit From Adjuvant Trastuzumab in the NSABP B-31 Trial. *J Clin Oncol.* 2015 Jan 5.
12. Sestak I, Cuzick J, Dowsett M. et al.. Prediction of Late Distant Recurrence After 5 Years of Endocrine Treatment: A Combined Analysis of Patients From the Austrian Breast and Colorectal Cancer Study Group 8 and Arimidex, Tamoxifen Alone or in Combination Randomized Trials Using the PAM50 Risk of Recurrence Score. *J Clin Oncol.* 2014 Oct 20. pii: JCO.2014.55.6894

Breast Cancer Index

1. Bartlett JMS, Sgroi DC, Treuner K et al. Breast Cancer Index and prediction of benefit from extended endocrine therapy in breast cancer patients treated in the Adjuvant Tamoxifen-To Offer More? (aTTom) trial. *Ann Oncol.* 2019 Nov 1;30(11):1776-1783.
2. Jerevall PL, Brommesson S, Strand C, et al. Exploring the two-gene ratio in breast cancer--independent roles for HOXB13 and IL17BR in prediction of clinical outcome. *Breast Cancer Res Treat.* 2008;107(2):225–234.
3. Jansen MP, Sieuwerts AM, Look MP, et al. HOXB13-to-IL17BR expression ratio is related with tumor aggressiveness and response to tamoxifen of recurrent breast cancer: a retrospective study. *J Clin Oncol.* 2007;25(6):662–668.
4. Ma XJ, Hilsenbeck SG, Wang W, et al. The HOXB13:IL17BR expression index is a prognostic factor in early-stage breast cancer. *J Clin Oncol.* 2006;24(28):4611–4619.

Commercially Available Molecular Tests					
	70 gene signature (MammaPrint®) §	21 gene Recurrence score (Oncotype DX®) §	8 gene signature (Endopredict®) §	PAM 50 (Prosigna®) §	Breast Cancer Index® (BCI)
Prognosis after 5 yrs (late recurrences)	not separately shown	yes	yes	yes	yes
Predictive impact (chemotherapy benefit)	poorly validated	yes	not shown	not shown	EAT after 5 yrs
Prospective- retrospective evidence (% of recruited patients)	Multicenter validation	NSABP B-14 (14%) NSABP B-20 (28%) ECOG 9127 SWOG 8814 (40%) ATAC (30%)	ABCSG 6 (19%) ABCSG 8 (36%) GEICAM-9906 (45%) ATAC (10%)	MA.12 (59%) MA.5 (66%) ABCSG 8 (44%) ATAC (16%)	TransATTOM (11%)
Prospective evidence	MINDACT (N0, N1) (8y DFS, OS)	TAILORx (12 y DFS, OS), N0, RS ≤ 25 vs. ≥ 26 PlanB (N0 highrisk/N+) (5 y DFS, OS) RxPONDER (5 y DFS, OS), N1, RS ≤ 25 vs. ≥ 26 ADAPT (5 y DFS, OS), N0-1, RS 0-11; RS 12- 25 / Ki67 response	–	–	–

§ Validated clinical data only available for this assay



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.
Guidelines Breast
Version 2023.1E

www.ago-online.de
FORSCHEN
LEBEN
HEILEN

Head to head comparisons

1. Varga Z, Sinn P, Seidman AD. Summary of head-to-head comparisons of patient risk classifications by the 21-gene Recurrence Score® (RS) assay and other genomic assays for early breast cancer. Int J Cancer. 2019 Aug 15;145(4):882-893.
2. Berchtold E, Vetter M, Gündert M et al. Comparison of six breast cancer classifiers using qPCR. Bioinformatics. 2019 Sep 15;35(18):3412-3420.
3. Sestak I, Buus R, Cuzick J, et al. Comparison of the Performance of 6 Prognostic Signatures for Estrogen Receptor-Positive Breast Cancer: A Secondary Analysis of a Randomized Clinical Trial. JAMA Oncol. 2018 Apr 1;4(4):545-553.

Endopredict

1. Blank PR, Filipits M, Dubsky P, et al.: Cost-effectiveness analysis of prognostic gene expression signature-based stratification of early breast cancer patients. Pharmacoeconomics. 2015 Feb;33(2):179-90.
2. Buus R, Sestak I, Kronenwett R, et al.: Comparison of EndoPredict and EPclin with Oncotype DX Recurrence Score for Prediction of Risk of Distant Recurrence after endocrine therapy J Natl Cancer Inst. 2016 Jul 10;108(11).

3. Denkert, C., Kronenwett, R., Schlake, W. et al. 2012. Decentral gene expression analysis for ER+/Her2- breast cancer: results of a proficiency testing program for the EndoPredict assay. *Virchows Arch.* 460, 251–259. doi:10.1007/s00428-012-1204-4.
4. Dubsky, P., Filipits, M., Jakesz, R. et al. 2012. EndoPredict improves the prognostic classification derived from common clinical guidelines in ER-positive, HER2-negative early breast cancer. *Ann Oncol.* 2013 Mar;24(3):640-7.
5. Dubsky P, Brase JC, Jakesz R et al.: Austrian Breast and Colorectal Cancer Study Group (ABCSG). T al.: The EndoPredict score provides prognostic information on late distant metastases in ER+/HER2- breast cancer patients. *Br J Cancer.* 2013 Dec 10;109(12):2959-64
6. Filipits, M., Rudas, M., Jakesz, R. et al. 2011. A new molecular predictor of distant recurrence in ER-positive, HER2-negative breast cancer adds independent information to conventional clinical risk factors. *Clin. Cancer Res.* 17, 6012–6020.
7. Kronenwett, R., Bohmann, K., Prinzler, J. et al. 2012. Decentral gene expression analysis: analytical validation of the Endopredict genomic multianalyte breast cancer prognosis test. *BMC Cancer* 12, 456.
8. Martin M, Brase JC, Calvo L, et al. Clinical validation of the EndoPredict test in node-positive, chemotherapy-treated ER+/HER2- breast cancer patients: results from the GEICAM 9906 trial. *Breast Cancer Res* 2014; 16(2): R38.

MammaPrint

1. Buyse, M., Loi, S., van't Veer, L., et al. 2006. Validation and clinical utility of a 70-gene prognostic signature for women with node-negative breast cancer. *J. Natl. Cancer Inst.* 98, 1183–1192.
2. Drukker CA, Elias SG, Nijenhuis MV, et al. Gene expression profiling to predict the risk of locoregional recurrence in breast cancer: a pooled analysis. *Breast Cancer Res Treat.* 2014 Dec;148(3):599-613.
3. Exner R, Bago-Horvath Z, Bartsch R et al. The multigene signature MammaPrint impacts on multidisciplinary team decisions in ER+, HER2- early breast cancer. *Br J Cancer.* 2014 Aug 26;111(5):837-42.
4. Jonsdottir K, Assmus J, Slewa A, et al. Prognostic value of gene signatures and proliferation in lymph-node-negative breast cancer. *PLoS One.* 2014 Mar 5;9(3):e90642.
5. Mook, S., Schmidt, M.K., Weigelt, B., et al. 2010. The 70-gene prognosis signature predicts early metastasis in breast cancer patients

between 55 and 70 years of age. *Ann. Oncol.* 21, 717–722.

6. Mook, S., Schmidt, M.K., Rutgers, E.J., et al. 2009a. Calibration and discriminatory accuracy of prognosis calculation for breast cancer with the online Adjuvant! program: a hospital-based retrospective cohort study. *Lancet Oncol.* 10, 1070–1076.
7. Mook, S., Schmidt, M.K., Viale, G. et al. 2009b. The 70-gene prognosis-signature predicts disease outcome in breast cancer patients with 1-3 positive lymph nodes in an independent validation study. *Breast Cancer Res. Treat.* 116, 295–302.
8. van de Vijver, M.J., He, Y.D., van't Veer, L.J., et al. 2002. A gene-expression signature as a predictor of survival in breast cancer. *N. Engl. J. Med.* 347, 1999–2009. doi:10.1056/NEJMoa021967.
9. van Veer, L.J. 't, Dai, H., van de Vijver, M.J., et al. 2002. Gene expression profiling predicts clinical outcome of breast cancer. *Nature* 415, 530–536. doi:10.1038/415530a.
10. Cardoso F, van't Veer LJ, Bogaerts J, et al. 70-Gene Signature as an Aid to Treatment Decisions in Early-Stage Breast Cancer. *N Engl J Med.* 2016 Aug 25;375(8):717-29.
11. Piccart M, van 't Veer LJ, Poncet C, et al. 70-gene signature as an aid for treatment decisions in early breast cancer: updated results of the phase 3 randomised MINDACT trial with an exploratory analysis by age. *Lancet Oncol.* 2021 Apr;22(4):476-488. doi: 10.1016/S1470-2045(21)00007-3. Epub 2021 Mar 12. PMID: 33721561.

Oncotype DX

1. Albain, K.S., Barlow, W.E., Shak, S. et al. 2010. Prognostic and predictive value of the 21-gene recurrence score assay in postmenopausal women with node-positive, oestrogen-receptor-positive breast cancer on chemotherapy: a retrospective analysis of a randomised trial. *Lancet Oncol.* 11, 55–65. Cronin, M., Sangli, C., Liu, M.-L., et al. 2007. Analytical validation of the Oncotype DX genomic diagnostic test for recurrence prognosis and therapeutic response prediction in node-negative, estrogen receptor-positive breast cancer. *Clin. Chem.* 53, 1084–1091.
2. Dowsett, M., Cuzick, J., Wale, C. et al. 2010. Prediction of risk of distant recurrence using the 21-gene recurrence score in node-negative and node-positive postmenopausal patients with breast cancer treated with anastrozole or tamoxifen: a TransATAC study. *J.*

Clin. Oncol. 28, 1829–1834.

3. Geyer CE Jr, Tang G, Mamounas EP, et al. 21-Gene assay as predictor of chemotherapy benefit in HER2-negative breast cancer. *NPJ Breast Cancer*. 2018 Nov 14;4:37.
4. Gluz O, Nitz UA, Christgen M, et al.: West German Study Group Phase III PlanB Trial: First Prospective Outcome Data for the 21-Gene Recurrence Score Assay and Concordance of Prognostic Markers by Central and Local Pathology Assessment. *J Clin Oncol*. 2016 Jul 10;34(20):2341-9.
5. Kalinsky K, Barlow WE, Gralow JR et al. 21-Gene Assay to inform chemotherapy benefit from node positive breast cancer. *Engl J Med* DOI: 10.1056/NEJMoa2108873
6. Khan SS, Karn T, Symmans WF, et al. Genomic predictor of residual risk of recurrence after adjuvant chemotherapy and endocrine therapy in high risk estrogen receptor-positive breast cancers. *Breast Cancer Res Treat*. 2015 Feb 5.
7. Mamounas, E.P., Tang, G., Fisher, B., et al. 2010. Association between the 21-gene recurrence score assay and risk of locoregional recurrence in node-negative, estrogen receptor-positive breast cancer: results from NSABP B-14 and NSABP B-20. *J. Clin. Oncol*. 28, 1677–1683.
8. Nitz U, Gluz O, Christgen M, et al. Reducing chemotherapy use in clinically high-risk, genomically low-risk pN0 and pN1 early breast cancer patients: five-year data from the prospective, randomised phase 3 West German Study Group (WSG) PlanB trial. *Breast Cancer Res Treat*. 2017 Oct;165(3):573-583.
9. Nitz UA, Gluz O, Kümmel S et al. Endocrine Therapy Response and 21-Gene Expression Assay for Therapy Guidance in HR+/HER2- Early Breast Cancer. *J Clin Oncol*. 2022 Aug 10;40(23):2557-2567. doi: 10.1200/JCO.21.02759. Epub 2022 Apr 11.
10. Paik, S., Shak, S., Tang, G. et al. 2004. A multigene assay to predict recurrence of tamoxifen-treated, node-negative breast cancer. *N. Engl. J. Med*. 351, 2817–2826. doi:10.1056/NEJMoa041588.
11. Paik, S., Tang, G., Shak, S., et al. 2006. Gene expression and benefit of chemotherapy in women with node-negative, estrogen receptor-positive breast cancer. *J. Clin. Oncol*. 24, 3726–3734.
12. Penault-Llorca F, Filleron T, Asselain B, et al. The 21-gene Recurrence Score® assay predicts distant recurrence in lymph node-positive, hormone receptor-positive, breast cancer patients treated with adjuvant sequential epirubicin- and docetaxel-based or epirubicin-

based chemotherapy (PACS-01 trial). BMC Cancer. 2018 May 4;18(1):526.

13. Sparano JA, Gray RJ, Makower DF, et al: Prospective Validation of a 21-Gene Expression Assay in Breast Cancer. N Engl J Med 2015, 373(21):2005-2014.
14. Sparano JA, Gray R, Oktay MH, et al. A metastasis biomarker (MetaSite Breast™ Score) is associated with distant recurrence in hormone receptor-positive, HER2-negative early-stage breast cancer. NPJ Breast Cancer. 2017 Nov 8;3:42.
15. Sparano JA, Gray RJ, Makower DF, et al. Adjuvant Chemotherapy Guided by a 21-Gene Expression Assay in Breast Cancer. N Engl J Med. 2018 Jul 12;379(2):111-121.
16. Tang, G., Cuzick, J., Costantino, J.P., et al. 2011. Risk of recurrence and chemotherapy benefit for patients with node-negative, estrogen receptor-positive breast cancer: recurrence score alone and integrated with pathologic and clinical factors. J. Clin. Oncol. 29, 4365–4372.
17. Zhao X, Rødland EA, Sørli T, et al. Systematic assessment of prognostic gene signatures for breast cancer shows distinct influence of time and ER status. BMC Cancer. 2014 Mar 19;14:211.
18. Sparano J, Gray RJ, Makower D, Albain KS, Hayes DF, Geyer C, Dees E, Goetz MP, Olson Jr JA, Lively TG, Badve S, Saphner T, Wagner LI, Whelan T, Kaklamani V, Sledge Jr GW. Trial Assigning Individualized Options for Treatment (TAILORx): An update including 12-year event rates. SABCS 2022, GS 1-05

Prosigna (ROR / PAM50)


1. Chia, S.K., Bramwell, V.H., Tu, Det al. 2012. A 50-gene intrinsic subtype classifier for prognosis and prediction of benefit from adjuvant tamoxifen. Clin. Cancer Res. 18, 4465–4472.
2. Gnant M, Filipits M, Greil R, et al. Austrian Breast and Colorectal Cancer Study Group. Predicting distant recurrence in receptor-positive breast cancer patients with limited clinicopathological risk: using the PAM50 Risk of Recurrence score in 1478 postmenopausal patients of the ABCSG-8 trial treated with adjuvant endocrine therapy alone. Ann Oncol. 2014 Feb;25(2):339-45
3. Jensen MB, Lænkholm AV, Nielsen TO, et al. The Prosigna gene expression assay and responsiveness to adjuvant cyclophosphamide-

based chemotherapy in premenopausal high-risk patients with breast cancer. *Breast Cancer Res.* 2018 Jul 27;20(1):79.

4. Lænkholm AV, Jensen MB, Eriksen JO, et al. PAM50 Risk of Recurrence Score Predicts 10-Year Distant Recurrence in a Comprehensive Danish Cohort of Postmenopausal Women Allocated to 5 Years of Endocrine Therapy for Hormone Receptor-Positive Early Breast Cancer. *J Clin Oncol.* 2018 Mar 10;36(8):735-740.
5. Liu S, Chapman JA, Burnell MJ, et al. Prognostic and predictive investigation of PAM50 intrinsic subtypes in the NCIC CTG MA.21 phase III chemotherapy trial. *Breast Cancer Res Treat.* 2015 Jan;149(2):439-48
Nielsen, T.O., Parker, J.S., Leung, S., et al. 2010. A comparison of PAM50 intrinsic subtyping with immunohistochemistry and clinical prognostic factors in tamoxifen-treated estrogen receptor-positive breast cancer. *Clin. Cancer Res.* 16, 5222–5232.
6. Ohnstad HO, Borgen E, Falk RS, et al. Prognostic value of PAM50 and risk of recurrence score in patients with early-stage breast cancer with long-term follow-up. *Breast Cancer Res.* 2017 Nov 14;19(1):120.
7. Parker, J.S., Mullins, M., Cheang, M.C.U., et al. 2009. Supervised risk predictor of breast cancer based on intrinsic subtypes. *J. Clin. Oncol.* 27, 1160–1167.
8. Prat, A., Cheang, M.C.U., Martín, M., et al. 2012b. Prognostic Significance of Progesterone Receptor-Positive Tumor Cells Within Immunohistochemically Defined Luminal A Breast Cancer. *J. Clin. Oncol.*
9. Prat, A., Parker, J.S., Fan, C., et al. 2012a. Concordance among gene expression-based predictors for ER-positive breast cancer treated with adjuvant tamoxifen. *Ann. Oncol.* 23, 2866–2873.
10. Perou, C.M., Sørlie, T., Eisen, M.B. et al. 2000. Molecular portraits of human breast tumours. *Nature* 406, 747–752.
11. Pogue-Geile KL, Song N, Jeong JH, et al. Intrinsic Subtypes, PIK3CA Mutation, and the Degree of Benefit From Adjuvant Trastuzumab in the NSABP B-31 Trial. *J Clin Oncol.* 2015 Jan 5.
12. Sestak I, Cuzick J, Dowsett M. et al.. Prediction of Late Distant Recurrence After 5 Years of Endocrine Treatment: A Combined Analysis of Patients From the Austrian Breast and Colorectal Cancer Study Group 8 and Arimidex, Tamoxifen Alone or in Combination Randomized Trials Using the PAM50 Risk of Recurrence Score. *J Clin Oncol.* 2014 Oct 20. pii: JCO.2014.55.6894

Breast Cancer Index

1. Bartlett JMS, Sgroi DC, Treuner K et al. Breast Cancer Index and prediction of benefit from extended endocrine therapy in breast cancer patients treated in the Adjuvant Tamoxifen-To Offer More? (aTTom) trial. *Ann Oncol.* 2019 Nov 1;30(11):1776-1783.
2. Jerevall PL, Brommesson S, Strand C, et al. Exploring the two-gene ratio in breast cancer--independent roles for HOXB13 and IL17BR in prediction of clinical outcome. *Breast Cancer Res Treat.* 2008;107(2):225–234.
3. Jansen MP, Sieuwerts AM, Look MP, et al. HOXB13-to-IL17BR expression ratio is related with tumor aggressiveness and response to tamoxifen of recurrent breast cancer: a retrospective study. *J Clin Oncol.* 2007;25(6):662–668.
4. Ma XJ, Hilsenbeck SG, Wang W, et al. The HOXB13:IL17BR expression index is a prognostic factor in early-stage breast cancer. *J Clin Oncol.* 2006;24(28):4611–4619.



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2023.1E

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

Prospective Clinical Trials (Oncotype DX® [TAILORx, PlanB, RxPONDER, ADAPT], MammaPrint® [MINDACT])

Prognosis in low-risk groups excellent for both tests: ~ 94% 5 J. DFS with only adjuvant endocrine therapy (ET)

	TailorX	RxPONDER	PlanB	ADAPT	MINDACT
Follow-up	median 7.5 years	median 5.1 years	5-year-DFS	median 60 months	median 8.7 years
Trial design (biomarker question)	pN0; Randomization RS 11-25 (+/- CTX)	pN1; Randomization RS0-25 (+/- CTX)	Prospective ODX testing: ET alone in RS 0-11 pN0-1	Non-inferiority (iDFS) ET alone: RS 0-11 vs RS12-25/ET response	Prospectively defined 5y-DMFS threshold for ET alone
Percentage clinically defined low-risk group	6615/9427 (70.2%, adj-online)	all 1-3 involved lymph nodes	all clinical CTX indication (pN0-1)	all clinical chemotherapy (CTX) indication (c/pN0-1)	3336/ 6693 (49.8%, adj-online)
Percentage high clinical risk and low genomic risk (clinical CTX indication)	16.7% (RS 0-10)	42.8% (RS 0-13)	15.3% (RS 0-11)	ET-trial (pN0-1): all RS 0-25, i.e. low genomic risk with ET alone	23.2% (high clinical/low genomic risk)
Test failure rate	n.r.	n.r.	2.9%	n.r.	26% (fresh frozen)
Percentage genomically intermediate-risk group (only for Oncotype DX, ODX)	69.1% (RS 11-25)	57.2% (RS 14-24)	60.4% (RS 12-25)	Included only RS 0-11 (37.9%) or RS 12-25/ET response (62.1%)	n.a.
Percentage genomically high-risk group (only for Oncotype DX)	14.3% (RS ≥ 26)	n.a.	24.3% (RS ≥ 26)	n.a.	27.0% (high clinical and high genomic risk)
12-year follow-up	reported	n.r.	n.r.	n.r.	n.r.

TailorX

1. Sparano JA, Gray RJ, Makower DF, et al. Prospective Validation of a 21-Gene Expression Assay in Breast Cancer. N Engl J Med. 2015 Nov 19;373(21):2005
2. Sparano JA, Gray RJ, Makower DF, et al. Adjuvant Chemotherapy Guided by a 21-Gene Expression Assay in Breast Cancer. N Engl J Med. 2018 Jul 12;379(2):111-121.
3. Sparano JA, Gray RJ, Makower DF, et al. 12-Year Recurrence and Survival Outcomes for Patients With Early-Stage Breast Cancer. SABCS 2022. GS1-05

RxPONDER

1. Kalinsky K, Barlow WE, Gralow JR et al. 21-Gene Assay to inform chemotherapy benefit in node-positive breast cancer. N Engl J Med DOI: 10.1056/NEJMoa2108873

Plan B

1. Gluz O, Nitz UA, Christgen M, et al. West German Study Group Phase III PlanB Trial: First Prospective Outcome Data for the 21-Gene Recurrence Score Assay and Concordance of Prognostic Markers by Central and Local Pathology Assessment. J Clin Oncol. 2016 Jul 10;34(20):2341-9.

2. Nitz U, Gluz O, Christgen M, et al. Reducing chemotherapy use in clinically high-risk, genomically low-risk pN0 and pN1 early breast cancer patients: five-year data from the prospective, randomised phase 3 West German Study Group (WSG) PlanB trial. *Breast Cancer Res Treat.* 2017 Oct;165(3):573-583.

ADAPT

1. Gluz O, Nitz UA, Christgen M, et al. Impact of age, recurrence score (RS) and ovarian function suppression (OFS) on endocrine response to short preoperative endocrine therapy (ET): Analysis of ADAPT and ADAPTCycle trials. *Annals of Oncology* (2022) 33 (suppl_7): S808-S869. 10.1016/annonc/annonc1089
2. Nitz UA, Gluz O, Kümmel S et al. Endocrine Therapy Response and 21-Gene Expression Assay for Therapy Guidance in HR+/HER2- Early Breast Cancer. *J Clin Oncol.* 2022 Aug 10;40(23):2557-2567. doi: 10.1200/JCO.21.02759. Epub 2022 Apr 11.

MINDACT

1. Cardoso F, van't Veer LJ, Bogaerts J, et al. 70-Gene Signature as an Aid to Treatment Decisions in Early-Stage Breast Cancer. *N Engl J Med.* 2016 Aug 25;375(8):717-29.
2. Piccart M, van 't Veer LJ, Poncet C, et al. 70-gene signature as an aid for treatment decisions in early breast cancer: updated results of the phase 3 randomised MINDACT trial with an exploratory analysis by age. *Lancet Oncol.* 2021 Apr;22(4):476-488. doi: 10.1016/S1470-2045(21)00007-3. Epub 2021 Mar 12. PMID: 33721561.

Several tests

1. Bartlett JM, Bayani J, Marshall A, et al; OPTIMA TMG. Comparing Breast Cancer Multiparameter Tests in the OPTIMA Prelim Trial: No Test Is More Equal Than the Others. *J Natl Cancer Inst.* 2016 Apr 29;108(9).
2. Varga Z, Sinn P, Seidman AD. Summary of head-to-head comparisons of patient risk classifications by the 21-gene Recurrence Score® (RS) assay and other genomic assays for early breast cancer. *Int J Cancer.* 2019 Aug 15;145(4):882-893.
3. Berchtold E, Vetter M, Gündert M et al. Comparison of six breast cancer classifiers using qPCR. *Bioinformatics.* 2019 Sep 15;35(18):3412-3420.
4. Sestak I, Buus R, Cuzick J, et al. Comparison of the Performance of 6 Prognostic Signatures for Estrogen Receptor-Positive Breast Cancer: A Secondary Analysis of a Randomized Clinical Trial. *JAMA Oncol.* 2018 Apr 1;4(4):545-553.

Adjuvant Endocrine Therapy

Predictive Factors for DFS

Therapy	Factor	Oxford		
		LoE	GR	AGO
▪ Endocrine therapy	▪ ER / PR status [%]	1a	A	++
	▪ IHC staining intensity (ER/PR)	1a	A	-
	▪ Ki-67 Re-Evaluation after short preoperative endocrine therapy (2-4 weeks) (ypT and ypN)*	1b	A	+
▪ Extended endocrine therapy (EAT)	▪ Breast Cancer Index® MammaPrint	2b	B	+/-
▪ Tamoxifen	▪ CYP2D6-polymorphism	2b	B	-
▪ Ovarian ablation or suppression	▪ Menopausal status	1c	A	++
▪ Aromatase inhibitors vs. tamoxifen	▪ Menopausal status	1c	A	++
	▪ ER / PR / HER2 as single factors	1c	A	-
	▪ Invasiv-lobular breast cancer	2b	B	+
	▪ Ki-67 high	2b	B	+/-
	▪ Obesity (BMI > 30 kg/m²)	2b	B	+/-

General publications

1. Clark GM et al. Prognostic and predictive factors. In: Diseases of the breast, 2nd edition: Seiten 489-514. Harris JR, Lippmann ME, Morrow M, Osborne CK (Hrsg). Lippincott-Raven Publishers, Philadelphia 2000.
2. Simon RM, Paik S, Hayes DF. Use of archived specimens in evaluation of prognostic and predictive biomarkers. J. Natl. Cancer Inst. 2009; 101(21): 1446 – 1452
3. Coates AS, Winer EP, Goldhirsch A, et al. Tailoring therapies--improving the management of early breast cancer: St Gallen International Expert Consensus on the Primary Therapy of Early Breast Cancer 2015. Ann Oncol. 2015 Aug;26(8):1533-46.

Endocrine therapy

1. Colleoni M et al.: Tamoxifen after adjuvant chemotherapy for premenopausal women with lymph node-positive breast cancer: International Breast Cancer Study Group Trial 13-93. J Clin Oncol 24 (9): 1332-41, 2006.
2. Early Breast Cancer Trialists' Collaborative Group (EBCTCG): Effects of chemotherapy and hormonal therapy for early breast cancer on recurrence and 15-year survival: an overview of the randomised trials. Lancet 365 (9472): 1687-717, 2005
3. Early Breast Cancer Trialists' Collaborative Group (EBCTCG), Davies C, Godwin J, et al. Relevance of breast cancer hormone receptors and other factors to the efficacy of adjuvant tamoxifen: patient-level meta-analysis of randomised trials. Lancet. 2011 Aug 27;378(9793):771-84

4. Harvey JM, Clark GM, Osborne CK, et al.: Estrogen receptor status by immunohistochemistry is superior to the ligand-binding assay for predicting response to adjuvant endocrine therapy in breast cancer. *J Clin Oncol* 17 (5): 1474-81, 1999.
5. Thürliman B et al: Is chemotherapy necessary for premenopausal women with lower-risk node-positive, endocrine responsive breast cancer? 10-year update of International Breast Cancer Study Group Trial 11-93. *Breast Cancer Res Treat.* 2009; 113:137-44
6. Ellis MJ, Suman VJ, Hoog J, et al. Ki67 Proliferation Index as a Tool for Chemotherapy Decisions During and After Neoadjuvant Aromatase Inhibitor Treatment of Breast Cancer: Results From the American College of Surgeons Oncology Group Z1031 Trial (Alliance). *J Clin Oncol.* 2017 Apr 1;35(10):1061-1069.
7. Dowsett M, Ellis MJ, Dixon JM, et al. Evidence-based guidelines for managing patients with primary ER+ HER2- breast cancer deferred from surgery due to the COVID-19 pandemic. *Breast Cancer.* 2020 Jun 8;6:21.
8. Smith I, Robertson J, Kilburn L, et al. Long-term outcome and prognostic value of Ki67 after perioperative endocrine therapy in postmenopausal women with hormone-sensitive early breast cancer (POETIC): an open-label, multicentre, parallel-group, randomised, phase 3 trial. *Lancet Oncol.* 2020 Nov;21(11):1443-1454.

EAT

1. Bartlett JMS, SgROI DC, Treuner K et al. Breast Cancer Index and prediction of benefit from extended endocrine therapy in breast cancer patients treated in the Adjuvant Tamoxifen-To Offer More? (aTTom) trial. *Ann Oncol.* 2019 Nov 1;30(11):1776-1783.
2. Rastogi, P., et al. (2021). "Utility of the 70-gene MammaPrint assay for prediction of benefit from extended letrozole therapy (ELT) in the NRG Oncology/NSABP B-42 trial." *Journal of Clinical Oncology* **39**(15_suppl): 502-502.
3. Liefers G-J, Meershoek-Klein Kranenbarg E, Duijm-de Carpentier M et al. Utility of the 70-gene MammaPrint test for prediction of extended endocrine therapy benefit in patients with early-stage breast cancer in the IDEAL Trial, SABCS 2022 GS5-10

Amenorrhoea

1. Anders C, Marcom PK, Peterson B, et al. A pilot study of predictive markers of chemotherapy-related amenorrhea among premenopausal women with early stage breast cancer. *Cancer Invest.* 2008 Apr-May;26(3):286-95
2. Anderson RA, Cameron DA. Pretreatment serum anti-müllerian hormone predicts long-term ovarian function and bone mass after chemotherapy for early breast cancer. *J Clin Endocrinol Metab.* 2011 May;96(5):1336-43.

Body Mass Index

1. Chan DSM et al. Body mass index and survival in women with breast cancer—systematic literature review and meta-analysis of 82 follow-up studies *Ann Oncol.* Oct 2014; 25(10): 1901–1914.
2. Xia X, Chen W, Li J, et al. Body mass index and risk of breast cancer: a nonlinear dose-response meta-analysis of prospective studies. *Sci Rep.* 2014 Dec 15;4:7480. doi: 10.1038/srep07480.

CYP2D6

1. Bezerra LS, Santos-Veloso MAO, Bezerra Junior NDS, et al. Impacts of Cytochrome P450 2D6 (CYP2D6) Genetic Polymorphism in Tamoxifen Therapy for Breast Cancer. *Rev Bras Ginecol Obstet.* 2018 Dec;40(12):794-799.
2. Brooks JD, Comen EA, Reiner AS, et al; WECARE Study collaborative group, Malone KE, Bernstein JL. CYP2D6 phenotype, tamoxifen, and risk of contralateral breast cancer in the WECARE Study. *Breast Cancer Res.* 2018 Dec 10;20(1):149.
3. Bai Y, Wu HW, Zhang YH. Effects of CYP2D6*10 polymorphism on tamoxifen pharmacokinetics in patients with breast cancer in Asia: a meta-analysis. *Cancer Chemother Pharmacol.* 2018 Oct 24. doi: 10.1007/s00280-018-3703-8. [Epub ahead of print] PMID: 30357449
4. Schroth W, Winter S, Mürdter T, et al. Improved Prediction of Endoxifen Metabolism by CYP2D6 Genotype in Breast Cancer Patients Treated with Tamoxifen. *Front Pharmacol.* 2017 Aug 24;8:582. doi: 10.3389/fphar.2017.00582. eCollection 2017. PMID: 28955222
5. Hertz DL, Kidwell KM, Hilsenbeck SG, et al. CYP2D6 genotype is not associated with survival in breast cancer patients treated with tamoxifen: results from a population-based study. *Breast Cancer Res Treat.* 2017 Nov;166(1):277-287.
6. Hwang GS, Bhat R, Crutchley RD, et al. Impact of CYP2D6 polymorphisms on endoxifen concentrations and breast cancer outcomes. *Pharmacogenomics J.* 2018 Apr;18(2):201-208.

Ki-67 Bestimmung nach kurzer präoperativer endokriner Therapie

1. Dowsett M, Torsten O, A´Hern R et al. Assessment of Ki-67 in Breast Cancer: Recommendations from the International Ki67 Breast Cancer Working Group. *J Natl Cancer Inst* 2011;103:1656-1664
2. Kim HJ, Noh WC, Lee ES et al. Efficacy of neoadjuvant endocrine therapy compared with neoadjuvant chemotherapy in premenopausal patients with estrogen receptor positive and HER2-negative lymph-node positive breast cancer. *Breast Cancer Res* 202;22(1):54-59.
3. Ellis MJ, Suman VJ, Hoog J et al. Ki-67 Proliferation Index as a tool for chemotherapy decision during and after neoadjuvant aromatase inhibitor treatment of breast cancer: Results from the American College of Surgeons Oncology Group Z1031 Trial

(Alliance). *J Clin Oncol* 2017;35(10):1061-1069.

4. Smith, I., et al. (2020). "Long-term outcome and prognostic value of Ki67 after perioperative endocrine therapy in postmenopausal women with hormone-sensitive early breast cancer (POETIC): an open-label, multicentre, parallel-group, randomised, phase 3 trial." The Lancet. Oncology **21**(11): 1443-1454.
5. Nitz, U. A., et al. (2022). "Endocrine Therapy Response and 21-Gene Expression Assay for Therapy Guidance in HR+/HER2- Early Breast Cancer." J Clin Oncol **40**(23): 2557-2567.

Adjuvant Chemotherapy and Targeted Therapy Predictive Factors for DFS

Therapy	Factor	Oxford		
		LoE	GR	AGO
■ Adjuvant Chemotherapy	70-Gene-signature (Mammaprint®)	1b	A	+
	21-Gene-signature (Oncotype DX RS®)	1b	A	+
	EPclin (Endopredict®)	2b	B	+
	PAM-50 (Prosigna®)	2b	B	+
	Histological type (lobular vs. NST)	2b	B	-
	TIL's in TNBC	2b	B	+/-
■ Anti-HER2-Therapy	HER2 (IHC, ISH)	1a	A	++
■ PARP-Inhibitors	gBRCA1/Mutation (HER2 neg.)	1a	A	+

*Consider decision according to age/menopausal status, prospective evidence available for Mammaprint and OncotypeDX only (see next slide)

70-Gene-Signature (Mammaprint®)

- Cardoso F, van't Veer LJ, Bogaerts J, et al. 70-Gene Signature as an Aid to Treatment Decisions in Early-Stage Breast Cancer. *N Engl J Med.* 2016;375(8):717–729.
- Cardoso F, van 't Veer L, Poncet C, et al. MINDACT: Long-term results of the large prospective trial testing the 70-gene signature MammaPrint as guidance for adjuvant chemotherapy in breast cancer patients. ASCO 2020, #506

OncotypeDX

- Paik, S., Tang, G., Shak, S., et al. 2006. Gene expression and benefit of chemotherapy in women with node-negative, estrogen receptor-positive breast cancer. *J. Clin. Oncol.* 24, 3726–3734.
- Sparano JA, Gray RJ, Makower DF, et al. Clinical Outcomes in Early Breast Cancer With a High 21-Gene Recurrence Score of 26 to 100 Assigned to Adjuvant Chemotherapy Plus Endocrine Therapy: A Secondary Analysis of the TAILORx Randomized Clinical Trial [published online ahead of print, 2019 Sep 30]. *JAMA Oncol.* 2019;e194794.
- Kalinsky K, Barlow WE, Meric-Bernstam F, et al. First results from a phase III randomized clinical trial of standard adjuvant endocrine therapy (ET) +/- chemotherapy (CT) in patients (pts) with 1-3 positive nodes, hormone receptor-positive (HR+) and HER2-negative (HER2-) breast cancer (BC) with recurrence score (RS) < 25: SWOG S1007 (RxPonder). SABCS 2020, GS3-00
- Harbeck N, Gluz O, Kuemmel S, et al. Endocrine therapy alone in patients with intermediate or high-risk luminal early breast cancer

(0-3 lymph nodes), Recurrence Score <26 and Ki67 response after preoperative endocrine therapy: Primary outcome results from the WSG-ADAPT HR+/HER2- trial. SABCS 2020, GS4-04.

EPclin (EndoPredict®)

1. Sestak I, Martín M, Dubsky P et al. Prediction of chemotherapy benefit by EndoPredict in patients with breast cancer who received adjuvant endocrine therapy plus chemotherapy or endocrine therapy alone. Breast Cancer Res Treat. 2019 Jul;176(2):377-386.

PAM-50 (Prosigna®)

1. Prat A, Galván P, Jimenez B et al. Prediction of Response to Neoadjuvant Chemotherapy Using Core Needle Biopsy Samples with the Prosigna Assay. Clin Cancer Res. 2016 Feb 1;22(3):560-6.
2. Jensen MB, Lærnholm AV, Nielsen TO et al. The Prosigna gene expression assay and responsiveness to adjuvant cyclophosphamide-based chemotherapy in premenopausal high-risk patients with breast cancer. Breast Cancer Res. 2018 Jul 27;20(1):79.

Histological type:

1. De Nonneville A, Jauffret C, Goncalves A. Adjuvant chemotherapy in lobular carcinoma of the breast: a clinicopathological score identifies high-risk patient with survival benefit Breast Cancer Res Treat. 2019 Jun;175(2):379-387.
2. Fu R, Yang J, Wang H et al.: A nomogram for determining the disease-specific survival in invasive lobular carcinoma of the breast: A population study. Medicine (Baltimore). 2020 Oct 23;99(43):e22807.

Anti-HER2 therapy

see evidence in chapter “Chemotherapy and targeted therapy”

PARPi

1. Tutt ANJ, Garber JE, Kaufmann B et al. Adjuvant Olaparib for patients with BRCA1- or BRCA2 mutated Breast Cancer. N Engl J Med 2021;384(25):2394.
2. Garber HR, Litton JR. Integrating poly(ADP-ribose) polymerase inhibitors in the treatment of early breast cancer. Curr Opin Oncol 2019;31(3):247-255.



© AGO e. V.
in der DGGO e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2023.1E

www.ago-online.de

FORSCHEN
LEBEN
HEILEN

Results for prospectively evaluated biomarkers (LOE1a) in early HR+/HER2- breast cancer

biomarker/ signature	Population (HR+/HER2- patients)	therapy options
Mammaprint (MINDACT n=2140)	Clinically high/genomic low risk (n=1550) N0-1, age >50 yrs N0-1, age ≤50 yrs (patients with OFS in the ET arm: 26%)	ET, no adjuvant CT adjuvant CT → ET*: 2.6% CT-benefit in 5-y DDFS (93.6 vs. 96.2%)
Oncotype DX (TAILORx n=6711)	TailorX (T1b-T2, N0, 74% clinically low risk, 13% OFS in premenopausal women) N0, RS 0-25 age >50 yrs. N0 RS 0-15 age ≤50 yrs N0 RS 16-25 age ≤50 yrs	ET, no adjuvant CHT ET, no adjuvant CHT adjuvant CT → ET*: (3.2-3.4% CT-benefit in 5-y DRFI (93→95-96% 5 y DRFI, in RS 16-20 if clinical high risk only, 16-20: HR=1.4 (n.s.), 21-25: HR=2.19 (sign) for ET vs. CT → ET
RxPonder (n=5018)	RxPonder: N1 RS 0-25: postmenopausal RS 0-25: premenopausal (patients with OFS in the ET arm: 19%)	ET, no adjuvant CT (neo)adjuvant CT → ET* 2.4% CT benefit in 5-y DRFI (5-y DRFI 93.9 vs. 96.3%, HR=0.062, p=0.02) explorative analysis: no effect of CT age 50 and older (p interaction 0.06)
RS + Ki-67post (ADAPT, n=2290 endocrine treated)	clinically intermediate/high risk , RS 0-25 (RS 12, 25+Ki67post≤10%) N0-1, age >50 yrs N0, RS 0-11 and age ≤50 yrs N0, RS 12-25 with Ki67post≤10% and age ≤50 yrs N1: RS 0-25 (+ Ki-67post≤10% in RS 12-25) and age ≤50 yrs N1: RS 0-25 and ki-67post>10%	ET, no adjuvant CT adjuvant ET, no adjuvant CT adjuvant ET +/- OFS, if RS >16 or clinically high risk +/- CT: 5-yr-DDFS: 97% with ET alone, no significant difference between RS 0-15 and 16-25 adjuvant ET+OFS or CT → ET 5-yrs. DDFS 97% with ET alone (neo)adjuvant CT → ET

* If CT is refused: alternative ET+OFS

DDFS=distant-disease-free-survival, DRFI= distant recurrence free interval, ET= endocrine treatment, CT= chemotherapy, OFS= ovarian function suppression, RS= Recurrence Score

1. Sparano JA, Gray RJ, Ravdin PM et al. Clinical and Genomic Risk to Guide the Use of Adjuvant Therapy for Breast Cancer. New England Journal of Medicine 2019; 380: 2395-2405.
2. Sparano JA, Gray RJ, Makower DF et al. Adjuvant Chemotherapy Guided by a 21-Gene Expression Assay in Breast Cancer. New England Journal of Medicine 2018
3. Kalinsky KM, Barlow WE, Gralow JR et al. Abstract GS2-07: Updated results from a phase 3 randomized clinical trial in participants (pts) with 1-3 positive lymph nodes (LN), hormone receptor-positive (HR+) and HER2-negative (HER2-) breast cancer (BC) with recurrence score (RS) ≤ 25 randomized to endocrine therapy (ET) +/- chemotherapy (CT): SWOG S1007 (RxPONDER). Cancer Research 2022; 82: GS2-07-GS02-07.
4. Piccart M, van 't Veer LJ, Poncet C et al. 70-gene signature as an aid for treatment decisions in early breast cancer: updated results of the phase 3 randomised MINDACT trial with an exploratory analysis by age. The Lancet Oncology 2021; 22: 476-488.
5. Gluz O, Nitz U, Christgen M et al. Prognostic impact of recurrence score, endocrine response and clinical-pathological factors in high-

risk luminal breast cancer: Results from the WSG-ADAPT HR+/HER2- chemotherapy trial. Journal of Clinical Oncology 2021; 39: 504-504.

6. Nitz UA, Gluz O, Kummel S et al. Endocrine Therapy Response and 21-Gene Expression Assay for Therapy Guidance in HR+/HER2- Early Breast Cancer. J Clin Oncol 2022; 40: 2557-2567.

Neoadjuvant Systemic Chemotherapy (NACT)

Predictive Factors for pCR I

Factor	pCR* Probability	Oxford		
		LoE	GR	AGO
▪ Young age	↑	1a	A	+
▪ Obesity	↓	2a	B	+
▪ cT1 / cT2 tumors o. N0 o. G3	↑↑	1a	A	++
▪ Negative hormone receptor status	↑↑	1a	A	++
▪ Triple negative breast cancer	↑↑	1a	A	++
▪ Positive HER2-status	↑↑	1a	A	++
▪ Early clinical response	↑	1b	A	+
▪ Lobular tumor type	↓	1a	A	+
▪ Metaplastic tumor type	↓↓	4	C	+

* High (↑) or very high (↑↑) probability to reach pCR, low (↓) or very low (↓↓) probability to reach pCR
 See also chapter „Prognostic and predictive factors“

General evidence

1. Cortazar P, Zhang L, Untch M, et al. Pathological complete response and long-term clinical benefit in breast cancer: the CTNeoBC pooled analysis. *Lancet* 2014;384: 164-72.
2. Gerber B, Loibl S, Eidtmann H, et al. Neoadjuvant bevacizumab and anthracycline-taxane-based chemotherapy in 678 triple-negative primary breast cancers; results from the geparquinto study (GBG 44). *Ann Oncol* 2013;24: 2978-84.
3. van Mackelenbergh MT, Denkert C, Nekljudova V, et al. Outcome after neoadjuvant chemotherapy in estrogen receptor-positive and progesterone receptor-negative breast cancer patients: a pooled analysis of individual patient data from ten prospectively randomized controlled neoadjuvant trials. *Breast Cancer Res Treat* 2017.
4. von Minckwitz G, Eidtmann H, Rezai M, et al. Neoadjuvant chemotherapy and bevacizumab for HER2-negative breast cancer. *N Engl J Med* 2012;366: 299-309.
5. von Minckwitz G, Untch M, Blohmer JU, et al. Definition and impact of pathologic complete response on prognosis after neoadjuvant chemotherapy in various intrinsic breast cancer subtypes. *J Clin Oncol* 2012;30: 1796-804.

Body mass index

1. Wang H, Zhang S, Yee D, et al. Impact of body mass index on pathological complete response following neoadjuvant chemotherapy in operable breast cancer: a meta analysis. *Breast Cancer* 2021;28(3):616-629

Lobular cancer

1. Loibl S, Volz C, Mau C, et al. Response and prognosis after neoadjuvant chemotherapy in 1,051 patients with infiltrating lobular breast carcinoma. *Breast Cancer Res Treat* 2014;144: 153-62.

Metaplastic breast cancer

1. McMullen ER, Zoumberos NA, Kleer CG. Metaplastic Breast Carcinoma: Update on Histopathology and Molecular Alterations. *Arch Pathol Lab Med*. 2019 Dec;143(12):1492-1496.
2. Tzanninis IG, Kotteas EA, Ntanasis-Stathopoulos I et al. Management and Outcomes in Metaplastic Breast Cancer. *Clin Breast Cancer*. 2016 Dec;16(6):437-443.
3. Al-Hilli Z, Choong G, Keeney MG, et al. Metaplastic breast cancer has a poor response to neoadjuvant systemic therapy. *Breast Cancer Res Treat*. 2019;176(3):709–716.

Neoadjuvant Systemic Chemotherapy (NACT)

Predictive Factors for pCR II

Factor	pCR* Probability	Oxford		
		LoE	GR	AGO
■ Gene expression profiles (gene signatures) (Mammaprint®, Endopredict®, Oncotype DX®, Prosigna®, Breast Cancer Index SM)	↑	2b	B	+/-
■ Ki-67	↑	2b	B	+
■ Tumor infiltrating lymphocytes**	↑	2a	B	+
■ PIK3CA mutation (for HER2-positive BC)	↑	2a	B	+/-
■ gBRCA-mutation (for the effect of chemotherapy)	↑	2b	B	+
■ gBRCA-mutation (for the effect of platinum)	↔	2b	B	+/-

* High (↑) or very high (↑↑) probability of pCR, low (↓) or very low (↓↓) probability of pCR

** Defined as dense lymphocytic infiltration of inner peritumoral stroma outside of the invasion front (lymphocytes make up > 50% of stroma area)

TIL

1. Denkert, C., Loibl, S., Noske, A., et al. Tumor-associated lymphocytes as an independent predictor of response to neoadjuvant chemotherapy in breast cancer. 2010. J. Clin. Oncol. 28, 105–113. doi:10.1200/JCO.2009.23.7370.
2. Denkert C, von Minckwitz G, Brase JC, et al. Tumor-Infiltrating Lymphocytes and Response to Neoadjuvant Chemotherapy With or Without Carboplatin in Human Epidermal Growth Factor Receptor 2-Positive and Triple-Negative Primary Breast Cancers. J Clin Oncol. 2014 Dec 22. pii: JCO.2014.58.1967.
3. Ibrahim EM, Al-Foheidi ME, Al-Mansour MM, et al. The prognostic value of tumor-infiltrating lymphocytes in triple-negative breast cancer: a meta-analysis. Breast Cancer Res Treat. 2014 Dec;148(3):467-76
4. Loi S, Michiels S, Salgado R, et al. Tumor infiltrating lymphocytes are prognostic in triple negative breast cancer and predictive for trastuzumab benefit in early breast cancer: results from the FinHER trial. Ann Oncol. 2014 Aug;25(8):1544-50. Doi
5. Mao Y, Qu Q, Zhang Y, et al. The Value of Tumor Infiltrating Lymphocytes (TILs) for Predicting Response to Neoadjuvant Chemotherapy in Breast Cancer: A Systematic Review and Meta-Analysis. PLoS One. 2014 Dec 12;9(12)
6. Tung NM, Winer EP. Tumor-Infiltrating Lymphocytes and Response to Platinum in Triple-Negative Breast Cancer. J Clin Oncol. 2015 Jan 5. pii: JCO.2014.59.6031.

7. Denkert C, von Minckwitz G, Darb—Esfahani S et al. Tumour-infiltrating lymphocytes and prognosis in different subtypes of breast cancer: a pooled analysis of 3771 patients treated with neoadjuvant therapy. *Lancet Oncol* 2018;19(1):40-50.

PIK3CA

1. Loibl S, von Minckwitz G, Schneeweiss A, et al. PIK3CA mutations are associated with lower rates of pathologic complete response to anti-human epidermal growth factor receptor 2 (her2) therapy in primary HER2-overexpressing breast cancer. *J Clin Oncol*. 2014 Oct
2. Nuciforo PG, Aura C, Holmes E, et al: Benefit to neoadjuvant anti-Human Epidermal Growth Factor Receptor 2 (HER2)-targeted therapies in HER2-positive primary breast cancer is independent of Phosphatase and tensin homolog deleted from chromosome 10 (PTEN) status. *Ann Oncol*. 2015 Jul;26(7):1494-500. doi: 10.1093/annonc/mdv175. Epub 2015 Apr 7.
3. Pogue-Geile KL, Song N, Jeong JH, et al: Intrinsic Subtypes, PIK3CA Mutation, and the Degree of Benefit From Adjuvant Trastuzumab in the NSABP B-31 Trial. *Clin Oncol*. 2015 Jan 5. pii: JCO.2014.56.2439
4. Sueta A, Yamamoto Y, Yamamoto-Ibusuki M, et al. An Integrative Analysis of PIK3CA Mutation, PTEN, and INPP4B Expression in Terms of Trastuzumab Efficacy in HER2-Positive Breast Cancer. *PLoS One*. 2014 Dec 26;9(12):e116054.
5. Loibl S, Majewski I, Guarneri V, et al. PIK3CA mutations are associated with reduced pathological complete response rates in primary HER2-positive breast cancer: pooled analysis of 967 patients from five prospective trials investigating lapatinib and trastuzumab. *Ann Oncol*. 2016 Aug;27(8):1519-25.
6. Loibl S, Majewski I, Guarneri V, et al. PIK3CA mutations are associated with reduced pathological complete response rates in primary HER2-positive breast cancer: pooled analysis of 967 patients from five prospective trials investigating lapatinib and trastuzumab. *Ann Oncol*. 2019 Jan 9. doi: 10.1093/annonc/mdy536. [Epub ahead of print]
7. Fan H, Li C, Xiang Q, et al. PIK3CA mutations and their response to neoadjuvant treatment in early breast cancer: A systematic review and meta-analysis. *Thorac Cancer*. 2018 May;9(5):571-579.
8. Zardavas D, Te Marvelde L, Milne RL, et al. Tumor PIK3CA Genotype and Prognosis in Early-Stage Breast Cancer: A Pooled Analysis of Individual Patient Data. *J Clin Oncol*. 2018 Apr 1;36(10):981-990.

gBRCA bei TNBC

1. Loibl S, Weber KE, Timms KM et al. Survival analysis of carboplatin added to an anthracycline/taxane-based neoadjuvant chemotherapy and HRD score as predictor of response-final results from GeparSixto. Ann Oncol. 2018 Dec 1;29(12):2341-2347.

Metastatic Breast Cancer (mBC)

Prognostic Factors

Factor	Oxford		
	LoE	GR	AGO
<ul style="list-style-type: none"> Circulating tumor cells (CTC in blood, Cell Search®) <ul style="list-style-type: none"> Prognosis Early response assessment (3w) Therapy decision solely based on dynamics of CTC numbers over time or CTC phenotype Cell-free DNA (cfDNA in blood) 	 1a 1b 1b 2a	 A B A A	 + + -* +/-

* Study participation recommended

CTC

1. Bidard FC, Peeters DJ, Fehm T, et al. 2014. Clinical validity of circulating tumour cells in patients with metastatic breast cancer: a pooled analysis of individual patient data. *Lancet Oncol.* 2014 Apr;15(4):406-14.
2. Cristofanilli, M., Budd, G.T., Ellis, M.J., et al. 2004. Circulating tumor cells, disease progression, and survival in metastatic breast cancer. *N. Engl. J. Med.* 351, 781–791. doi:10.1056/NEJMoa040766.
3. Cristofanilli, M., Hayes, D.F., Budd, G.T. et al 2005. Circulating tumor cells: a novel prognostic factor for newly diagnosed metastatic breast cancer. *J. Clin. Oncol.* 23, 1420–1430. doi:10.1200/JCO.2005.08.140.
4. Giuliano, M., Giordano, A., Jackson, S., et al. 2011. Circulating tumor cells as prognostic and predictive markers in metastatic breast cancer patients receiving first-line systemic treatment. *Breast Cancer Res.* 13, R67. doi:10.1186/bcr2907.
5. Smerage JB, Barlow WE, Hortobagyi GN, et al. 2014. Circulating tumor cells and response to chemotherapy in metastatic breast cancer: SWOG S0500. *J Clin Oncol.* 2014 Nov, 1;32(31):3483-9
6. Sparano J, O'Neill A, Alpaugh K, et al. Association of Circulating Tumor Cells With Late Recurrence of Estrogen Receptor-Positive Breast Cancer: A Secondary Analysis of a Randomized Clinical Trial. *JAMA Oncol.* 2018 Dec 1;4(12):1700-1706.
7. Jauch SF, Riethdorf S, Sprick MR, et al. Sustained prognostic impact of circulating tumor cell status and kinetics upon further progression of metastatic breast cancer. *Breast Cancer Res Treat.* 2018 Oct 1. doi: 10.1007/s10549-018-4972-y. [Epub ahead of print] PMID: 30276763

Cell-free DNA

1. Cheng J, Holland-Letz T, Wallwiener M, et al. Circulating free DNA integrity and concentration as independent prognostic markers in metastatic breast cancer. *Breast Cancer Res Treat*. 2018 May;169(1):69-82.
2. Yang J, Cheng L, Zhang J, et al. Predictive value of circulating cell-free DNA in the survival of breast cancer patients: A systemic review and meta-analysis. *Medicine (Baltimore)*. 2018 Jul 97(28):e11417.
3. Zill OA, Banks KC, Fairclough SR, et al. The Landscape of Actionable Genomic Alterations in Cell-Free Circulating Tumor DNA from 21,807 Advanced Cancer Patients. *Clin Cancer Res*. 2018 Aug 1;24(15):3528-3538.

Treatment of Metastatic Breast Cancer

Predictive Factors for response

Therapy	Factor	Oxford		
		LoE	GR	AGO
▪ Endocrine therapy	ER / PR (prim. tumor, better: metastasis)	1a	A	++
	Response to prior therapy	2b	B	++
	Autocrine receptor mutation (<i>ESR1</i>)	2b	B	+
▪ Alpelisib	<i>PIK3CA</i> mutation (prim. tumor, metastases, plasma)	1b	A	++
▪ Trastuzumab Deruxtecan	HER2-low or HER2-positive	1b	A	++
▪ Chemotherapy	Response to prior therapy	1b	A	++
▪ Anti-HER2-therapy	HER2 (prim. tumor, better: metastasis)	1a	A	++
▪ Checkpoint-Inhibitors	PD-L1 positivity [#] (PD-L1c, CPS) in TNBC (primary tumor or metastasis)	1b	B	++
▪ PARP-Inhibitors	<i>gBRCA1/2</i> -mutation	1a	A	++
▪ Bone modifying drugs	Bone metastasis	1a	A	++

see chapter „pathology“

Endocrine therapy

1. Campbell FC, Blamey RW, Elston CW, et al. Quantitative oestradiol receptor values in primary breast cancer and response of metastases to endocrine therapy. *Lancet*. 1981;2(8259):1317–1319.

Endocrine therapy - ESR1:

1. Dustin D, Gu G, Fuqua SAW (2019) ESR1 mutations in breast cancer. *Cancer* 125:3714-3728 doi: 10.1002/cncr.32345.
2. Fribbens C, Garcia Murillas I, Beaney M et al. (2018) Tracking evolution of aromatase inhibitor resistance with circulating tumour DNA analysis in metastatic breast cancer. *Ann Oncol*.29:145-153. doi: 10.1093/annonc/mdx483
3. Fribbens C, O'Leary B, Kilburn L et al. (2016) Plasma ESR1 Mutations and the Treatment of Estrogen Receptor-Positive Advanced Breast Cancer. *J Clin Oncol*. 34:2961-8. doi: 10.1200/JCO.2016.67.3061

Alpelisib

1. André F, Ciruelos E, Rubovszky G et al. (2019) Alpelisib for *PIK3CA*-Mutated, Hormone Receptor-Positive Advanced Breast Cancer. *N Engl J Med*. 380:1929-1940. doi: 10.1056/NEJMoa1813904

Chemotherapy

1. Cardoso F, Senkus E, Costa A, et al. 4th ESO-ESMO International Consensus Guidelines for Advanced Breast Cancer (ABC 4)[†]. *Ann Oncol*. 2018;29(8):1634–1657.

Anti-HER2-Therapy

1. Seidman AD, Fornier MN, Esteva FJ, et al. Weekly trastuzumab and paclitaxel therapy for metastatic breast cancer with analysis of efficacy by HER2 immunophenotype and gene amplification. *J Clin Oncol*. 2001;19(10):2587–2595.

Checkpoint-Inhibitors

1. Schmid P, Adams S, Rugo HS, et al. Atezolizumab and Nab-Paclitaxel in Advanced Triple-Negative Breast Cancer. *N Engl J Med*. 2018 Nov 29;379(22):2108-2121.
2. Cortes J, Cescon DW, Rugo HS et al.: KEYNOTE-355 Investigators. Pembrolizumab plus chemotherapy versus placebo plus chemotherapy for previously untreated locally recurrent inoperable or metastatic triple-negative breast cancer (KEYNOTE-355): a randomised, placebo-controlled, double-blind, phase 3 clinical trial. *Lancet*. 2020 Dec 5;396(10265):1817-1828.

PARP-Inhibitors

1. Robson M, Im SA, Senkus E, et al. Olaparib for Metastatic Breast Cancer in Patients with a Germline BRCA Mutation. *N Engl J Med*. 2017;377(6):523-533.
2. Litton JK, Rugo HS, Ettl J, et al. Talazoparib in Patients with Advanced Breast Cancer and a Germline BRCA Mutation. *N Engl J Med*. 2018;379(8):753-763.

Bone modifying drugs

1. Aktas B, Kasimir-Bauer S, Lehmann N, et al.: Validity of bone marker measurements for monitoring response to bisphosphonate therapy with zoledronic acid in metastatic breast cancer. *Oncol Rep.* 2013;30(1):441–447.
2. Loftus LS, Edwards-Bennett S, Sokol GH. Systemic therapy for bone metastases. *Cancer Control.* 2012;19(2):145–153.
3. Coleman R, Gnant M, Morgan G, Clezardin P. Effects of bone-targeted agents on cancer progression and mortality. *J Natl Cancer Inst.* 2012;104(14):1059–1067.

CTC monitoring (any therapy)

1. Bidard FC, Peeters DJ, Fehm T, et al. Clinical validity of circulating tumour cells in patients with metastatic breast cancer: a pooled analysis of individual patient data. *Lancet Oncol.* 2014;15:406-14.
2. Smerage JB, Barlow WE, Hortobagyi GN, et al. Circulating tumor cells and response to chemotherapy in metastatic breast cancer: SWOG S0500. *J Clin Oncol.* 2014;32(31):3483-9.

Mutation Diagnostics* in mBC: „Precision Medicine“ for Targeted Therapies

Altered genes	Therapeutic relevance	Gene region	Material	Oxford		
				LOE	GR	AGO
BRCA1, BRCA2	Olaparib, Talazoparib Olaparib	All exons	Germline: Blood cells Somatic: Tissue	1b 2b	A B	++ +/-
PALB2	Olaparib		Germline: Blood cells	2b	B	+
PIK3CA	Alpelisib	Exons 7, 9 and 20	Primary tumor, metastases, plasma	1b	A	++
HER2-mutation (independent of HER2-status)	Neratinib, lapatinib	Kinase- and extracellular domains; S310, L755, V777, Y772_A775dup	Primary tumor, metastases, plasma particul. lobular BC	4	C	+/-
ESR1	Resistance against AI Response oral SERDs	Exons 4, 7 and 8	Metastases, plasma Metastases, plasma	2b 1b	B B	+/- +/-
NTRK gene fusion	Larotrectinib, entrectinib	Fusion- and splice variants	Tumor tissue, particul. secretory breast cancer	2a	B	+
MSI * Ideally panel diagnostics	Pembrolizumab	Microsatellite-instability	Tissue	2a	B	+

BRCA 1/2:

1. Robson M, Im SA, Senkus E et al. (2017) Olaparib for Metastatic Breast Cancer in Patients with a Germline BRCA Mutation. N Engl J Med. 377:523-533. doi: 10.1056/NEJMoa1706450
2. Kaufman B, Shapira-Frommer R, Schmutzler RK et al. (2015) Olaparib monotherapy in patients with advanced cancer and a germline BRCA1/2 mutation. J Clin Oncol. 2015 Jan 20;33(3):244-50. doi: 10.1200/JCO.2014.56.2728.
3. Tung NM, Robson ME, Ventz S, et al. TBCRC 048: Phase II Study of Olaparib for Metastatic Breast Cancer and Mutations in Homologous Recombination-Related Genes. J Clin Oncol. 2020 Dec 20;38(36):4274-4282.

PIK3CA:

1. André F, Ciruelos E, Rubovszky G, Campone M et al. (2019) Alpelisib for PIK3CA-Mutated, Hormone Receptor-Positive Advanced Breast Cancer. N Engl J Med. 380:1929-1940. doi: 10.1056/NEJMoa1813904B
2. Baselga J, Im SA, Iwata H, et al. Buparlisib plus fulvestrant versus placebo plus fulvestrant in postmenopausal, hormone receptor-positive, HER2-negative, advanced breast cancer (BELLE-2): a randomised, double-blind, placebo-controlled, phase 3 trial. Lancet Oncol. 2017 Jul;18(7):904-916.
3. Campone M, Im SA, Iwata H, et al. Buparlisib plus fulvestrant versus placebo plus fulvestrant for postmenopausal, hormone receptor-positive, human epidermal growth factor receptor 2-negative, advanced breast cancer: Overall survival results from BELLE-2. Eur J Cancer. 2018 Nov;103:147-154.

4. Dickler MN, Saura C, Richards DA, et al. Phase II Study of Taselisib (GDC-0032) in Combination with Fulvestrant in Patients with HER2-Negative, Hormone Receptor-Positive Advanced Breast Cancer. *Clin Cancer Res*. 2018 Sep 15;24(18):4380-4387.

HER2-Mutation:

1. Hyman DM, Piha-Paul SA, Won H, et al. (2018) HER kinase inhibition in patients with HER2- and HER3-mutant cancers. *Nature* 554:189-194. doi: 10.1038/nature25475.
2. Ma CX, Bose R, Gao F, et al. (2017) Neratinib Efficacy and Circulating Tumor DNA Detection of HER2 Mutations in HER2 Nonamplified Metastatic Breast Cancer. *Clin Cancer Res*. 23:5687-5695. doi: 10.1158/1078-0432.CCR-17-0900.
3. Hanker AB, Brewer MR, Sheehan JH, et al. An Acquired HER2(T798I) Gatekeeper Mutation Induces Resistance to Neratinib in a Patient with HER2 Mutant-Driven Breast Cancer. *Cancer Discov* 2017;7: 575-85.
4. Xu X, De Angelis C, Burke KA, et al. HER2 Reactivation through Acquisition of the HER2 L755S Mutation as a Mechanism of Acquired Resistance to HER2-targeted Therapy in HER2(+) Breast Cancer. *Clin Cancer Res* 2017;23: 5123-34.

ESR1:

1. Bidard FC, Kaklamani VG, Neven P, et al. (2022) Elacestrant (oral selective estrogen receptor degrader) Versus Standard Endocrine Therapy for Estrogen Receptor-Positive, Human Epidermal Growth Factor Receptor 2-Negative Advanced Breast Cancer: Results From the Randomized Phase III EMERALD Trial. *J Clin Oncol*. 2022 Oct 1;40(28):3246-3256. doi: 10.1200/JCO.22.00338. Epub 2022 May 18. PMID: 35584336; PMCID: PMC9553388.
2. Dustin D, Gu G, Fuqua SAW (2019) ESR1 mutations in breast cancer. *Cancer* 125:3714-3728 doi: 10.1002/cncr.32345.
3. Fribbens C, Garcia Murillas I, Beaney M et al. (2018) Tracking evolution of aromatase inhibitor resistance with circulating tumour DNA analysis in metastatic breast cancer. *Ann Oncol*.29:145-153. doi: 10.1093/annonc/mdx483
4. Fribbens C, O'Leary B, Kilburn L et al. (2016) Plasma ESR1 Mutations and the Treatment of Estrogen Receptor-Positive Advanced Breast Cancer. *J Clin Oncol*. 34:2961-8. doi: 10.1200/JCO.2016.67.3061

NTRK:


1. Cocco E, Scaltriti M, Drilon A (2018) NTRK fusion-positive cancers and TRK inhibitor therapy. *Nat Rev Clin Oncol*. 15(:731-747. doi: 10.1038/s41571-018-0113-0.

MSI:

FDA approval across tumor entities (23.5.17): see full prescribing information for pembrolizumab

PALB2:

1. Antoniou AC, Casadei S, Heikkinen T et al. Breast Cancer Risk in families with mutations in PALB2. N Engl J Med 2014;371(6):497-506.
2. Breast Cancer Association Consortium, Dolling L, Carvahlo S, Allen J et al. Breast Cancer Risk Genes - Association Analysis in more than 113,000 women. N Engl J Med 2021;384(5):428-439.
3. Couch FJ, Shimelis H, Hu C et al. Associations between cancer predisposition

 Decision guidance prospectively evaluated biomarkers (LOE1a) and therapy options (mBC)		
<small>© AGO e. V. in der DGGG e.V. sowie in der DKG e.V. Guidelines Breast Version 2023.1E www.ago-online.de FORSCHEN LEBEN HEILEN</small>		
Biomarker / Signature-therapy option	Subtyp / Population	Therapy option
PDL-L1 $\geq 1\%$	TNBC	First line Atezolizumab + nab Paclitaxel
CPS > 10	TNBC	First line Pembro + chemotherapy
PIK3CA mutation	HR+ / HER2-	Fulvestrant + Alplisib after failure offirst line ET
BRCA1/2 mutation (OlympiAD, EMBRACA)	HER2 –	Olaparib, Talazoparib

Head to head comparisons

1. Varga Z, Sinn P, Seidman AD. Summary of head-to-head comparisons of patient risk classifications by the 21-gene Recurrence Score® (RS) assay and other genomic assays for early breast cancer. Int J Cancer. 2019 Aug 15;145(4):882-893.
2. Berchtold E, Vetter M, Gündert M et al. Comparison of six breast cancer classifiers using qPCR. Bioinformatics. 2019 Sep 15;35(18):3412-3420.
3. Sestak I, Buus R, Cuzick J, et al. Comparison of the Performance of 6 Prognostic Signatures for Estrogen Receptor-Positive Breast Cancer: A Secondary Analysis of a Randomized Clinical Trial. JAMA Oncol. 2018 Apr 1;4(4):545-553.

Endopredict

1. Blank PR, Filipits M, Dubsky P, et al.: Cost-effectiveness analysis of prognostic gene expression signature-based stratification of early breast cancer patients. Pharmacoeconomics. 2015 Feb;33(2):179-90.
2. Buus R, Sestak I, Kronenwett R, et al.: Comparison of EndoPredict and EPclin with Oncotype DX Recurrence Score for Prediction of Risk of Distant Recurrence after endocrine therapy J Natl Cancer Inst. 2016 Jul 10;108(11).
3. Denkert, C., Kronenwett, R., Schlake, W. et al. 2012. Decentral gene expression analysis for ER+/Her2- breast cancer: results of a

proficiency testing program for the EndoPredict assay. *Virchows Arch.* 460, 251–259. doi:10.1007/s00428-012-1204-4.

4. Dubsky, P., Filipits, M., Jakesz, R. et al. 2012. EndoPredict improves the prognostic classification derived from common clinical guidelines in ER-positive, HER2-negative early breast cancer. *Ann Oncol.* 2013 Mar;24(3):640-7.
5. Dubsky P, Brase JC, Jakesz R et al. Austrian Breast and Colorectal Cancer Study Group (ABCSG). T al.: The EndoPredict score provides prognostic information on late distant metastases in ER+/HER2- breast cancer patients. *Br J Cancer.* 2013 Dec 10;109(12):2959-64
6. Filipits, M., Rudas, M., Jakesz, R. et al. 2011. A new molecular predictor of distant recurrence in ER-positive, HER2-negative breast cancer adds independent information to conventional clinical risk factors. *Clin. Cancer Res.* 17, 6012–6020.
7. Kronenwett, R., Bohmann, K., Prinzler, J. et al. 2012. Decentral gene expression analysis: analytical validation of the Endopredict genomic multianalyte breast cancer prognosis test. *BMC Cancer* 12, 456.
8. Martin M, Brase JC, Calvo L, et al. Clinical validation of the EndoPredict test in node-positive, chemotherapy-treated ER+/HER2- breast cancer patients: results from the GEICAM 9906 trial. *Breast Cancer Res* 2014; 16(2): R38.
9. Sinn P, Aulmann S, Wirtz R, et al. Multigene Assays for Classification, Prognosis, and Prediction in Breast Cancer: a Critical Review on the Background and Clinical Utility. *A. Geburtshilfe Frauenheilkd.* 2013 Sep;73(9):932-940

MammaPrint

1. Buyse, M., Loi, S., van't Veer, L., et al. 2006. Validation and clinical utility of a 70-gene prognostic signature for women with node-negative breast cancer. *J. Natl. Cancer Inst.* 98, 1183–1192.
2. Drukker CA, Elias SG, Nijenhuis MV, et al. Gene expression profiling to predict the risk of locoregional recurrence in breast cancer: a pooled analysis. *Breast Cancer Res Treat.* 2014 Dec;148(3):599-613.
3. Exner R, Bago-Horvath Z, Bartsch R et al. The multigene signature MammaPrint impacts on multidisciplinary team decisions in ER+, HER2- early breast cancer. *Br J Cancer.* 2014 Aug 26;111(5):837-42.
4. Jonsdottir K, Assmus J, Slewa A, et al. Prognostic value of gene signatures and proliferation in lymph-node-negative breast cancer. *PLoS One.* 2014 Mar 5;9(3):e90642.
5. Mook, S., Schmidt, M.K., Weigelt, B., et al. 2010. The 70-gene prognosis signature predicts early metastasis in breast cancer patients between 55 and 70 years of age. *Ann. Oncol.* 21, 717–722.

6. Mook, S., Schmidt, M.K., Rutgers, E.J., et al. 2009a. Calibration and discriminatory accuracy of prognosis calculation for breast cancer with the online Adjuvant! program: a hospital-based retrospective cohort study. *Lancet Oncol.* 10, 1070–1076.
7. Mook, S., Schmidt, M.K., Viale, G. et al. 2009b. The 70-gene prognosis-signature predicts disease outcome in breast cancer patients with 1-3 positive lymph nodes in an independent validation study. *Breast Cancer Res. Treat.* 116, 295–302.
8. van de Vijver, M.J., He, Y.D., van't Veer, L.J., et al. 2002. A gene-expression signature as a predictor of survival in breast cancer. *N. Engl. J. Med.* 347, 1999–2009. doi:10.1056/NEJMoa021967.
9. van Veer, L.J. 't, Dai, H., van de Vijver, M.J., et al. 2002. Gene expression profiling predicts clinical outcome of breast cancer. *Nature* 415, 530–536. doi:10.1038/415530a.
10. Cardoso F, van't Veer LJ, Bogaerts J, et al. 70-Gene Signature as an Aid to Treatment Decisions in Early-Stage Breast Cancer. *N Engl J Med.* 2016 Aug 25;375(8):717-29.
11. Cardoso F, van 't Veer L, Poncet C, et al. MINDACT: Long-term results of the large prospective trial testing the 70-gene signature MammaPrint as guidance for adjuvant chemotherapy in breast cancer patients. *ASCO 2020*, #506

Oncotype DX

1. Albain, K.S., Barlow, W.E., Shak, S. et al. 2010. Prognostic and predictive value of the 21-gene recurrence score assay in postmenopausal women with node-positive, oestrogen-receptor-positive breast cancer on chemotherapy: a retrospective analysis of a randomised trial. *Lancet Oncol.* 11, 55–65. Cronin, M., Sangli, C., Liu, M.-L., et al. 2007. Analytical validation of the Oncotype DX genomic diagnostic test for recurrence prognosis and therapeutic response prediction in node-negative, estrogen receptor-positive breast cancer. *Clin. Chem.* 53, 1084–1091.
2. Dowsett, M., Cuzick, J., Wale, C. et al. 2010. Prediction of risk of distant recurrence using the 21-gene recurrence score in node-negative and node-positive postmenopausal patients with breast cancer treated with anastrozole or tamoxifen: a TransATAC study. *J. Clin. Oncol.* 28, 1829–1834.
3. Gluz O, Nitz UA, Christgen M, et al.: West German Study Group Phase III PlanB Trial: First Prospective Outcome Data for the 21-Gene Recurrence Score Assay and Concordance of Prognostic Markers by Central and Local Pathology Assessment. *J Clin Oncol.* 2016 Jul 10;34(20):2341-9.

4. Khan SS, Karn T, Symmans WF, et al. Genomic predictor of residual risk of recurrence after adjuvant chemotherapy and endocrine therapy in high risk estrogen receptor-positive breast cancers. *Breast Cancer Res Treat.* 2015 Feb 5.
5. Mamounas, E.P., Tang, G., Fisher, B., et al. 2010. Association between the 21-gene recurrence score assay and risk of locoregional recurrence in node-negative, estrogen receptor-positive breast cancer: results from NSABP B-14 and NSABP B-20. *J. Clin. Oncol.* 28, 1677–1683.
6. Paik, S., Shak, S., Tang, G. et al. 2004. A multigene assay to predict recurrence of tamoxifen-treated, node-negative breast cancer. *N. Engl. J. Med.* 351, 2817–2826. doi:10.1056/NEJMoa041588.
7. Paik, S., Tang, G., Shak, S., et al. 2006. Gene expression and benefit of chemotherapy in women with node-negative, estrogen receptor-positive breast cancer. *J. Clin. Oncol.* 24, 3726–3734.
8. Sparano JA, Gray RJ, Makower DF, et al: Prospective Validation of a 21-Gene Expression Assay in Breast Cancer. *N Engl J Med* 2015, 373(21):2005-2014.
9. Tang, G., Cuzick, J., Costantino, J.P., et al. 2011. Risk of recurrence and chemotherapy benefit for patients with node-negative, estrogen receptor-positive breast cancer: recurrence score alone and integrated with pathologic and clinical factors. *J. Clin. Oncol.* 29, 4365–4372.
10. Zhao X, Rødland EA, Sørli T, et al. Systematic assessment of prognostic gene signatures for breast cancer shows distinct influence of time and ER status. *BMC Cancer.* 2014 Mar 19;14:211.
11. Geyer CE Jr, Tang G, Mamounas EP, et al. 21-Gene assay as predictor of chemotherapy benefit in HER2-negative breast cancer. *NPIJ Breast Cancer.* 2018 Nov 14;4:37.
12. Penault-Llorca F, Filleron T, Asselain B, et al. The 21-gene Recurrence Score® assay predicts distant recurrence in lymph node-positive, hormone receptor-positive, breast cancer patients treated with adjuvant sequential epirubicin- and docetaxel-based or epirubicin-based chemotherapy (PACS-01 trial). *BMC Cancer.* 2018 May 4;18(1):526.
13. Sparano JA, Gray R, Oktay MH, et al. A metastasis biomarker (MetaSite Breast™ Score) is associated with distant recurrence in hormone receptor-positive, HER2-negative early-stage breast cancer. *NPIJ Breast Cancer.* 2017 Nov 8;3:42.
14. Sparano JA, Gray RJ, Makower DF, et al. Adjuvant Chemotherapy Guided by a 21-Gene Expression Assay in Breast Cancer. *N Engl J Med.* 2018 Jul 12;379(2):111-121.
15. Nitz U, Gluz O, Christgen M, et al. Reducing chemotherapy use in clinically high-risk, genomically low-risk pN0 and pN1 early breast

cancer patients: five-year data from the prospective, randomised phase 3 West German Study Group (WSG) PlanB trial. *Breast Cancer Res Treat.* 2017 Oct;165(3):573-583.

16. Harbeck N, Gluz O, Kuemmel S, et al. Endocrine therapy alone in patients with intermediate or high-risk luminal early breast cancer (0-3 lymph nodes), Recurrence Score <26 and Ki67 response after preoperative endocrine therapy: Primary outcome results from the WSG-ADAPT HR+/HER2- trial. *SABCS 2020*, GS4-04.
17. Kalinsky K, Barlow WE, Meric-Bernstam F, et al. First results from a phase III randomized clinical trial of standard adjuvant endocrine therapy (ET) +/- chemotherapy (CT) in patients (pts) with 1-3 positive nodes, hormone receptor-positive (HR+) and HER2-negative (HER2-) breast cancer (BC) with recurrence score (RS) < 25: SWOG S1007 (RxPonder). *SABCS 2020*, GS3-00.

Prosigna (ROR / PAM50)

1. Chia, S.K., Bramwell, V.H., Tu, Det al. 2012. A 50-gene intrinsic subtype classifier for prognosis and prediction of benefit from adjuvant tamoxifen. *Clin. Cancer Res.* 18, 4465–4472.
2. Gnant M, Filipits M, Greil R, et al. Austrian Breast and Colorectal Cancer Study Group. Predicting distant recurrence in receptor-positive breast cancer patients with limited clinicopathological risk: using the PAM50 Risk of Recurrence score in 1478 postmenopausal patients of the ABCSG-8 trial treated with adjuvant endocrine therapy alone. *Ann Oncol.* 2014 Feb;25(2):339-45
3. Liu S, Chapman JA, Burnell MJ, et al. Prognostic and predictive investigation of PAM50 intrinsic subtypes in the NCIC CTG MA.21 phase III chemotherapy trial. *Breast Cancer Res Treat.* 2015 Jan;149(2):439-48.
4. Nielsen, T.O., Parker, J.S., Leung, S., et al. 2010. A comparison of PAM50 intrinsic subtyping with immunohistochemistry and clinical prognostic factors in tamoxifen-treated estrogen receptor-positive breast cancer. *Clin. Cancer Res.* 16, 5222–5232.
5. Parker, J.S., Mullins, M., Cheang, M.C.U., et al. 2009. Supervised risk predictor of breast cancer based on intrinsic subtypes. *J. Clin. Oncol.* 27, 1160–1167.
6. Prat, A., Cheang, M.C.U., Martín, M., et al. 2012b. Prognostic Significance of Progesterone Receptor-Positive Tumor Cells Within Immunohistochemically Defined Luminal A Breast Cancer. *J. Clin. Oncol.*
7. Prat, A., Parker, J.S., Fan, C., et al. 2012a. Concordance among gene expression-based predictors for ER-positive breast cancer treated with adjuvant tamoxifen. *Ann. Oncol.* 23, 2866–2873.

8. Perou, C.M., Sørli, T., Eisen, M.B. et al. 2000. Molecular portraits of human breast tumours. *Nature* 406, 747–752.
9. Pogue-Geile KL, Song N, Jeong JH, et al. Intrinsic Subtypes, PIK3CA Mutation, and the Degree of Benefit From Adjuvant Trastuzumab in the NSABP B-31 Trial. *J Clin Oncol.* 2015 Jan 5.
10. Sestak I, Cuzick J, Dowsett M. et al.. Prediction of Late Distant Recurrence After 5 Years of Endocrine Treatment: A Combined Analysis of Patients From the Austrian Breast and Colorectal Cancer Study Group 8 and Arimidex, Tamoxifen Alone or in Combination Randomized Trials Using the PAM50 Risk of Recurrence Score. *J Clin Oncol.* 2014 Oct 20. pii: JCO.2014.55.6894
11. Jensen MB, Lænkholm AV, Nielsen TO, et al. The Prosigna gene expression assay and responsiveness to adjuvant cyclophosphamide-based chemotherapy in premenopausal high-risk patients with breast cancer. *Breast Cancer Res.* 2018 Jul 27;20(1):79.
12. Lænkholm AV, Jensen MB, Eriksen JO, et al. PAM50 Risk of Recurrence Score Predicts 10-Year Distant Recurrence in a Comprehensive Danish Cohort of Postmenopausal Women Allocated to 5 Years of Endocrine Therapy for Hormone Receptor-Positive Early Breast Cancer. *J Clin Oncol.* 2018 Mar 10;36(8):735-740.
13. Ohnstad HO, Borgen E, Falk RS, et al. Prognostic value of PAM50 and risk of recurrence score in patients with early-stage breast cancer with long-term follow-up. *Breast Cancer Res.* 2017 Nov 14;19(1):120.

Breast Cancer Index

1. Bartlett JMS, Sgroi DC, Treuner K et al. Breast Cancer Index and prediction of benefit from extended endocrine therapy in breast cancer patients treated in the Adjuvant Tamoxifen-To Offer More? (aTTom) trial. *Ann Oncol.* 2019 Nov 1;30(11):1776-1783.
2. Jerevall PL, Brommesson S, Strand C, et al. Exploring the two-gene ratio in breast cancer--independent roles for HOXB13 and IL17BR in prediction of clinical outcome. *Breast Cancer Res Treat.* 2008;107(2):225–234.
3. Jansen MP, Sieuwerts AM, Look MP, et al. HOXB13-to-IL17BR expression ratio is related with tumor aggressiveness and response to tamoxifen of recurrent breast cancer: a retrospective study. *J Clin Oncol.* 2007;25(6):662–668.
4. Ma XJ, Hilsenbeck SG, Wang W, et al. The HOXB13:IL17BR expression index is a prognostic factor in early-stage breast cancer. *J Clin Oncol.* 2006;24(28):4611–4619.

PARPi

1. Robson M, Im SA, Senkus E et al. Olaparib for metastatic breast cancer in patients with germline BRCA Mutation. N Engl J Med 2017;377(6):523-533.
2. Litton JK, Rugo HS, Ettl J et al. Talazoparib in patients with advanced breast cancer and a germline BRCA mutation N Engl J Med 2018;379(8):753-763.

Therapy-Relevant Mutational Analysis for „Actionable“ Genomic Alterations in BC


Diagnostic Tool*	Outcome	Oxford			
		LoE	GR	AGO	
Evidence from studies with other cancer patients („tumor-agnostic testing“)					
▪ Companion Diagnostics for therapies of other tumor entities (e.g. BRAF, FGFR1, ...)	Efficacy of diverse therapies	4	D	+/-**	
▪ Large Panel Gene Analysis (e.g. FoundationOne, GPS Cancer, NeoSelect, Molecular Health Guide, local „hand-selected„ panels)	Efficacy of diverse therapies, prognosis	3a	C	+/-**	
▪ Next Generation Sequencing (NGS) (recommended only in Tier 1 + 2)	Efficacy of evaluated drugs	1b	B	+/-**	

* Assessment method for somatic mutations (tumor tissue, cf-DNA) is not taken into consideration for LoE

** Participation in clinical trials or structured registries recommended

NGS in breast cancer:

1. Mosele F, Remon J, Mateo J, et al. Recommendations for the use of next-generation sequencing (NGS) for patients with metastatic cancers: a report from the ESMO Precision Medicine Working Group. Ann Oncol. 2020 Nov;31(11):1491-1505.
2. André F, Bachelot T, Commo F, et al. Comparative genomic hybridisation array and DNA sequencing to direct treatment of metastatic breast cancer: a multicentre, prospective trial (SAFIR01/UNICANCER). Lancet Oncol. 2014 Mar;15(3):267-74.
3. van Geelen CT, Savas P, Teo ZL, et al. Clinical implications of prospective genomic profiling of metastatic breast cancer patients. Breast Cancer Res. 2020 Aug 18;22(1):91.
4. Hempel D, Ebner F, Garg A, et al. Real world data analysis of next generation sequencing and protein expression in metastatic breast cancer patients. Sci Rep. 2020 Jun 26;10(1):10459.
5. Sultova E, Westphalen CB, Jung A, et al. NGS-guided precision oncology in metastatic breast and gynecological cancer: first experiences at the CCC Munich LMU. Gynecol Obstet. 2020 Dec 4. doi: 10.1007/s00404-020-05881-z.



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2023.1E

www.ago-online.de

FORSCHEN
LEBEN
HEILEN

Joint Consensus Recommendations of AMP, ACMG, ASCO and CAP for Reporting Genetic Variants in Cancer

Tier	LoE		Explanation
Tier 1	A.1	Biomarkers that predict response or resistance to FDA-approved therapies for a specific type of cancer	Variants of strong clinical significance
	A.2	Biomarkers included in professional guidelines that predict response to therapies for a specific type of tumor	
	B	Biomarkers that predict response or resistance to therapies for a specific type of tumor based on well-powered studies with consensus from experts in the field	
Tier 2	C.1	Biomarkers that predict response or resistance to therapies approved by the FDA or professional societies for a different type of tumor	Variants of potential clinical significance
	C.2	Biomarkers that serve as inclusion criteria for clinical trials	
	D	Biomarkers that show plausible therapeutic significance based on preclinical studies	
Tier 3		Not observed at a significant allele frequency in the general or specific subpopulation databases, or pan-cancer or tumor-specific variant databasis. No convincing published evidence or cancer association	Variants of unknown clinical significance
Tier 4		Observed at significant allele frequency in the general or specific subpopulation Databases. No existing published evidence of cancer association	Benign or likely benign variants

Treatment Recommendations for genetic variants

1. Li MM, Datto M, Duncavage EJ, et al. Standards and guidelines for the interpretation and reporting of sequence variants in cancer: a joint consensus recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists. J Mol Diagn 2017;19:4–23.
2. Leichsenring J, Horak P, Kreuzfeldt S et al. Variant classification in precision oncology. Int J. Cancer 2019;145:2996–3010
3. Condorelli R, Mosele F, Verret B et al. Genomic alterations in breast cancer: level of evidence for actionability according to ESMO Scale for Clinical Actionability of molecular Targets (ECAT). Ann Oncol 2019;30(3):365-373.