



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.


Guidelines Breast
Version 2025.1E

FORSCHEN
LEHREN
HEILEN

Diagnosis and Treatment of Patients with early and advanced Breast Cancer

Breast Cancer Follow-Up

Breast Cancer Follow-Up




- **Versions 2002-2024:**

**Bauerfeind / Bischoff / Blohmer / Böhme / Costa / Diel / Friedrich /
Gerber / Gluz / Hanf / Heinrich / Huober / Janni / Kaufmann /
Kolberg-Liedtke / Kümmel / Lüftner / Lux / Maass / Möbus / Müller-
Schimpfle/ Mundhenke / Oberhoff / Rody / Scharl / Schmidt / Solbach
/ Solomayer / Stickeler / Thomssen / Wöckel**

- **Version 2025:**

Witzel / Wöckel

1. Aktualisierung der Therapieempfehlungen nach Durchsicht der ASCO, NCCN und ACS Guidelines*, sowie der S3 Leitlinie
2. Durchgeführte „Medline“ und „PubMed“-Suche nach „Surveillance Breast Cancer“ und „Follow up primary breast cancer“ (2020/01-2025/01)



Breast Cancer Follow-Up Objectives

	Oxford		
	LoE	GR	AGO
Early detection of curable events			
▪ In-breast recurrence	1a	B	++
▪ Loco-regional recurrence*	1a	B	++
Early detection of contralateral cancers	1a	B	++
Early detection of metastasis			
▪ Early detection of symptomatic metastases	3b	C	+
▪ Early detection of asymptomatic metastases	1a	A	-

* loco-regional recurrence is associated with a higher risk of mortality in node-positive, PR-negative, younger patients and in patients with a short time between primary diagnosis and recurrence


1. De Bock GH, Bonnema J, van Der Hage J et al., Effectiveness of Routine Visits and Routine Tests in Detecting Isolated Locoregional Recurrences After Treatment for Early-Stage Invasive Breast Cancer: A Meta-Analysis and Systematic Review. J Clin Oncol 2004; 22 (19): 4010-4018.
2. Margenthaler JA, Allan E, Cheng L, et al. Surveillance of Patients With Breast Cancer After Curative-Intent Primary Treatment: Current Practice Patterns. Journal of Oncology Practice 2012; 8(2): 79 – 83.
3. Parmeshwar R, Margenthaler JA, Allam E, et al. Patient Surveillance After Initial Breast cancer Therapy Variation by Physician Specialty. Am J Surg 2013; 206(2): 218-222.
4. Jochelson M, Hayes DF, Ganz PA. Surveillance and Monitoring in Breast Cancer Survivors: Maximizing Benefit and Minimizing Harm. ASCO Educational Book 2013 e13 – e18.
5. Khatcheressian JL, Hurley P, Bantug E, et al. Breast Cancer Follow-up and Management After Primary Treatment: American Society of Clinical Oncology Clinical Practice Guideline Update . J Clin Oncol. 2013 March 1; 31(7):961-965.
6. Runowicz CD, Leach CR, Henry NL, et al. American Cancer Society/American Society of Clinical Oncology Breast Cancer Survivorship Care Guideline. Journal of Clinical Oncology 2016;34:611-35.
7. Cardoso F, Kyriakides S, Ohno S, et al. Early breast cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. Annals of Oncology 2019;30:1194-220.
8. Lee et al. Cancer Res Treat. 2015 Oct;47(4):765-73 Reduced mortality according to regular follow up Impact on Survival

of Regular Postoperative Surveillance for Patients with Early Breast Cancer

9. Moschetti I, Cinquini M, Lambertini M et al., Follow-up strategies for women treated for early breast cancer. Cochrane Database Syst Rev. 2016 May 27;(5):1CD001768.
10. NCCN Clinical Practice Guidelines in Oncology, Breast Cancer Version 3.17-10.17;
https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf
11. NCCN Clinical Practice Guidelines in Oncology, Breast Cancer Version 6.2020
https://www.nccn.org/professionals/physician_gls/pdf/breast_blocks.pdf
12. Lin J, Hu H, Shriver CD et al., Survival among Breast Cancer Patients: Comparison of the U.S. Military Health System with the Surveillance, Epidemiology and End Results Program. Clin Breast Cancer. 2021 Dec 1:S1526-8209(21)00348-7.
13. Viuff JH, Greiber IK, Karlsen MA et al., Survival in Women Diagnosed With Breast Cancer During Pregnancy. Clin Breast Cancer. 2021 Dec 1:S1526-8209(21)00346-3. doi: 10.1016/j.clbc.2021.11.009.

Statement: risk factors of mortality after loco-regional recurrence

1. Dent R, Valentini H, Hanna W, et al.. Factors associated with breast cancer mortality after local recurrence. Curr Oncol 2014; 21 (3): e418-25.



Breast Cancer Follow-Up Objectives

	Oxford		
	LoE	GR	AGO
▪ Improve quality of life	2b	B	+
▪ Improve physical performance	2a	B	+
▪ Reduction and / or early detection of therapy-related side effects (such as osteoporosis, cardiac failure, fatigue, endocrinopathy, neurotoxicity, lymphedema, web axillary pain syndrome (abacterial lymphangitis), sexual disorders, cognitive impairment, sterility, and secondary tumors) and start of necessary therapies	2b	B	+
▪ Participation in interventional programs during follow-up for breast cancer survivors in order to maximize therapy adherence, assess life-style interventions, and improve quality of life	3b	B	+

Statement: Outcome measurements

1. Browall M, Forsberg C, Wengström Y. Assessing patient outcomes and cost-effectiveness of nurse-led follow-up for women with breast cancer - have relevant and sensitive evaluation measures been used? J Clin Nurs. 2017 Jul;26(13-14):1770-1786.
2. Cheng KKF, Lim YTE, Koh ZM et al. Home-based multidimensional survivorship programmes for breast cancer survivors. Cochrane Database Syst Rev. 2017 Aug 24;8:CD011152.
3. Dent S, Fergusson D, Aseyev O et al., A Randomized Trial Comparing 3- versus 4-Monthly Cardiac Monitoring in Patients Receiving Trastuzumab-Based Chemotherapy for Early Breast Cancer. Curr Oncol. 2021 Dec 3;28(6):5073-5083.
4. Franzoi MA, Degousée L, Martin et al. (2023) Implementing a PROACTIVE Care Pathway to Empower and Support Survivors of Breast Cancer. JCO Oncol Pract 19(6):353–361. doi:10.1200/OP.23.00016
5. Lewinter C, Nielsen TH, Edfors LR et al., A systematic review and meta-analysis of beta-blockers and renin-angiotensin system inhibitors for preventing left ventricular dysfunction due to anthracyclines or trastuzumab in patients with breast cancer. Eur Heart J. 2021 Dec 24;ehab843. doi: 10.1093/eurheartj/ehab843.
6. Ong WL, Schouwenburg MG, van Bommel ACM et al. A Standard Set of Value-Based Patient-Centered Outcomes for Breast Cancer: The International Consortium for Health Outcomes Measurement (ICHOM) Initiative. JAMA Oncol. 2017 May 1;3(5):677-685.

7. Rassaf T, Totzeck M, Backs J et al. for Committee for Clinical Cardiovascular Medicine of the German Cardiac Society. Onco-Cardiology: Consensus Paper of the German Cardiac Society, the German Society for Pediatric Cardiology and Congenital Heart Defects and the German Society for Hematology and Medical Oncology. Clin Res Cardiol. 2020 Oct;109(10):1197-1222.

Statement: Obesity, physical activity and quality of life

1. Baumann FT, Bieck O, Oberste M et al.. Sustainable impact of an individualized exercise program on physical activity level and fatigue syndrome on breast cancer patients in two German rehabilitation centers. Support Care Cancer. 2017 Apr;25(4):1047-1054.
2. Bower JE, Garet D, Sternlieb B et al. Yoga for persistent fatigue in breast cancer survivors: A randomized controlled trial. Cancer 2012; 118(15): 3766-75.
3. Cramp F, Byron-Daniel J. Exercise for the management of cancer-related fatigue in adults. Cochrane Database Syst Rev. 2012 Nov 14;11:CD006145.
4. Curigliano G, Lenihan D, Fradley M, et al. Management of cardiac disease in cancer patients throughout oncological treatment: ESMO consensus recommendations. Annals of Oncology 2020;31:171-90.
5. Fabi A, Bhargava R, Fatigoni S, et al. Cancer-related fatigue: ESMO Clinical Practice Guidelines for diagnosis and treatment. Annals of Oncology 2020;31:713-23.
6. Lisevick A, Cartmel B, Harrigan M, Li F, et al. Effect of the Lifestyle, Exercise, and Nutrition (LEAN) Study on Long-Term Weight Loss Maintenance in Women with Breast Cancer. Nutrients. 2021 Sep 18;13(9):3265.
7. Nagy AC, GulAcsi-BARDos P, CserEp Z et al.Late cardiac effect of anthracycline therapy in physically active breast cancer survivors - a prospective study. Neoplasma. 2017;64(1):92-100.
8. Vaskuil DW, van Nes JG, Junngeburt JM et al. Maintenance of physical activity and body weight in relation to subsequent quality of life in postmenopausal breast cancer patients. Annals of Oncology 2010; 21: 2094–2101.
9. Yang H, Brand JS, Fang F et al. Time-dependent risk of depression, anxiety, and stress-related disorders in patients with invasive and in situ breast cancer. Int J Cancer. 2017 Feb 15;140(4):841-852.

Statement: Obesity and breast cancer prognosis

1. Ewertz M, Jensen MB, Gunnarsdóttir KÁ et al.. Effect of obesity on prognosis after early-stage breast cancer. J Clin Oncol 2011; 29(1): 25-31.

2. Cespedes Feliciano EM, Kroenke CH, Bradshaw PT et al..Postdiagnosis Weight Change and Survival Following a Diagnosis of Early-Stage Breast Cancer. *Cancer Epidemiol Biomarkers Prev.* 2017 Jan;26(1):44-50.

Statement: Lymphedema

1. Soran A, Ozmen T, McGuire KP et al..The importance of detection of subclinical lymphedema for the prevention of breast cancer-related clinical lymphedema after axillary lymph node dissection; a prospective observational study.*Lymphat Res Biol* 2014;12(4): 289-9.
2. Basta MN, Wu LC, Kanchwala SK et al.. Reliable prediction of postmastectomy lymphedema: the Risk Assessment Tool Evaluating Lymphedema. *Am J Surg.* 2017 Jun;213(6):1125-1133.
3. Ferguson CM, Swaroop MN, Horick N, et al. Impact of Ipsilateral Blood Draws, Injections, Blood Pressure Measurements, and Air Travel on the Risk of Lymphedema for Patients Treated for Breast Cancer. *Journal of Clinical Oncology* 2016;34:691-8.
4. McNeely ML, Dolgoy ND, Rafn BS et. al. Nighttime compression supports improved self-management of breast cancer-related lymphedema: A multicenter randomized controlled trial. *Cancer.* 2021 Oct 6. doi: 10.1002/cncr.33943.
5. Donahue PMC, MacKenzie A, Filipovic A, Koelmeyer L (2023) Advances in the prevention and treatment of breast cancer-related lymphedema. *Breast Cancer Res Treat* 200(1):1–14. doi:10.1007/s10549-023-06947-7
6. Jeffers EJ, Wagner JL, Korentager SS et al. (2023) Breast Cancer-Related Lymphedema (BCRL) and Bioimpedance Spectroscopy: Long-Term Follow-Up, Surveillance Recommendations, and Multidisciplinary Risk Factors. *Ann Surg Oncol* 30(10):6258–6265. doi:10.1245/s10434-023-13956-9

Statement: Neurotoxicity:

1. Jordan B, Margulies A, Cardoso F et al. Systemic anticancer therapy-induced peripheral and central neurotoxicity: ESMO-EONS-EANO Clinical Practice guidelines for diagnosis, prevention, treatment and follow-up. *Ann Oncology* 2020 Oct; 31(10):1306-1319. doi 10.1016/j.annonc.2020.07.003

Statement: web axillary pain syndrome (Morbus Mondor):

1. Agostini F, Attanasi C, Bernetti A et al., Web Axillary Pain Syndrome-Literature Evidence and Novel Rehabilitative Suggestions: A Narrative Review. *Int J Environ Res Public Health.* 2021 Oct 2;18(19):10383.

Statement: sexual disorders and cognitive impairment:


1. Runowcz CD, Leach CR, Henry L et al.. American Cancer Society/American Society of Clinical Oncology breast cancer survivorship care guideline. CA Cancer J Clin 2016; 66: 43-73.
2. Janelins MC, Heckler CE, Peppone LJ et al.. Cognitive Complaints in Survivors of Breast Cancer After Chemotherapy Compared With Age-Matched Controls: An Analysis From a Nationwide, Multicenter, Prospective Longitudinal Study. J Clin Oncol. 2017 Feb 10;35(5):506-514.
3. Bernstein LJ, McCreath GA, Komeylian Z et al..Cognitive impairment in breast cancer survivors treated with chemotherapy depends on control group type and cognitive domains assessed: A multilevel meta-analysis. Neurosci Biobehav Rev. 2017 Dec;83:417-428.

Statement kognitive Einschränkungen

1. Small BJ, Lange M, Zhai W et al. for Thinking Living with Cancer C. O. G.-Age Studies. Impact of taxane-based chemotherapy among older women with breast cancer on cognition and quality of life: a longitudinal pooled analysis. Breast Cancer Res Treat. 2021 Nov 24. doi: 10.1007/s10549-021-06455-6.

Statement: Secondary tumors:

1. Hoekstra N, Fleury E, Merino Lara TR, et al. Long-term risks of secondary cancer for various whole and partial breast irradiation techniques. Radiother Oncol. 2018 Sep;128(3):428-433



Monitoring after Cardiotoxic Therapy (e.g. Anthracyclines, anti-HER2, Immuno-oncology)

After anthracyclines / Trastuzumab:

- ECG and echocardiography:
 - 6, 12, 24 months and yearly up to 5 years after therapy
 - after 5th year, every 5 years and if patient is symptomatic
- If cardiovascular risk factors:
 - blood pressure at least yearly
 - lipids and HbA1c in serum yearly
- Modify risk factors if possible:
 - nicotine, body weight, bmi
- Education about individual risk profile and lifestyle

Risk factors:

radiotherapy of left breast, nicotine, hypertonus, diabetes mell., dyslipidaemia, adiposity, age > 60, cardiac diseases: reduced ejection fraction, post-myocardial infarction status, ≥ moderate heart defects

1. Curigliano G, Lenihan D, Fradley M, et al. Management of cardiac disease in cancer patients throughout oncological treatment: ESMO consensus recommendations. *Annals of Oncology* 2020;31:171-90.
2. Dent S, Fergusson D, Aseyev O et al. „A Randomized Trial Comparing 3- versus 4-Monthly Cardiac Monitoring in Patients Receiving Trastuzumab-Based Chemotherapy for Early Breast Cancer. *Curr Oncol.* 2021 Dec 3;28(6):5073-5083.
3. Michel L, Rassaf T, Totzeck M. Biomarkers for the detection of apparent and subclinical cancer therapy-related cardiotoxicity. *J Thorac Dis* 2018;10:S4282-S95.
4. Panuccio G, Correale P, d'Apolito M, et al. Immuno-related cardio-vascular adverse events associated with immuno-oncological treatments: an under-estimated threat for cancer patients. *Basic Res Cardiol.* 2024 Sep 3. doi: 10.1007/s00395-024-01077-7.
5. Rassaf T, Totzeck M, Backs J, et al. Onco-Cardiology: Consensus Paper of the German Cardiac Society, the German Society for Pediatric Cardiology and Congenital Heart Defects and the German Society for Hematology and Medical Oncology. *Clin Res Cardiol* 2020;109:1197-222.

Breast Cancer Follow-Up Objectives

	Oxford		
	LoE	GR	AGO
<ul style="list-style-type: none"> Evaluation of current adjuvant therapy <ul style="list-style-type: none"> incl. monitoring of adherence to endocrine therapies Control of menopausal status, e.g. in case of CT-induced amenorrhea (FSH/2 or bleeding history) and addition of GnRH analogs (up to 2 years after CT) if premenopausal status in women < 45 years old, or switch to aromatase inhibitors (if postmenopausal) Pro-active improvement of therapy adherence <ul style="list-style-type: none"> Patient information about efficacy data for 5-10 years endocrine therapy Early therapy of side effects (sports, NSAIDs, vitamin D / calcium) 	2b	B	++
	5	D	++

Evaluation of current adjuvant therapy

- Hershman DL, Kushi LH, Shao T et al. Early Discontinuation and Nonadherence to Adjuvant Hormonal Therapy in a Cohort of 8,769 Early-Stage Breast Cancer Patients. J Clin Oncol 2010; 28: 4120-4128.
- Lueck H-J, Hadji P, Harbeck N et al. 24 Months Follow-Up Results from PACT (Patient's Anastrozole Compliance to Therapy Programme), a Non-Interventional Study Evaluating the Influence of a Standardized Information Service on Compliance in Postmenopausal Women with Early Breast Cancer. SABCS 2011 [P5-17-05].
- Neven P, Markopoulos C, Tanner M et al. The impact of educational materials on compliance and persistence rates with adjuvant aromatase inhibitor treatment: first-year results from the compliance of aromatase inhibitors assessment in daily practice through educational approach (CARIATIDE) study. Breast. 2014 Aug;23(4):393-9.
- Hershman DL, Kushi LH, Hillyer GC et al. Psychosocial factors related to non persistence with adjuvant endocrine therapy among women with breast cancer: the Breast Cancer Quality of Care Study (BQUAL). Breast Cancer Res Treat. 2016 May;157(1):133-43.
- Goss PE, Ingle JN, Pritchard KI et al. Extending Aromatase-Inhibitor Adjuvant Therapy to 10 Years. N Engl J Med. 2016 Jul 21;375(3):209-19.
- Nabieva N, Kellner S, Fehm T et al. Patient and tumor characteristics and their influence on early therapy persistence with letrozole in postmenopausal patients with early breast cancer. Ann Oncol. 2017 Oct 10. doi: 10.1093/annonc/mdx630.


7. Laroche F, Perrot S, Medkour T et al. Quality of life and impact of pain in women treated with aromatase inhibitors for breast cancer. A multicenter cohort study. PLoS One. 2017 Nov 8;12(11):e0187165.
8. Kim, H. A., Lee, J. W., Nam, S. J., et al. Adding Ovarian Suppression to Tamoxifen for Premenopausal Breast Cancer: A Randomized Phase III Trial J Clin Oncol 2020 Feb 10;38(5):434-443

Adhärenz erhöhen durch Verhaltenstherapie/-training

1. Ream ME, Walsh EA, Jacobs JM, et al. Brief relaxation training is associated with long-term endocrine therapy adherence among women with breast cancer: post hoc analysis of a randomized controlled trial. Breast Cancer Res Treat. 2021 Nov;190(1):79-88. doi: 10.1007/s10549-021-06361-x.

Statement zur Analgesie

1. Lu YC, Chen PT, Lin MC et al., Nonsteroidal Anti-Inflammatory Drugs Reduce Second Cancer Risk in Patients With Breast Cancer: A Nationwide Population-Based Propensity Score-Matched Cohort Study in Taiwan. Front Oncol. 2021 Nov 24;11:756143. doi: 10.3389/fonc.2021.756143



Breast Cancer Follow-Up Objectives

	Oxford		
	LoE	GR	AGO
<ul style="list-style-type: none"> ▪ Psycho-social aspects of support and counseling <ul style="list-style-type: none"> ▪ Pregnancy, contraception, sexuality, quality of life, menopausal symptoms, fear of recurrence ▪ Inclusion of related persons (partner, family, friends, caregivers) 	4	C	+
<ul style="list-style-type: none"> ▪ Second opinion regarding primary therapy 	2c	B	++
<ul style="list-style-type: none"> ▪ General counseling (e.g. changes in family history of breast, ovarian, prostate, pancreas carcinoma with new indication for genetic counseling, HRT, prophylactic surgery, breast reconstruction) 	2c	C	+

Statement: Psycho-social aspects

1. Cathcart-Rake EJ, Ruddy KJ (2023) Evidence-Based Guidance for Breast Cancer Survivorship. Hematol Oncol Clin North Am 37(1):225–243. doi:10.1016/j.hoc.2022.08.019
2. Drolet M, Maunsell E, Brisson J et al. Not Working 3 Years After Breast Cancer: Predictors in a Population-Based Study. J Clin Oncol 2005; 23(33): 8305-8312.
3. Fors EA, Bertheussen GF, Thune I et al.: Psychosocial interventions as part of breast cancer rehabilitation programs? Results from a systematic review. Psycho-Oncology 2011; 20: 909-918.
4. Gudenkauf LM, Ehlers SL. Psychosocial interventions in breast cancer survivorship care. Breast. 2017 Nov 20;38:1-6.
5. Knerr S, Guo B, Wernli KJ et al. (2023) Longitudinal adherence to breast cancer surveillance following cancer genetic testing in an integrated health care system. Breast Cancer Res Treat 201(3):461–470. doi:10.1007/s10549-023-07007-w
6. Luke B, Brown MB, Missmer SA et al.. Assisted reproductive technology use and outcomes among women with a history of cancer. Hum Reprod. 2016 Jan;31(1):183-9.
7. Scheier MF, Helgeson VS, Schulz R et al.. Interventions to Enhance Physical and Psychological Functioning Among Younger Women Who Are Ending Nonhormonal Adjuvant Treatment for Early-Stage Breast Cancer. J Clin Oncol 2005; 23(19): 4298-4311.
8. Silva C, Caramelo O, Almeida-Santos T et al.. Factors associated with ovarian function recovery after chemotherapy for

breast cancer: a systematic review and meta-analysis. Hum Reprod. 2016 Dec;31(12):2737-2749.

9. Rogers LQ, Courneya KS, Anton PM et al.. Effects of a multicomponent physical activity behavior change intervention on fatigue, anxiety, and depressive symptomatology in breast cancer survivors: randomized trial. Psychooncology. 2017 Nov;26(11):1901-1906.
10. Tran TXM, Jung S, Lee EG et al., Fear of Cancer Recurrence and Its Negative Impact on Health-Related Quality of Life in Long-term Breast Cancer Survivors. Cancer Res Treat. 2021 Dec 9. doi: 10.4143/crt.2021.835.

Statement: prophylactic surgery

1. Rhiem K, Engel C, Graeser M et al.. The risk of contralateral breast cancer in patients from BRCA 1/2 negative high risk families as compared to patients from BRCA1 or BRCA2 positive families: a retrospective cohort study. Breast Cancer Res. 2012; 14(6): R156.

Breast Cancer Follow-Up Recommended Interventions

Interventions regarding lifestyle risks and comorbidity in order to reduce an unfavorable impact on disease outcome

	Oxford		
	LoE	GR	AGO
<ul style="list-style-type: none"> ▪ Treatment of type II-diabetes (> 25% undetected DM in postmenopausal BC patients, endocrine therapy improves risk for DM) 	2a	B	++
<ul style="list-style-type: none"> ▪ Weight/lifestyle intervention (if BMI < 18.5 and > 30) 	2a	B	+
<ul style="list-style-type: none"> ▪ Nightly fastening > 13 h 	2b	B	+
<ul style="list-style-type: none"> ▪ Reduction of dietary intake (at least 15 % calories from fat) in HR-negative BC is associated with improved overall survival 	2b	B	+
<ul style="list-style-type: none"> ▪ Stop smoking (smoking causes 2-fold increase in BC-specific and 4-fold increase in not directly BC-associated mortality) 	2b	B	++
<ul style="list-style-type: none"> ▪ Alcohol consumption reduction (below 6g/d) 	2b	B	+
<ul style="list-style-type: none"> ▪ Moderate sport (in patients with reduced physical activity prior to diagnosis) (at least 150 minutes/w, 2x/w) 	1b	A	++
<ul style="list-style-type: none"> ▪ Distress reduction 	3b	B	+

1. Anderson C, Sandler DP, Weinberg CR et al. Age- and treatment-related associations with health behavior change among breast cancer survivors. *Breast*. 2017 Jun;33:1-7.
2. Chlebowski RT, Aragaki AK, Anderson GL et al. Low-Fat Dietary Pattern and Breast Cancer Mortality in the Women's Health Initiative Randomized Controlled Trial. *J Clin Oncol*. 2017 Sep 1;35(25):2919-2926.
3. Chlebowski RT, Aragaki AK, Pan K, et al. Breast cancer incidence and mortality by metabolic syndrome and obesity: The Women's Health Initiative. *Cancer*. 2024 Sep 15;130(18):3147-3156. doi: 10.1002/cncr.35318. Epub 2024 May 13. PMID: 38736319.
4. D'cunha K, Park Y, Protani MM, Reeves MM (2023) Circadian rhythm disrupting behaviours and cancer outcomes in breast cancer survivors: a systematic review. *Breast Cancer Res Treat* 198(3):413–421. doi:10.1007/s10549-022-06792-0
5. Gudenkauf LM, Ehlers SL. Psychosocial interventions in breast cancer survivorship care. *Breast*. 2017 Nov 20;38:1-6. Review.
6. Mehra K, Berkowitz A, Sanft T.D et al. Physical Activity, and Body Weight in Cancer Survivorship. *Med Clin North Am*. 2017 Nov;101(6):1151-1165. Review
7. Haykowsky MJ, Scott JM, Hudson K et al. Lifestyle Interventions to Improve Cardiorespiratory Fitness and Reduce Breast Cancer Recurrence. *Am Soc Clin Oncol Educ Book*. 2017;37:57-64.
8. Marinac CR, Nelson SH, Breen CI et al. Prolonged Nightly Fasting and Breast Cancer Prognosis. *JAMA Oncol*. 2016 Aug

1;2(8):1049-55.

9. Onitilo AA, Donald M, Stankowski RV et al. Breast and prostate cancer survivors in a diabetic cohort: results from the Living with DiabetesStudy. Clin Med Res. 2013 Dec;11(4):210-8.
10. Sonnenblick A, Agbor-Tarh D, Bradbury I, et al. Impact of Diabetes, Insulin, and Metformin Use on the Outcome of Patients With Human Epidermal Growth Factor Receptor 2–Positive Primary Breast Cancer: Analysis From the ALTO Phase III Randomized Trial. Journal of Clinical Oncology 2017;35:1421-9.
11. Syrowatka A, Motulsky A, Kurteva S et al. Predictors of distress in female breast cancer survivors: a systematic review. Breast Cancer Res Treat. 2017 Sep;165(2):229-245. Review.
12. Berrino F, Villarini A, Gargano G et al. The Effect of Diet on Breast Cancer Recurrence: The DIANA-5 Randomized Trial. Clin Cancer Res. 2024 Mar 1;30(5):965-974. doi: 10.1158/1078-0432.CCR-23-1615. PMID: 37847493; PMCID: PMC10905522.

AHT erhöht Diabetes mellitus

1. Ye F, Wen J, Yang A et al., The Influence of Hormone Therapy on secondary diabetes mellitus in Breast Cancer: A Meta-analysis. Clin Breast Cancer. 2021 Jul 21:S1526-8209(21)00174-9. doi: 10.1016/j.clbc.2021.06.014.

Statement: for all statements see most recent literature see at Survivorship care guidelines of ASC and ASCO

1. Runowcz CD, Leach CR, Henry L et al. American Cancer Society/American Society of Clinical Oncology breast cancer survivorship care guideline. CA Cancer J Clin 2016; 66: 43-73.
2. Rock CL, Doyle C, Demark-Wahnefried W, et al. Nutrition and physical activity guidelines for cancer survivors. CA: a cancer journal for clinicians 2012;62:243-74.

Weight intervention.

1. Chajès V, Romieu I. Nutrition and breast cancer. Maturitas, 2014; 77 (1): 7–11.
2. Goodwin PJ, Segal RJ, Vallis M, et al. The LISA randomized trial of a weight loss intervention in postmenopausal breast cancer. npj Breast Cancer 2020;6:6.
3. Hauner D, Rack B, Friedl T, et al. Rationale and description of a lifestyle intervention programme to achieve moderate weight loss in women with non-metastatic breast cancer: the lifestyle intervention part of the SUCCESS C Study. BMJ

Nutr Prev Health. 2020 Sep 21;3(2):213-219. doi: 10.1136/bmjnph-2020-000119.

4. Roberts SA, Gillespie TC, Shui AM et al., Weight loss does not decrease risk of breast cancer-related arm lymphedema. *Cancer*. 2021 Nov 1;127(21):3939-3945. doi: 10.1002/cncr.33819.
5. Shaikh H, Bradhurst P, Ma LX et al.: Body weight management in overweight and obese breast cancer survivors. *The Cochrane database of systematic reviews* 2020;12:Cd012110.
6. Trestini I, Sperduti I, Caldart A et al., Evidence-based tailored nutrition educational intervention improves adherence to dietary guidelines, anthropometric measures and serum metabolic biomarkers in early-stage breast cancer patients: A prospective interventional study. *Breast*. 2021 Dec;60:6-14. doi: 10.1016/j.breast.2021.08.008.

Moderate sport intervention when physical activity was reduced


1. Chlebowski RT. Nutrition and physical activity influence on breast cancer incidence and outcome. *Breast* 2013; Aug;22 Suppl 2: S30-7.
2. Friedenreich CM, Stone CR, Cheung WY, et al. Physical Activity and Mortality in Cancer Survivors: A Systematic Review and Meta-Analysis. *JNCI cancer spectrum* 2020;4:pkz080.
3. Kehm RD, MacInnis RJ, John EM et al., Recreational Physical Activity and Outcomes After Breast Cancer in Women at High Familial Risk. *JNCI Cancer Spectr*. 2021 Dec 8;5(6):pkab090. doi: 10.1093/jncics/pkab090.
4. Patsou ED, Alexias GD, Anagnostopoulos FG et al.. Effects of physical activity on depressive symptoms during breast cancer survivorship: a meta-analysis of randomised control trials. *ESMO Open*. 2017 Dec 11;2(5):e000271

Nut consumption improves OS und DFS

1. Wang C, Gu K, Wang F et al., Nut consumption in association with overall mortality and recurrence/disease-specific mortality among long-term breast cancer survivors. *Int J Cancer*. 2022 Feb 15;150(4):572-579.

Bariatric surgery

1. Lee E, Kawaguchi ES, Zhang J et al. Bariatric surgery in patients with breast and endometrial cancer in California: population-based prevalence and survival. *Surg Obes Relat Dis*. 2022 Jan;18(1):42-52. doi: 10.1016/j.soard.2021.09.017.



Routine Follow-Up Examinations in Asymptomatic Patients

Tests:	Oxford		
	LoE	GR	AGO
▪ History (specific symptoms)	1a	A	++
▪ Physical examination	1a	B	++
▪ Breast self-examination	5	D	+
▪ Mammography	1a	A	++
▪ Sonography of the breast	2a	B	++
▪ Routine MRI of the breast*	3a	B	+/-
▪ Breast MRI if conventional imaging is inconclusive	3b	B	+
▪ Pelvic examination	5	D	++
▪ DXA-scan at baseline and repeat scan according to individual risk in women with premature menopause or women taking an AI	5	D	+

* Consider in case of increased risk (age < 50 y, HR-neg., diagnostic assessability C/D in mammography + ultrasound)

1. Bychkovsky BL, Lin NU. Imaging in the evaluation and follow-up of early and advanced breast cancer: When, why, and how often? *Breast*. 2017 Feb;31:318-324 Review.
2. Expert Panel on Breast Imaging: Moy L, Bailey L, D'Orsi C et al. ACR Appropriateness Criteria[®] Stage I Breast Cancer: Initial Workup and Surveillance for Local Recurrence and Distant Metastases in Asymptomatic Women. *J Am Coll Radiol*. 2017 May;14(5S):S282-S292.
3. Lam DL, Houssami N, Lee JM. Imaging Surveillance After Primary Breast Cancer Treatment. *AJR Am J Roentgenol*. 2017 Mar;208(3):676-686. Review.
4. Saltbæk L, Horsboel TA, Offersen BV, et al.. Patterns in detection of recurrence among patients treated for breast cancer. *Breast Cancer Res Treat*. 2020 Nov;184(2):365-373. doi: 10.1007/s10549-020-05847-4. Epub 2020 Aug 6. PMID: 32761415.
5. Sartor H, Hagberg O, Hemmingsson O, et al.. Breast cancer recurrence in relation to mode of detection: implications on personalized surveillance. *Breast Cancer Res Treat*. 2024 Sep 10. doi: 10.1007/s10549-024-07475-8. Epub ahead of print. PMID: 39251456.
6. Swinnen J, Keupers M, Soens J et al. Breast imaging surveillance after curative treatment for primary non-metastasised breast cancer in non-high-risk women: a systematic review. *Insights Imaging*. 2018 Dec;9(6):961-970. doi: 10.1007/s13244-018-0667-5. Epub 2018 Nov 8. PMID: 30411278; PMCID: PMC6269345.

Statement: Physical examination

1. Margenthaler JA, Allan E, Cheng L, et al. Surveillance of Patients With Breast Cancer After Curative-Intent Primary Treatment: Current Practice Patterns. *Journal of Oncology Practice* 2012; 8(2): 79 – 83.
2. Khatcheressian JL, Hurley P, Bantug E, et al. Breast Cancer Follow-up and Management After Primary Treatment: American Society of Clinical Oncology Clinical Practice Guideline Update . *J Clin Oncol*. 2013 March 1; 31(7):961-965.
3. Jochelson M, Hayes DF, Ganz PA. Surveillance and Monitoring in Breast Cancer Survivors: Maximizing Benefit and Minimizing Harm. *ASCO Educational Book* 2013 e13 – e18.
4. Parmeshwar R, Margenthaler JA, Allam E et al.. Patient Surveillance After Initial Breast cancer Therapy Variation by Physician Specialty. *Am J Surg* 2013; 206(2): 218-222.

Statement: Mammography

1. De Bock GH, Bonnema J, van Der Hage J et al. Effectiveness of Routine Visits and Routine Tests in Detecting Isolated Locoregional Recurrences After Treatment for Early-Stage Invasive Breast Cancer: A Meta-Analysis and Systematic Review. *J Clin Oncol* 2004; 22(19): 4010-4018 .
2. Elder K, Matheson J, Nickson C et al. (2023) Contrast enhanced mammography in breast cancer surveillance. *Breast Cancer Res Treat* 199(2):221–230. doi:10.1007/s10549-023-06916-0
3. Evans A, Dunn J, Donnelly PK. Mammographic surveillance after breast cancer. *Br J Radiol*. 2024 May 7;97(1157):882-885. doi: 10.1093/bjr/tqae043. PMID: 38450420; PMCID: PMC11075979.
4. Khatcheressian JL, Hurley P, Bantug E et al. Breast Cancer Follow-up and Management After Primary Treatment: American Society of Clinical Oncology Clinical Practice Guideline Update . *J Clin Oncol*. 2013 March 1; 31(7):961-965.
5. Matheson J, Elder K, Nickson C, et al. Contrast-enhanced mammography for surveillance in women with a personal history of breast cancer. *Breast Cancer Res Treat*. 2024 Nov;208(2):293-305.
6. Moschetti, I, Cinquini, M, Lambertini, M, Levaggi, A, Liberati, A, *Follow-up strategies for women treated for early breast cancer*. *Cochrane Database Syst Rev*, 2016. 2016 (5)(5): p. CD001768
7. NCCN Guidelines, Breast Cancer, 2024. Version 5.2024. <https://www.nccn.org/guidelines/guidelines-detail?category=1&id=1419>

Statement: Sonography of the breast

1. Dillon MF, Hill AD, Quinn CM et al. The accuracy of ultrasound, stereotactic, and clinical core biopsies in the diagnosis of breast cancer, with an analysis of false-negative cases. *Ann Surg.* 2005; 242(5):701-7. .
2. Jung Hyun-Yun., Min Jung Kim, Eun-Kyung Kim et al.. Imaging Surveillance of Patients with Breast Cancer after Primary Treatment: Current Recommendations. *Korean J Radiol* 2015;16(2):219-228.
3. Song SE, Cho N, Chang JM et al. Diagnostic performances of supplemental breast ultrasound screening in women with personal history of breast cancer. *Acta Radiol.* 2017 Jan 1:284185117725779.

Statement: MRI of the breast

1. Cho N, Han W, Han BK et al. Breast Cancer Screening With Mammography Plus Ultrasonography or Magnetic Resonance Imaging in Women 50 Years or Younger at Diagnosis and Treated With Breast Conservation Therapy. *JAMA Oncol.* 2017 Nov 1;3(11):1495-1502.
2. Kim EJ, Kang BJ, Kim SH et al. Diagnostic Performance of and Breast Tissue Changes at Early Breast MR Imaging Surveillance in Women after Breast Conservation Therapy. *Radiology.* 2017 Sep;284(3):656-666.
3. Lehman CD, Lee JM, DeMartini WB et al. Screening MRI in Women With a Personal History of Breast Cancer. *J Natl Cancer Inst.* 2016 Jan 7;108(3).
4. Shah C, Ahlawat S, Khan A et al. The Role of MRI in the Follow-up of Women Undergoing Breast-conserving Therapy. *Am J Clin Oncol.* 2016 Jun;39(3):314-9.
5. Tadros A, Arditi B, Weltz C et al.:Utility of surveillance MRI in women with a personal history of breast cancer. *Clin Imaging.* 2017 Nov - Dec;46:33-36.
6. Warner E. The role of magnetic resonance imaging in screening women at high risk of breast cancer. *Top Magn Reson Imaging.* 2008; 19(3):163-9. Review.
7. Yoon J, Han K, Nahm S, et al. Surveillance Breast MRI in Women with a History of Breast Cancer: Association with Occurrence of Advanced Second Breast Cancer. *Radiology.* 2025 Jan;314(1):e240119.


Statement: Pelvic examination Expert Opinion

1. Cohen I, Beyth Y, Tepper R. The role of ultrasound in the detection of endometrial pathologies in asymptomatic postmenopausal breast cancer patients with tamoxifen treatment. *Obstet Gynecol Surv* 1998; 53(7): 429-38.
2. Geurts SM, de Vegt F, Siesling S et al. Pattern of follow up care and early relapse detection in breast cancer patients. *Breast Cancer Res Treat* 2012; 136(3): 859-68.

3. Giorda G, Crivellari D, Veronesi A et al. Comparison of ultrasonography, hysteroscopy, and biopsy in the diagnosis of endometrial lesions in postmenopausal tamoxifen-treated patients. *Acta Obstet Gynecol Scand* 2002; 81(10):975-80.
4. Robertson C1, Arcot Ragupathy SK, Boachie C et al. The clinical effectiveness and cost-effectiveness of different surveillance mammography regimens after the treatment for primary breast cancer: systemic reviews registry database analyses and economic evaluation. *Health Technol Assess*. 2011;15(34): 1-322.

Statement: DEXA scan Expert Opinion

1. Mahon SM, Williams MT, Spies MA: Screening for second cancers and osteoporosis in long-term survivors. *Cancer Pract* 2000; 8(6): 282-90.
2. Runowcz CD, Leach CR, Henry L et al. American Cancer Society/American Society of Clinical Oncology breast cancer survivorship care guideline. *CA Cancer J Clin* 2016; 66: 43-73.
3. Shapiro CL, Van Poznak C, Lacchetti C, et al. Management of Osteoporosis in Survivors of Adult Cancers With Nonmetastatic Disease: ASCO Clinical Practice Guideline. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2019;37:2916-46.



Routine Follow-Up Examinations in Asymptomatic Patients

	Oxford		
	LoE	GR	AGO
▪ Routine biochemistry (incl. tumor markers)	1a	A	-
▪ Blood tests for monitoring of acute and late toxicities	5	D	+
▪ Ultrasound of the liver / Bone scan / Chest X-ray	1a	A	-
▪ CT of chest, abdomen, and pelvis	2a	D	-
▪ Detection of isolated / circulating tumor cells*	2a	D	-
▪ ctDNA*	2a	D	-
▪ PET / Whole body MRI	2b	B	-

* Study participation recommended

1. Bychkovsky BL, Lin NU. Imaging in the evaluation and follow-up of early and advanced breast cancer: When, why, and how often? *Breast*. 2017 Feb;31:318-324 Review.
2. Expert Panel on Breast Imaging; Moy L, Bailey L, D'Orsi C, Green ED et al. ACR Appropriateness Criteria[®] Stage I Breast Cancer: Initial Workup and Surveillance for Local Recurrence and Distant Metastases in Asymptomatic Women. *J Am Coll Radiol*. 2017 May;14(5S):S282-S292.
3. Lam DL, Houssami N, Lee JM. Imaging Surveillance After Primary Breast Cancer Treatment. *AJR Am J Roentgenol*. 2017 Mar;208(3):676-686. Review.
4. Lafranconi A, Pylkkänen L, Deandrea S et al. Intensive follow-up for women with breast cancer: review of clinical, economic and patient's preference domains through evidence to decision framework. *Health Qual Life Outcomes*. 2017 Oct 19;15(1):206.
5. Mullins MA, Atluri N, Abrahamse P et al. (2023) Primary care provider attitudes about and tendency to use non-recommended surveillance tests after curative breast cancer treatment. *Breast Cancer Res Treat* 200(3):391–398. doi:10.1007/s10549-023-06994-0
6. Sheng JY, Snyder CF, Smith KC et al. (2023) Evaluating potential overuse of surveillance care in cancer survivors. *Cancer Med* 12(5):6139–6147. doi:10.1002/cam4.5346

Statement: Routine biochemistry (incl. tumor markers)

1. Cardoso F, Kyriakides S, Ohno S, et al. Early breast cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Annals of Oncology* 2019;30:1194-220.
2. McShane LM, Altman DG, Sauerbrei W et al. Statistics Subcommittee of the NCI-EORTC Working Group on Cancer Diagnostics. Reporting recommendations for tumor marker prognostic studies. *J Clin Oncol* 2005; 23(36): 9067-72.

Statement: Ultrasound of the liver

Statement: Bone scan

Statement: Chest X-ray

Statement: CT of chest, abdomen and pelvis

1. Dull B, Linkugel A, Margenthaler JA, Cyr AE. Overuse of Chest CT in Patients With Stage I and II Breast Cancer: An Opportunity to Increase Guidelines Compliance at an NCCN Member Institution. *J Natl Compr Canc Netw*. 2017 Jun;15(6):783-789.
2. Emens LA, Davidson NE. The follow-up of breast cancer. *Semin Oncol*. 2003; 30(3): 338-48. Review.
3. Moschetti I, Cinquini M, Lambertini M et al. Follow-up strategies for women treated for early breast cancer. *Cochrane Database Syst Rev* 2005 Jan 25;(1) Review. Update in: *Cochrane Database Syst Rev* 2016; (5) CD001768.

Statement: Detection of isolated/circulating tumor cells/ctDNA

1. Janni W, Vogl FD, Wiedswang G et al. Persistence of disseminated tumor cells (DTC) in bone marrow (BM) during Follow-up predicts increased risk for relapse – Up-date of the pooled European data. *Clin Cancer Res* 2011; 17(9): 2967-76.
2. Rack B, Schindlbeck C, Jüeckstock J et al. Circulating tumor cells predict survival in early average-to-high risk breast cancer patients. SUCCESS Study Group. *J Natl Cancer Inst*. 2014 May 15;106(5).
3. Turner NC, Swift C, Jenkins B et al. (2023) Results of the c-TRAK TN trial: a clinical trial utilising ctDNA mutation tracking to detect molecular residual disease and trigger intervention in patients with moderate- and high-risk early-stage triple-negative breast cancer. *Ann Oncol* 34(2):200–211. doi:10.1016/j.annonc.2022.11.005
4. Turner N. et al. GS3-01: Circulating tumor DNA surveillance in ZEST, a randomized, phase 3, double-blind study of niraparib or placebo in patients w/ triple-negative breast cancer or HER2+ BRCA-mutated breast cancer with molecular residual disease after definitive therapy SABCS 2024. Abstract Number: SESS-2027

Statement: PET / WB-MRI

1. Cho IH, Kong EJ. Potential Clinical Applications of ¹⁸F-Fluorodeoxyglucose Positron Emission Tomography/Magnetic Resonance Mammography in Breast Cancer. Nucl Med Mol Imaging. 2017 Sep;51(3):217-226. Review.
2. Ide M. Cancer screening with FDG-PET. Q J Nucl Med Mol Imaging 2006; 50(1): 23-7.
3. Lei L, Wang X, Chen Z. PET/CT Imaging for Monitoring Recurrence and Evaluating Response to Treatment in Breast Cancer. Adv Clin Exp Med. 2016 Mar-Apr;25(2):377-82.
4. Lee H, Choi JY, Park YH et al. (2023) Diagnostic Value of FDG PET/CT in Surveillance after Curative Resection of Breast Cancer. Cancers (Basel) 15(9). doi:10.3390/cancers15092646
5. Melsaether A, Moy L. Breast PET/MR Imaging. Radiol Clin North Am. 2017 May;55(3):579-589. Review.
6. Schöder H, Gönen M. Screening for cancer with PET and PET/CT: potential and limitations. J Nucl Med 2007; 48 Suppl 1:4S-18S. Review.



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.
Guidelines Breast
Version 2025.1E

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

Background for Toxicity Management

Tamoxifen:	Cholesterol, Triglycerides, Bilirubin, ALAT, ASAT, gamma-GT, Glucose
Aromatase inhibitors:	Cholesterol, Triglycerides, Bilirubin, ALAT, ASAT, gamma-GT
Anthracyclines:	pro-BNP, possibly Troponin
Trastuzumab:	pro-BNP, possibly Troponin
Checkpoint inhibitors:	Bilirubin, ALAT, ASAT, gamma-GT, Creatinine, TSH, fT3/T4, Myoglobin

Early Detection of Potentially Curable Events

	Oxford		
	LoE	GR	AGO
Locoregional recurrence (chest wall, in-breast):			
▪ Incidence 7-20% (depending on time of F/U)			
▪ Breast self-examination	5	D	+
▪ Physical examination, mammography & US	1a	A	++
▪ Magnetic resonance imaging (MRI)*	3a	B	+/-

* Consider in case of increased risk (age < 50 y, HR-neg., diagnostic assessability C/D in mammography + ultrasound)

Statement incidence

1. Cheun J-H, Kim H-K, Moon H-G, Han W, Lee H-B (2023) Locoregional Recurrence Patterns in Patients With Different Molecular Subtypes of Breast Cancer. JAMA Surg 158(8):841–852. doi:10.1001/jamasurg.2023.2150
2. Perry NM. Quality assurance in the diagnosis of breast disease. EUSOMA Working Party. Eur J Cancer 2001; 37: 159-172
3. Wapnir IL, Anderson SJ, Mamounas EP et al. Prognosis after ipsilateral breast tumor recurrence and locoregional recurrences in five National Surgical Adjuvant Breast and Bowel Project node-positive adjuvant breast cancer trials. J Clin Oncol 2006; 24:2028-2037

Statement breast self examination

1. Khatcheressian JL, Wolff AC, Smith TJ. American Society of Clinical Oncology 2006 update of the breast cancer follow-up and management guidelines in the adjuvant setting. J Clin Oncol. 2006 Nov 1;24(31):5091-7.
2. Montgomery DA, Krupa K, Cooke TG. Follow-up in breast cancer: does routine clinical examination improve outcome? A systematic review of the literature. Br J Cancer 2007; 97(12): 1632-41.
3. Thomas DB, Gao DL, Ray RM et al. Randomized trial of breast self-examination in Shanghai: final results. J Natl Cancer Inst 2002; 94(19): 1445-57.

Statement physical examination, mammography & US & MRI

1. Beinart G, Gonzalez-Angulo AM, Broglio K. Clinical course of 771 patients with bilateral breast cancer: characteristics associated with overall and recurrence-free survival. *Clin Breast Cancer* 2007; 7(11): 867-74.
2. Buist DS1, Abraham LA, Barlow WE et al. Diagnosis of second breast cancer events after initial diagnosis of early stage breast cancer. Breast Cancer Surveillance Consortium. *Breast Cancer Res Treat* 2010; 124(3): 863-73.
3. Kim JY, Cho N, Koo HR et al. Unilateral breast cancer: screening of contralateral breast by using preoperative MR imaging reduces incidence of metachronous cancer. *adiology*. 2013 Apr;267(1):57-66.
4. Khatcheressian JL, Hurley P, Bantug E et al. Breast Cancer Follow-up and Management After Primary Treatment: American Society of Clinical Oncology Clinical Practice Guideline Update . *J Clin Oncol*. 2013 March 1; 31(7):961-965.
5. Montgomery DA, Krupa K, Cooke TG. Follow-up in breast cancer: does routine clinical examination improve outcome? A systematic review of the literature. *Br J Cancer*. 2007; 97(12): 1632-41.
6. Neuman HB, Schumacher JR, Francescatti AB et al. Utility of Clinical Breast Examinations in Detecting Local-Regional Breast Events After Breast-Conservation in Women with a Personal History of High-Risk Breast Cancer. *Ann Surg Oncol*. 2016 Oct;23(10):3385-91.
7. Tsai WC, Wei HK, Hung CF et al. Better Overall Survival for Breast Cancer Patients by Adding Breast Ultrasound to Follow-Up Examinations for Early Detection of Locoregional Recurrence-A Survival Impact Study. *Ultrasound Med Biol*. 2016 Sep;42(9):2058-64.
8. Taros A, Arditi B, Weltz C et al. Utility of surveillance MRI in women with a personal history of breast cancer. *Clin Imaging*. 2017 Nov - Dec;46:33-36.

Early Detection of Potentially Curable Events

	Oxford		
	LoE	GR	AGO
Contralateral breast cancer:			
▪ Relative risk: 2.5-5			
▪ Incidence: 0.5-1.0 %/year			
▪ Breast self-examination	5	D	+
▪ Physical examination, mammography & US	1a	A	++
▪ Routine breast MRI*	3b	B	+/-
▪ Male breast cancer: analogous to BC in women**	5	D	+

* Consider in case of increased risk: age < 50 y, HR-neg., diagnostic assessability C/D in mammography + ultrasound.

** See chapter "Breast Cancer Specific Situations"

Statement risk and incidence

1. Bertelsen L, Mellekjær L, Christensen J et al. Age-Specific Incidence of Breast Cancer in Breast Cancer Survivors and Their First-Degree Relatives. *Epidemiology* 2009; 20(2): 175 – 80.
2. Chao C, Bhatia S, Xu L, et al. Incidence, Risk Factors, and Mortality Associated With Second Malignant Neoplasms Among Survivors of Adolescent and Young Adult Cancer. *JAMA Network Open* 2019;2:e195536-e.
3. Ramin C, Veiga LHS, Vo JB et al. (2023) Risk of second primary cancer among women in the Kaiser Permanente Breast Cancer Survivors Cohort. *Breast Cancer Res* 25(1):50. doi:10.1186/s13058-023-01647-y
4. Yerushalmi R, Kennecke H, Woods R et al. Does multicentric/multifocal breast cancer differ from unifocal breast cancer? An analysis of survival and contralateral breast cancer incidence. *Breast Cancer Res Treat* 2009; 117(2): 365-70.

Statement breast self examination

1. Khatcheressian JL, Hurley P, Bantug E et al.: Breast Cancer Follow-up and Management After Primary Treatment: American Society of Clinical Oncology Clinical Practice Guideline Update. *J Clin Oncol.* 2013 March 1; 31(7):961-965.
2. Thomas DB, Gao DL, Ray RM et al. Randomized trial of breast self-examination in Shanghai: final results. *J Natl Cancer Inst* 2002; 94(19): 1445-57.
3. Montgomery DA, Krupa K, Cooke TG et al. Follow-up in breast cancer: does routine clinical examination improve

outcome? A systematic review of the literature. Br J Cancer 2007; 97(12): 1632-41.

Statement physical examination, mammography & US&MRI

1. Freedman RA, Keating NL, Partridge AH et al. Surveillance Mammography in Older Patients With Breast Cancer-Can We Ever Stop?: A Review. JAMA Oncol. 2017 Mar 1;3(3):402-409.
2. Vapiwala N, Hwang WT, Kushner CJ et al. No impact of breast magnetic resonance imaging on 15-year outcomes in patients with ductal carcinoma in situ or early-stage invasive breast cancer managed with breast conservation therapy. Cancer. 2017 Apr 15;123(8):1324-1332.
3. van Bodegraven EA, van Raaij JC, Van Goethem M et al. Guidelines and recommendations for MRI in breast cancer follow-up: A review. Eur J Obstet Gynecol Reprod Biol. 2017 Nov;218:5-11.

Statement surveillance of male breast cancer

1. Auvinen A, Curtis RE, Ron E. Risk of subsequent cancer following breast cancer in men. J Natl Cancer Inst. 2002;94(17):1330–1332.
2. Ferzoco RM, Ruddy KJ. Optimal delivery of male breast cancer follow-up care: improving outcomes, Breast Cancer: Targets and Therapy 2015;7 371–379

Early Detection of Potentially Curable Events

	Oxford		
	LoE	GR	AGO
Unrelated site carcinoma:			
▪ MDS (RR 10.9), AML (RR 2.6-5.3), Colon RR 3.0; endometrium RR 1.6; ovary RR 1.5; lymphoma RR 7			
▪ Screening for secondary malignancies according to current guidelines	5	D	++
▪ Pelvic examination and PAP smear	5	D	++
▪ Routine endometrial ultrasound / biopsy	1b	B	-

Statement: Risk

1. Allen I, Hassan H, Sofianopoulou E, Eccles D et al. (2023) Risks of second non-breast primaries following breast cancer in women: a systematic review and meta-analysis. *Breast Cancer Res* 25(1):18. doi:10.1186/s13058-023-01610-x
2. Freedman RA, Seisler DK, Foster JC et al. Risk of acute myeloid leukemia and myelodysplastic syndrome among older women receiving anthracycline-based adjuvant chemotherapy for breast cancer on Modern Cooperative Group Trials (Alliance A151511). *Breast Cancer Res Treat.* 2017 Jan;161(2):363-373. doi: 10.1007/s10549-016-4051-1.
3. Kim H, Kim SS, Lee JS et al. (2023) Epidemiology of Second Non-breast Primary Cancers among Survivors of Breast Cancer: A Korean Population-Based Study by the SMARTSHIP Group. *Cancer Res Treat* 55(2):580–591. doi:10.4143/crt.2022.410
4. Ramin C, Veiga LHS, Vo JB et al. (2023) Risk of second primary cancer among women in the Kaiser Permanente Breast Cancer Survivors Cohort. *Breast Cancer Res* 25(1):50. doi:10.1186/s13058-023-01647-y
5. Sung H, Hyun N, Leach CR, et al. Association of First Primary Cancer With Risk of Subsequent Primary Cancer Among Survivors of Adult-Onset Cancers in the United States. *JAMA* 2020;324:2521-35.

Statement: Marrow neoplasms after adjuvant breast cancer therapy

1. Wolff AC, Blackford AL, Visvanathan K et al. Risk of marrow neoplasms after adjuvant breast cancer therapy: the national

comprehensive cancer network experience. J Clin Oncol. 2015; 33(4): 340-8.

Statement: Secondary lung tumors:

1. Burt LM, Ying J, Poppe MM, et al. Risk of secondary malignancies after radiation therapy for breast cancer: Comprehensive results. Breast (Edinburgh, Scotland) 2017;35:122-9.
2. Hoekstra N, Fleury E, Merino Lara TR et al. Long-term risks of secondary cancer for various whole and partial breast irradiation techniques. Radiother Oncol. 2018 Sep;128(3):428-433

Statement: Screening for secondary malignancies according to current guidelines

1. Allen, I, Hassan, H, Sofianopoulou, E, Eccles, D, Turnbull, C, Tischkowitz, M, et.al. Risks of second non-breast primaries following breast cancer in women: a systematic review and meta-analysis.. Breast Cancer Res, 2023. 25 (1)(1): p. 18
2. Runowicz CD, Leach CR, Henry NL, et al. American Cancer Society/American Society of Clinical Oncology Breast Cancer Survivorship Care Guideline. Journal of Clinical Oncology 2016;34:611-35.

Statement: Pelvic examination and PAP smear

1. Gerber B, Krause A, Müller H et al. Ultrasonographic detection of asymptomatic endometrial cancer in postmenopausal patients offers no prognostic advantage over symptomatic disease discovered by uterine bleeding. Eur J Cancer 2001; 37(1): 64-71.
2. Chan JK, Manuel MR, Cheung MK et al. Breast cancer followed by corpus cancer: is there a higher risk for aggressive histologic subtypes? Gynecol Oncol 2006; 102(3): 508-12.

Statement: Endometrial ultrasound / biopsy

1. Barakat RR, Gilewski TA, Almadrones L et al. Effect of adjuvant tamoxifen on the endometrium in women with breast cancer: a prospective study using office endometrial biopsy. J Clin Oncol 2000;18(20): 3459-63.
2. Fung MF, Reid A, Faught W et al. Prospective longitudinal study of ultrasound screening for endometrial abnormalities in women with breast cancer receiving tamoxifen. Gynecol Oncol 2003; 91(1): 154-9.
3. Gerber B, Krause A, Müller H, et al. Effects of adjuvant tamoxifen on the endometrium in postmenopausal women with breast cancer: a prospective long-term study using transvaginal ultrasound. J Clin Oncol 2000; 18(20): 3464-70

Follow-Up Care for invasive / non-invasive Breast Cancer

Recommendations for asymptomatic pts.

(mod. according to ASCO-ACS recommendations 2016, NCCN 2021, ESMO 2019 and S3-guidelines 2017)

Clinical follow-up		Follow-up*				Screening/ Follow-up	
Years after primary therapy		1	2	3	4	5	> 5
History, physical examination, counseling		every 3 months DCIS every 6 months			every 6 months		inv.: every 12 months
Self-examination		monthly					
Imaging modalities and biochemistry		indicated only if complaints, clinical findings, or suspicion of recurrence Monitoring of side effects of therapy					
Mammo-graphy and additional sonography	BCT**	both sides: every 12 months					
	Mastectomy	contralateral every 12 months					
Echocardiography		6,12,24 months and yearly up to 5 years after completion of cardiotoxic therapy, after 5th year, every 5 years and if patient is symptomatic.					

* Continued follow-up visits if still on adjuvant treatment.

** In pts after breast-conserving therapy (BCT): First mammography 1 year after initial mammography or at least 6 months after completion of radiotherapy.

1. Cardoso F, Kyriakides S, Ohno S, et al. Early breast cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Annals of Oncology* 2019;30:1194-220.
2. Curigliano G, Lenihan D, Fradley M, et al. Management of cardiac disease in cancer patients throughout oncological treatment: ESMO consensus recommendations. *Annals of Oncology* 2020;31:171-90.
3. Dent S, Fergusson D, Aseyev O et al. ,A Randomized Trial Comparing 3- versus 4-Monthly Cardiac Monitoring in Patients Receiving Trastuzumab-Based Chemotherapy for Early Breast Cancer. *Curr Oncol.* 2021 Dec 3;28(6):5073-5083.
4. Evans A, Dunn J, Donnelly PK. Mammographic surveillance after breast cancer. *Br J Radiol.* 2024 May 7;97(1157):882-885. doi: 10.1093/bjr/tqae043. PMID: 38450420; PMCID: PMC11075979.
5. Interdisziplinäre S3-Leitlinie für die Früherkennung, Diagnostik, Therapie und Nachsorge des Mammakarzinoms. Langversion 4.0 – Dezember 2017 AWMF-Registernummer: 032-045OL; http://www.leitlinienprogramm-onkologie.de/fileadmin/user_upload/LL_Mammakarzinom_Langversion_4.0.pdf
6. Merkow RP, Korenstein D, Yeahia R et al. Quality of Cancer Surveillance Clinical Practice Guidelines: Specificity and Consistency of Recommendations. *JAMA Intern Med.* 2017 May 1;177(5):701-709.
7. Muradali D, Kennedy EB, Eisen A et al. Breast screening for survivors of breast cancer: A systematic review. *Prev Med.* 2017 Oct;103:70-75.
8. Michel L, Rassaf T, Totzeck M. Biomarkers for the detection of apparent and subclinical cancer therapy-related

cardiotoxicity. J Thorac Dis 2018;10:S4282-S95.

9. NCCN Clinical Practice Guidelines in Oncology, Breast Cancer Version 02.2022
https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf
10. Rassaf T, Totzeck M, Backs J, et al. Onco-Cardiology: Consensus Paper of the German Cardiac Society, the German Society for Pediatric Cardiology and Congenital Heart Defects and the German Society for Hematology and Medical Oncology. Clin Res Cardiol 2020;109:1197-222.
11. Runowcz CD, Leach CR, Henry L et al. American Cancer Society/American Society of Clinical Oncology breast cancer survivorship care guideline. CA Cancer J Clin 2016; 66: 43-73

Breast Cancer Follow-up Duration and Breast Nurses

	Oxford		
	LoE	GR	AGO
<ul style="list-style-type: none"> ▪ Duration of follow-up <ul style="list-style-type: none"> ▪ up to 5 years ▪ up to 10 years ▪ Surveillance by specialized breast nurses 	1c	A	++
	1c	A	+
	2b	B	+/-*

* Study participation recommended

1. Freedman RA, Keating NL, Partridge AH et al.: Surveillance Mammography in Older Patients With Breast Cancer-Can We Ever Stop?: A Review. JAMA Oncol. 2017 Mar 1;3(3):402-409.
2. Sheppard C. Breast cancer follow-up: literature review and discussion. Eur J Oncol Nurs 2007;11(4):340-7.
3. Interdisziplinäre S3-Leitlinie für die Früherkennung, Diagnostik, Therapie und Nachsorge des Mammakarzinoms. Langversion 4.0 – Dezember 2017 AWMF-Registernummer: 032-045OL; http://www.leitlinienprogramm-onkologie.de/fileadmin/user_upload/LL_Mammakarzinom_Langversion_4.0.pdf
4. NCCN Clinical Practice Guidelines in Oncology, Breast Cancer Version 3.17-10.17; https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf